

Immunogenicity and Safety of Sinopharm Covid-19 Vaccine in Young Mice

Emhemmed Ali Elgallal¹, Inass A Sadawe², Abdelmonam Hadi Elghobassa¹, Abdulathim A A Alshoushan¹, Salah M Bensaber³, Jamal Ali M. Elbakay³, Nisreen H Meiqal³, Massaud Salem Maamar⁴, Mohammed A. Gbaj⁵, Anton Hermann⁶ and Abdul M Gbaj^{3*}

¹Food and Drug Control Centre (LFDA), Tripoli, Libya.

²Medical Supply Organization (MSO), Tripoli, Libya.

³Department of Medicinal Chemistry, Faculty of Pharmacy, University of Tripoli, Libya.

⁴Zoology Department, Faculty of Science, Tripoli University, Libya.

⁵Chemical Engineering Department, Faculty of Engineering, Tripoli University, Libya.

⁶Department of Biosciences, University of Salzburg, Salzburg, Austria.

***Corresponding Author:** Abdul M Gbaj, Professor of Genetics and Biochemistry, Department of Medicinal Chemistry, Faculty of Pharmacy, University of Tripoli, Libya.

Received Date: 15 October 2021 | **Accepted Date:** 10 January 2022 | **Published Date:** 17 January 2022

Citation: Nisreen H Meiqal, Massaud S Maamar, Mohammed A Gbaj, A Hermann, Abdul M Gbaj, et al. (2022). Immunogenicity and Safety of Sinopharm Covid-19 Vaccine in Young Mice. *International Journal of Clinical Case Reports and Reviews*. 10(3); DOI:10.31579/2690-4861/185

Copyright: © 2022 Abdul M Gbaj, This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background and Aim: Vaccines to prevent SARS-CoV-2 infection may be considered a promising way for reduction of the pandemic. Many different vaccines have become obtainable for use in many countries. The present study aims to evaluate the immune response and the safety of Sinopharm COVID-19 vaccine on 14 days old mice.

Materials and Methods: Our experimental study was performed on two weeks old mice, selected by random allocation. The mice were divided into three groups of 12. Group one received a single dose of 0.5 ml Sinopharm COVID-19 vaccine, group two received two doses of 0.5 ml Sinopharm COVID-19 vaccine, and group three (control) received two doses of 0.5 ml of 0.9 % NaCl.

Results: Our study shows that Sinopharm COVID-19 vaccine is safe and induces good immunity in young mice.

Conclusions: The Sinopharm COVID-19 vaccine was safe and immunogenic in 14 days old mice. The two doses of Sinopharm COVID-19 vaccine elicit a safe antibody response in young mice. Further post-marketing toxicity studies are required to assess potential hazards for children to evaluate the histopathological characteristics.

Keywords: sinopharm covid-19 vaccine; mice; coronavirus disease; immunogenic; safety

Introduction

The severe acute respiratory syndrome caused by coronavirus 2 (SARS-CoV-2), also called COVID-19, has quickly spread over the whole world and raised severe public health distresses. The scientific society is intensely requested investigating treatments that would potentially be effective in fighting COVID-19 [1, 2]. During July, 2021 WHO revealed that the COVID-19 pandemic caused more than four million deaths worldwide. Vaccination has been established to limit the further spread of SARS-CoV-2 virus. Children are also vulnerable to SARS-CoV-2 infection, although they display milder clinical symptoms of the disease [3]. This susceptibility raises the possibility of transmission between

family members and risk to elderly members who are more prone to the disease [4].

There are four categories of vaccines in use: whole virus, protein subunits, viral vectors and nucleic acid (RNA and DNA). There are additional vaccine candidates currently in the pipeline for COVID-19. All vaccines are trying to attain immunity to the virus, and some may be capable to stop transmission. By finding a proper molecule on the virus they are initiating an immune response to the antigen. In the case of COVID-19 the antigen is usually a characteristic spike protein found on the surface of the virus which assist attacks of human cells [5-7]. In case of using the entire virus (such as Sinovac and Sinopharm) it produces an immune response with the help of antigen presenting cells (APCs) such as

dendritic cells (DCs) [8]. In particular, DCs have essential functions in capturing molecules, fragmenting them into smaller peptides and presenting the antigenic peptides on their major histocompatibility complex (MHC) I and II to prime T cells for the start of cellular and humoral immunity against the virus [9]. The study aims to evaluate the immune response of Sinopharm COVID-19 vaccine and its safety in young mice aged two weeks.

Materials and Methods

Experimental animals

Young Swiss Albino male mice (10 ± 2 g) with 14 days old were used for experiments. In order to reduce the contact caused by environmental alterations and handling during behavioral studies, mice were acclimatized to the Laboratory Animal Holding Center and laboratory surroundings for three days and at least one hour before experimentation, respectively. Mice were kept under standard conditions with food (low protein diet) and water available *ad libitum*. The animals were housed six per cage in a light-controlled room (12 h light/dark cycle, light on 07:00 h) at 27°C and 65% relative humidity. All experiments were carried out between 09:30 and 15:00 h. Each test group consisted of 12 mice, and each mouse was used only once. All animal experiments were conducted according to guidelines set by the Institutional Animal Ethics Committee of the National Medical Research Centre (NMRC35/2009).

Clinical and necropsy observations

This study represents one constituent of the safety evaluation program for using Sinopharm COVID-19 vaccine for very young mice to assess efficacy and toxicity. The aim was to evaluate these parameters following the administration of the proposed human vaccine dose. The mice were divided into three groups of 12. Group one received a single dose of 0.5 ml Sinopharm COVID-19 vaccine, Group two received two doses of 0.5 ml Sinopharm COVID-19 vaccine and the second dose was given after 21 days, and group three (control) received two doses of 0.5 ml of 0.9 % NaCl. Mice were examined every day for 40 days. Any signs of ill health were recorded daily. Blood samples for IgM and IgG were taken from animals on day 14 and day 30 after first vaccine application. At necropsy a full macroscopic examination was performed on each animal. Organs macroscopically examined were the spleen, lungs, liver, kidney, heart, brain, testes, and ovaries.

Statistical analysis

The difference among various treated groups and the control groups were analyzed using one-way-ANOVA followed by using unpaired Student's t-test. The results were expressed as the mean \pm SEM of the number of experiments, with $p < 0.05$ indicating a significant difference between groups. All p values reported are for a one-tailed test. The significance level was chosen at $\alpha = 0.05$.

Results and discussion

Mice have been the most generally used animals in scientific research [10, 11]. This is could be attributed to the fact that the mouse genome is 99% identical to the human genome, and mice have similar patterns with respect to human organs and systemic physiology.

The Sinopharm Beijing Covid-19 vaccine is produced by Beijing Institute of Biological Products (BIBP), subsidiary of China National Biotec Group (CNBG), they uses inactive or weakened virus (19nCoV-CDC-Tan-HB02) strain as antigen which based on a form of the virus that has been inactivated or weakened so that it does not cause disease, but is still able to produce an immune response. It has been reported that the effectiveness of the vaccine is approximately 87.5% for the prevention of hospitalizations of Covid-19 patients, 65.9% for prevention of Covid-19, 90.3% for the prevention of intensive care unit admissions, and 86.3% for

the prevention of Covid-19-related deaths [12]. Furthermore, children younger than 12 years old are at their crucial phase of growth and development; concern must be taken to assess the long-term effect of COVID-19 vaccine on their growth and development. In addition, children whom are going to be vaccinated should have enough immunity and safety against COVID-19 vaccine [13].

It has been reported that Pfizer and Moderna messenger RNA (mRNA) vaccines studied in children older than 12 years and were found safe and effective. In addition, Pfizer and Moderna vaccines were also tested in children under 12, with the aim of involving babies from just six months old. Although teenagers only seldom get badly sick with Covid-19, they are able to spread the infection. Hence, vaccination will be able to assist stopping the pandemic [14]. Sinopharm's institute in Wuhan approved for emergency utilization on children aged between three and seventeen by the China National Biotec Group on August, 2021. China began to permit people aged between three and seventeen to obtain a dose of COVID-19 vaccines in early June 2021, making it the first country to declare the endorsement of vaccines for such a young age group [15]. None of the mice used in the study showed any sign of abnormality or ill health throughout the 42 days post-immunization observation for the three groups after the first dose of immunization. At necropsies no macroscopic treatment related changes were observed. Antibody binding the SARS-CoV-2 spike protein was induced by vaccination, and, as expected, the temporal induction of anti-spike IgM was faster than that of IgG. The mice injected with Sinopharm COVID-19 vaccine or 0.9% sodium chloride solutions were generally in good condition, no obvious clinical unusual symptoms were observed, and no death occurred during the observation period. The mice body weights in control and vaccine groups increased but the increase was more with the vaccinated group by around 1.5 times compared with the initial weights. There were significant differences in body weights and food intakes noted between the vaccine and negative control groups throughout the study period. Furthermore, no abnormal changes were found in the gross autopsy results of all mice investigated.

Conclusion

Our study shows that the Sinopharm COVID-19 vaccine given to 14 days old mice produces an immune response with no side effects ascertain its safety and protection efficacy against COVID-19. We highly recommend post-marketing surveillance of the vaccine safety when given for children for a longer period than that in adults.

References

1. Dhama K, Khan S, Tiwari R, Sircar S, Bhat S, Malik YS, Singh KP, Chaicumpa W, Bonilla-Aldana DK, Rodriguez-Morales AJ. (2020). Coronavirus Disease 2019 COVID-19. *Clin Microbiol Rev.* 33(4).
2. Abd El-Aziz TM, Stockand JD. (2020). Recent progress and challenges in drug development against COVID-19 coronavirus (SARS-CoV-2) - an update on the status. *Infect Genet Evol.* 83:104327.
3. Qiu H, Wu J, Hong L, Luo Y, Song Q, Chen D. (2020). Clinical and epidemiological features of 36 children with coronavirus disease 2019 (COVID-19) in Zhejiang, China: an observational cohort study. *Lancet Infect Dis.* 20(6):689-696.
4. Weiss P, Murdoch DR. (2020). Clinical course and mortality risk of severe COVID-19. *Lancet.* 395(10229):1014-1015.
5. Koirala A, Joo YJ, Khatami A, Chiu C, Britton PN. (2020). Vaccines for COVID-19: The current state of play. *Paediatr Respir Rev.* 35:43-49.
6. Wang J, Peng Y, Xu H, Cui Z, Williams RO, III. (2020). The COVID-19 Vaccine Race: Challenges and Opportunities in Vaccine Formulation. *AAPS Pharm SciTech.* 21(6):225.

7. Chowdhury MA, Hossain N, Kashem MA, Shahid MA, Alam A. (2020). Immune response in COVID-19: A review. *J Infect Public Health*. 13(11):1619-1629.
8. Li W, Joshi MD, Singhania S, Ramsey KH, Murthy AK. Peptide Vaccine: Progress and Challenges.
9. Rock KL, Reits E, Neefjes J. (2016). Present Yourself! By MHC Class I and MHC Class II Molecules. *Trends Immunol*. 37(11):724-737.
10. Justice MJ, Dhillon P. (2016). Using the mouse to model human disease: increasing validity and reproducibility. *Dis Model Mech*. 9(2):101-103.
11. Wang S, Lai X, Deng Y, Song Y. (2020). Correlation between mouse age and human age in anti-tumor research: Significance and method establishment. *Life Sci*. 242:117242.
12. Jara A, Undurraga EA, Gonzalez C, Paredes F, Fontecilla T, Jara G, Pizarro A, Acevedo J, Leo K, Leon F, Sans C, Leighton P, Suñeriz P, Garca-a-Escorza H, Araos R. (2021). Effectiveness of an Inactivated SARS-CoV-2 Vaccine in Chile. *E Med*. 385(10): 875-884.
13. Zou X, Cao B. (2021). COVID-19 vaccines for children younger than 12 years: are we ready? *Lancet Infect Dis*.
14. Wodi AP, Ault K, Hunter P, McNally V, Szilagyi PG, Bernstein H. (2021). Advisory Committee on Immunization Practices Recommended Immunization Schedule for Children and Adolescents Aged 18 Years or Younger-United States. *MMWR Morb Mortal Wkly Rep*. 70(6):189-192.
15. Xia S, Zhang Y, Wang Y, Wang H, Yang Y, Gao GF, Tan W, Wu G, Xu M, Lou Z, Huang W, Xu W, Huang B, Wang W, Zhang W, Li N, Xie Z, Zhu X, Ding L, You W, Zhao Y, Zhao J, Huang L, Shi X, Yang Y, Xu G, Wang W, Liu P, Ma M, Qiao Y, Zhao S, Chai J, Li Q, Fu H, Xu Y, Zheng X, Guo W, Yang X. (2021). Safety and immunogenicity of an inactivated COVID-19 vaccine, BBIBP-CorV, in people younger than 18 years: a randomised, double-blind, controlled, phase 1/2 trial. *Lancet Infect Dis*.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

Submit Manuscript

DOI: [10.31579/2690-4861/185](https://doi.org/10.31579/2690-4861/185)

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/international-journal-of-clinical-case-reports-and-reviews>