

The Case Number 112 of Congenital Chylous Ascites and Its Successful Treatment

Aamir Jalal Al-Mosawi*

Advisor in Pediatrics and Pediatric Psychiatry Children Teaching Hospital of Baghdad Medical City and the National Training and Development Center

***Corresponding Author:** Aamir Jalal Al-Mosawi, Advisor in Pediatrics and Pediatric Psychiatry Children Teaching Hospital of Baghdad Medical City and the National Training and Development Center.

Received Date: March 30, 2022; **Accepted Date:** April 15, 2022; **Published Date:** April 30, 2022.

Citation: Aamir Jalal Al-Mosawi, (2022) The Case Number 112 of Congenital Chylous Ascites and Its Successful Treatment Aamir Jalal Al-Mosawi. *Biomedical Research and Clinical Reviews*, 7(1); DOI: [10.31579/2692-9406/118](https://doi.org/10.31579/2692-9406/118)

Copyright: © 2022 Aamir Jalal Al-Mosawi, This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: Congenital chylous ascites is a very rare condition that has been well recognized as early as the 1950s, and was mostly called “Congenital chyloperitoneum” during that time. Few cases have been reported during the 1960s, and in 1967, Craven et al attributed the development of congenital chylous ascites to obstruction of the cisterna chyli and chylous reflux into the peritoneal space and small intestine and they demonstrated lymphangiographically.

Patients and methods: The use of a unique dietary therapy for the treatment of congenital chylous ascites is briefly described and the relevant literature is described.

Results: An infant with congenital chylous ascites was seen at about the age of two and half months. He was initially treated with intravenous somatostatin infusion, and thereafter by repeated paracentesis, and continued to receive breast and bottle milk feedings. However, when he was first seen, ultrasound was still showing moderate ascites. The boy was successfully treated a combination of a unique form of very low-fat diet and octreotide acetate subcutaneous injections. Treatment was not associated with a negative effect on growth or development, and no evidence of nutritional deficiency was observed.

Conclusion: The case number 112 of congenital chylous ascites is reported and its unique therapeutic approach is described.

Keywords: congenital chylous ascites; Iraq

Introduction

Congenital chylous ascites is a very rare condition that has been well recognized as early as the 1950s, and was mostly called “Congenital chyloperitoneum” during that time [1, 2, 3]. Few cases have been reported during the 1960s [4, 5, 6], and in 1967, Craven et al attributed the development of congenital chylous ascites to obstruction of the cisterna chyli and chylous reflux into the peritoneal space and small intestine and they demonstrated lymphangiographically [6].

In 1975, Weber et al attributed the development of congenital chylous ascites to abdominal lymphdysplasia, and reported the 52nd case of this rare disorder [7].

In 1979, Flores and colleagues reported the occurrence of an autosomal recessive cases congenital chylous ascites in a brother and sister. The parents were first cousins [8].

About seventy-six cases of congenital chylous ascites were reported during half century from the fifties to the end of the 1990s [1-24].

Patients and methods

The use of a unique dietary therapy for the treatment of congenital chylous ascites is briefly described and the relevant literature is described.

Results

An infant with congenital chylous ascites was seen at about the age of two and half months. He was initially treated with intravenous somatostatin infusion, and thereafter by repeated paracentesis, and continued to receive

breast and bottle milk feedings. However, when he was first seen, ultrasound was still showing moderate ascites. The boy was treated a combination of very low-fat and octreotide acetate subcutaneous injections.

Low-fat infant formula was not easily or rapidly available, and the dietary treatment included replacing milk with rice milk with early gradual introduction of juices including orange, grape, and melon juices. Nutritional supplements included amino acids and royal jell capsule.

Octreotide acetate 50 microgram was given by subcutaneous injections weekly for one month, and thereafter, every 10 days for month.

Ascites was completely disappeared after one month of treatment, and didn't recur when the child was seen at about the age of nine months.

Treatment was not associated with a negative effect on growth or development, and no evidence of nutritional deficiency was observed.

Discussion

Congenital chylous ascites is a very rare disorder with about seventy-six cases reported in the medical literature during the 1950s, 1960s, 1970s, 1980s and 1990s, including the case reported by Matsuzaki et al in 1987 which was associated with infantile sialidosis [7-24].

In 1975, Weber et al reported a girl with neonatal chylous ascites that had been treated with repeated paracentesis until about the age of three years when treatment with a low fat diet was started, and after 8 weeks, ascites was no longer recurring [7].

In 1978, Portier et al reported the treatment of a difficult case of congenital chylous ascites by peritoneo-venous shunt [9]. Guttman and colleagues (1982) also reported the treatment of two patients with severe chylous ascites associated with generalized lymphatic malformations, and causing difficulty with breathing with peritoneo-venous shunts [13].

Alliët and colleagues (1992) reported the occurrence of congenital chylous ascites in girl with Down syndrome who experienced respiratory distress secondary at birth because of severe abdominal distension. The patient was treated with total parenteral nutrition and for 10 weeks [18].

In 1996, Caty et al reported a successful treatment of congenital chylous ascites with a somatostatin analog [20].

By the end of the year 2010, there were about 89 cases of congenital chylous ascites reported in the medical literature [25-33].

Machmouchi et al (2000) reported four patients (3 males and one female) with congenital chylous ascites, three of them responded to low-fat diet and medium-chain triglycerides and multiple paracentesis. The fourth patient was a male who had a macro-digit affecting the right index and medius. He continued to have ascites after 10 weeks, but the ascites completely resolved during the third week of complete parenteral alimentation, and no recurrence occurred during one-year of follow-up [25].

Mitsunaga et al (2001) treated two patients with congenital chylous ascites initially conservatively with medium-chain triglyceride milk and total parenteral nutrition, but more than one month treatment was not effective. Thereafter, they successfully treated the patients by lymphatic duct ligation after the identification of the area of chylous leakage using a laparoscopic lymphoid dye test [26].

Lee (2002) reported a patient with congenital chylous ascites who was fully recovered after 26 days of total parenteral nutrition [27].

te Pas et al (2004) reported a boy with congenital chylous ascites who was initially treated with total parental feeding and later by diet with medium-chain triglycerides, and octeotride, a somatostatin analogue. The patient

didn't respond to these therapies and required surgical ligation of the leaking lymphatic duct [28].

Huang and Xu (2005) the treatment of congenital chylous ascites with total parenteral nutrition and somatostatin [29].

Siahanidou et al (2009) reported the occurrence of gall stones in a patient with congenital chylous ascites.

Karagol et al (2010) emphasized the rarity of congenital chylous ascites and reported a patient neonate.

During March, 2022, there were about 111 cases of congenital chylous ascites reported in the medical literature [34-49].

In 2011, Huang et al reported a premature neonate with congenital chylous ascites who was treated successfully with intravenous infusion of octreotide, a synthetic somatostatin analogue after failure to respond to traditional supportive therapies [34].

Spagnol et al (2011) reported two patients with refractory congenital chylous ascites treated surgically successfully by cauterization, and fibrin glue application [35].

Mouravas et al (2012) reported the successful treatment of 4 patients having congenital chylous ascites including one patient treated conservatively and three required surgery. Two patients were treated with excision of intra-abdominal cysts and one patient was treated with ligation of the left lumbar lymphatic trunk and cisterna chyli and the use of fibrin glue [36].

Olivieri et al (2012) reported the treatment of a patient with congenital chylous ascites with a failed conservative management, but thereafter was treated successfully with fasting, total parenteral nutrition and octreotide for a period of 15 days followed by medium chain triglyceride feeds. On day 25, breast feeding started, and was discharged on day 33 [37].

Bhattacharya and colleagues (2012) reported the second case of congenital chylous ascites occurring in association with Down's syndrome, and was also associated with lymphedema [38].

Moreira Dde et al (2013) reported a patient with recurrent chylous ascites who didn't respond to medical treatment. Exploratory laparotomy couldn't identify lymph leakage site that could be ligated. The patient was treated successfully by the placement of a hemostatic oxidized cellulose mesh on lymph exudation areas, with a thin layer of fibrinogen/thrombin glue over the mesh [39].

Purkait et al (2014) reported a 2 and half month infant with congenital chylous ascites who didn't respond to treatment with medium-chain triglycerides formula and paracentesis, but was successfully treated with octreotide intravenous infusion [40].

Sadovnikova and colleagues (2015) emphasized the variable pathogenesis of congenital chylous ascites and reported a patient with congenital chylous ascites caused by peritoneal cavity lymphangioma [41].

Cao et al (2016) reported two patients congenital chylous ascites treated successfully with rice soup and parenteral nutrition [42].

Ermarth (2016) reported the association of congenital chylous ascites caused by primary lymphatic defect with additional vascular anomaly, with Type VI Ehlers-Danlos Syndrome. The patient was treated with octreotide and total parenteral nutritional [43].

Long et al (2018) reported a patient with congenital chylous ascites caused mesenteric lymphatic obstruction associated with intestinal malrotation [44].

Xu et al (2018) reported the association of congenital chyloous ascites phacomatosis pigmentovascularis which is associated with pigmentary nevus and capillary malformation in a five-month old [45].

Carr et al (2019) reported the successful treatment of four patients who had refractory chyloous ascites with complete retroperitoneal exposure with application of fibrin glue and Vicryl mesh [46].

Saxena et al (2020) a patient with congenital chyloous ascites refractory to initial treatment with medium-chain triglyceride based diet followed treatment with total parenteral nutrition plus octreotide. Thereafter, the patient was treated successfully surgically using fibrin glue the area of mesenteric lymphatic leakage [47].

Sooklin et al (2020) a patient with severe congenital chyloous ascites in a preterm neonate who experienced severe respiratory distress at birth and required immediate paracentesis. Treatment with medium-chain triglyceride was not satisfactory, but octreotide treatment was successful [48].

Tamaoka et al (2021) reported the third case of congenital chyloous ascites occurring in association with Down syndrome. The case was refractory to traditional therapies, but was successfully treated with midodrine which is an alpha-1 adrenoreceptor agonist [49].

Successful dietary treatment of congenital chyloous ascites with a low fat diet was probably first reported in 1975 by Weber et al. A variety of surgical treatments have been reported including peritoneo-venous shunts (Portier et al, 1978) and lymphatic duct ligation (Pearl and colleague, 1977; Mitsunaga et al, 2001; Densupsoontorn et al, 2009). Successful treatment with rather lengthy period has also been reported total parenteral nutrition (Alliët and colleagues, 1992; Machmouchi et al, 2000; Lee, 2002).

Successful treatment of congenital chyloous ascites with a somatostatin analog has also been reported (Coty et al, 1996; Purkait et al, 2014; Sooklin et al, 2020), but was not successful all the time (te Pas et al, 2004; Olivieri, et al, 2012; Saxena et al. 2020).

Treatment with low-fat diet and medium-chain triglycerides may not always be successful (Machmouchi et al, 2000; Saxena et al, 2020; Sooklin et al, 2020).

Treatment with total parenteral nutrition with or without medium-chain triglyceride milk also may not always be successful (Mitsunaga et al, 2001; te Pas et al, 2004; Olivieri, et al, 2012; Purkait et al, 2014; Saxena et al, 2020).

Conclusion

The case number 112 of congenital chyloous ascites is reported and its unique therapeutic approach is described.

Conflict of interest: None.

References

- Hurez A, Leveque C. Le chylopéritoine congénital [Congenital chyloperitoneum]. *Gaz Med Fr.* 1952 Jun; 59(11):669-672.
- Janet H, Haquin C, Civatte J, Giraud G, Grandin L. (1953). Chyloperitoine congenital [Congenital chyloperitoneum]. *Nourrisson Jan-Feb;* 41(1):1-9.
- Niznikowska-Marks MJ. Przepadek wrodzonego wylewu mleczu do otrzewnej [A case of congenital chyloperitoneum]. *Pediatr Pol* 1954 Nov; 29(11):1117-1120.
- Alvarado A. Congenital chyloous ascites. *Clin Proc Child Hosp Dist Columbia* 1962 Apr; 18:109-113.
- Freundlich E, Herz N, Falk W. Congenital chyloous ascites and chylothorax. *Case report. Isr Med J* 1962 Jul-Aug; 21:171-177.
- Craven CE, Goldman AS, Larson DL, Patterson M, Hendrick CK. Congenital chyloous ascites: lymphangiographic demonstration of obstruction of the cisterna chyli and chyloous reflux into the peritoneal space and small intestine. *J Pediatr* 1967 Mar; 70(3):340-345.
- Weber HP, Emons D, Knöpfle G, Kowalewski S. (1975). Congenitaler chylöser Ascites. Eigene Beobachtung und Literaturübersicht über 51 Kinder [Congenital chyloous ascites. Case reports and review of 51 recorded cases. *Klin Padiatr Jul;* 187(4):370-376.
- Flores S, Leungas J, Arredondo-Vega F, Guízar-Vázquez J. (1979). Chyloous ascites in sibs from a consanguineous marriage. *Am J Med Genet;* 3(2):145-148.
- Portier F, Hattab R, de Saint Florent G. (1978). Ascite chyleuse congénitale irréductible. Traitement par dérivation péritonéo-veineuse [Irreducible congenital chyloous ascites. Therapy by peritoneo-venous shunt]. *Nouv Presse Med Dec 9;*7(44):4048.
- Evers KG, Huth F. (1977). Congenital lymphatic vessel dysplasia with intestinal lymphangiectasis and chyloous ascites in twins. *Monatsschr Kinderheilkd;* 125:526-529.
- Pearl J, Joyner J, Collins DL. (1977). Chyloous ascites: the first reported surgical cure by direct ligation. *J Pediatr Surg;* 12:687-691.
- Volkman RW. (1978). Chyloous ascites: the first reported surgical cure by direct ligation. *J Pediatr Surg;*13:351.
- Guttman FM, Montupet P, Bloss RS. (1982). Experience with peritoneo-venous shunting for congenital chyloous ascites in infants and children. *J Pediatr Surg Aug;* 17(4):368-372.
- Hogeman PH, Van der Vorm DH, Puyenbroek JI, Exalto N. (1983). Chyloperitoneum in a newborn infant. *Tijdschr Kindergeneesk;* 51:143-146.
- Matsuzaki K, Matsumoto Y, Yoshihara W, Okada S, Midorikawa M, et al. (1987). [A case of infantile sialidosis associated with congenital chyloous ascites]. *No To Hattatsu May;* 19(3):249-253.
- Forg A, Pochard J, Nisand I, Flori J. (1987). Medical treatment of chyloous ascites: apropos of a case. *Pediatrics;* 42:253-254.
- Asada M, Ishikawa K, Yamaguchi K, Miwa S, Shibata H, et al. (1989).[A case of congenital chyloous ascites]. *Nihon Sanka Fujinka Gakkai Zasshi Feb;* 41(2):231-233
- Pallas Alonso CR, Gómez Castillo E, Moreno Ruiz MJ, Martín Puerto MJ, Manzanares López-Manzanares J, Urruzuno Tellería P. (1990). Ascitis quílosa congénita. A propósito de un caso [Congenital chyloous ascites. Report of a case]. *An Esp Pediatr Nov;* 33(5):487-488.
- Liao HB, Hwang RL, Chu DM, Chu ML. (1990). Neonatal chyloous ascites: report of two cases. *Acta Pediatr Sin;* 31:47-52.
- Alliët P, Young C, Lebenthal E. (1992). Chyloous ascites: total parenteral nutrition as primary therapeutic modality. *Eur J Pediatr Mar;* 151(3):213-214.
- Woods CG, Pearce JL, Husson SM. (1992). Recessively determined chyloous ascites: a case report and possible mouse model. *Clin Dysmorphol;* 1:179-181.
- Coty MG, Hilfiker MI, Azizkhan RG, Glick PL. (1996). Successful treatment of congenital chyloous ascites with a somatostatin analog. *Pediatr Surg Int;* 11:396-397.
- Chye JK, Lim CT, van der Heuvel M. (1997). Neonatal chyloous ascites: report of three cases and review of the literature. *Pediatr Surg Int;* 12:296-298.
- D'Agastino S, Costal L, Fbbro MA, Spina P, Musi L. (1998). Neonatal chyloous ascites: a case report. *Pediatr Med Chir;* 20:285-286.

25. Bappal B, Ghani SA, Chaudhary R, Vaishnav A. (1998). Successful surgical treatment of congenital chylous ascites. *Indian Pediatr Mar*; 35(3):276-277.
26. Bagolan P, Bilancioni E, Spina V, Nahoum A. (1999). Fetal tachycardia and chylous ascites. *Br J Obstet Gynecol*; 106:376-378.
27. Machmouchi M, Amin A, Lanjaoui I, Jacobs A, Hatoum C, Al Zahrani D. (2000). Congenital chylous ascites: report of four cases and review of the literature. *Ann Saudi Med Sep-Nov*; 20(5-6):436-439.
28. Mitsunaga T, Yoshida H, Iwai J, Matsunaga T, Kouchi K, et al. (2001). Successful surgical treatment of two cases of congenital chylous ascites. *J Pediatr Surg Nov*; 36(11):1717-1719.
29. Lee YY, Soong WJ, Lee YS, Hwang B. (2002). Total parenteral nutrition as a primary therapeutic modality for congenital chylous ascites: report of one case. *Acta Paediatr Taiwan Jul-Aug*; 43(4):214-216.
30. te Pas AB, vd Ven K, Stokkel MP, Walther FJ. (2004). Intractable congenital chylous ascites. *Acta Paediatr Oct*; 93(10):1403-1405.
31. Huang Y, Xu H. (2005). [Treatment of congenital chylous ascites using total parenteral nutrition and somatostatin: a case report]. *Zhonghua Er Ke Za Zhi Feb*; 43(2):152-153.
32. Densupsoontorn N, Jirapinyo P, Aanpreung P, Laohapensang M, Parichatikanond P. (2009). Congenital chylous ascites: the roles of fibrin glue and CD31. *Acta Paediatr Nov*; 98(11):1847-1849.
33. Siahaidou T, Mikou C, Karra P, Pappas D, Zellos A, Mandyla H. (2010). Cholelithiasis in a neonate with chylous ascites. *J Perinatol Mar*; 30(3):227-230.
34. Karagol BS, Zenciroglu A, Gokce S, Kundak AA, (2010). Ipek MS. Therapeutic management of neonatal chylous ascites: report of a case and review of the literature. *Acta Paediatr Sep*; 99(9):1307-1310.
35. Melo-Filho AA, Souza IJ, Leite CA, Leite RD, Colares JH, Correia JM. (2010). Refractory congenital chylous ascites. *Indian J Pediatr Nov*; 77(11):1335-1337.
36. Huang Y, Zhuang S, Li Y, Liu M, Chen H, Du M. (2011). Successful management of congenital chylous ascites in a premature infant using somatostatin analogue. *Indian J Pediatr. Mar*; 78(3):345-347.
37. Spagnol L, Conforti A, Valfrè L, Morini F, Bagolan P. (2011). Preoperative administration of Sudan III and successful treatment of persistent chylous ascites in a neonate. *J Pediatr Surg May*; 46(5):994-997.
38. Mouravas V, Dede O, Hatzioannidis H, Spyridakis I, (2012). Filippopoulos A. Diagnosis and management of congenital neonatal chylous ascites. *Hippokratia Apr*; 16(2):175-180.
39. Olivieri C, Nanni L, Masini L, Pintus C. Successful management of congenital chylous ascites with early octreotide and total parenteral nutrition in a newborn. *BMJ Case Rep 2012 Sep 25*; 2012:bcr2012006196.
40. Bhattacharya S, Das NK, Chatterjee A. (2012). Congenital chylous ascites and lymphedema in Down's syndrome. *Indian J Pediatr Nov*; 79(11):1532.
41. Moreira Dde A, Santos MM, Tannuri AC, Tannuri U. (2013). Congenital chylous ascites: a report of a case treated with hemostatic cellulose and fibrin glue. *J Pediatr Surg Feb*; 48(2):e17-19.
42. Purkait R, Saha A, Tripathy I, Roy B. (2014). Congenital chylous ascites treated successfully with MCT-Based formula and octreotide. *J Indian Assoc Pediatr Surg Jul*; 19(3):175-177.
43. Sadovnikova IV, Shirokova N Yu, (2015). Tihomirov GV. [Congenital chylous ascites: A case report]. *Eksp Klin Gastroenterol* ;(1):84-87.
44. Cao Y, Yan W, Lu L, Tao Y, Lu W, Chen Y, Tang Q, Cai W. (2016). Parenteral nutrition combined with rice soup can be a safe and effective intervention for congenital chylous ascites. *Asia Pac J Clin Nutr*; 25(3):631-635.
45. Ermarth AK, Pohl J, Esty B, Sempler JK, Carey JC, O'Gorman MA. (2016). Congenital Chylous Ascites and Ehlers-Danlos Syndrome Type VI. *ACG Case Rep J Dec 21*; 3(4):e186.
46. Long L, Zhen C, Yandong W, Ning D, Qi L, Qing G. (2018). Congenital chylous ascites in infants: another presentation of intestinal malrotation. *J Pediatr Surg. Mar*; 53(3):537-539.
47. Xu S, Zhang Q, Liu T, Zhang Y, Sun N. (2018). A female infant with phacomatosis pigmentovascularis and congenital chylous ascites: A case report. *Medicine (Baltimore) Aug*; 97(34):e12012.
48. Carr BD, Grant CN, Overman RE, Gadepalli SK, Geiger JD. (2019). Retroperitoneal exploration with Vicryl mesh and fibrin tissue sealant for refractory chylous ascites. *J Pediatr Surg Mar*; 54(3):604-607.
49. Saxena R, Suchiang B, Pathak M, Sinha A. (2020). Refractory congenital chylous ascites: role of fibrin glue in its management. *J Indian Assoc Pediatr Surg Jul-Aug*; 25(4):245-247.
50. Sooklin L, Anand AJ, Rajadurai VS, Chandran S. (2020). Management of large congenital chylous ascites in a preterm infant: fetal and neonatal interventions. *BMJ Case Rep Sep 2*; 13(9):e235849.
51. Tamaoka S, Osada A, Kin T, Arimitsu T, Hida M. Midodrine, (2021). an oral alpha-1 adrenoreceptor agonist, successfully treated refractory congenital chylous pleural effusion and ascites in a neonate. *Chest Apr*; 159(4):e189-e191.