

Recent Advances in Foot and Ankle Surgery

Nithin Sunku *

Department of Orthopaedics University/Organisation: Global Hospital Trauma Centre, India.

Corresponding author: Nithin Sunku, Department of Orthopaedics University/Organisation: Global Hospital Trauma Centre, India. Email: drsnithin@gmail.com

Received date: July 28, 2022; **Accepted date:** August 19, 2022; **Published date:** August 30, 2022

Citation: Nithin Sunku, (2022), Recent advances in foot and ankle surgery, *J Clinical Orthopaedics and Trauma Care*, 4(4); DOI: 10.31579/2694-0248/042

Copyright: © 2022, Nithin Sunku. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Foot and Ankle has in itself developed as a subspecialty of importance. With growing advance in technology and imaging modalities the treatment and surgery has grown enormously in the field of foot and ankle joints. From simple bony fractures to ankle replacement to arthroscopy it's a fast pace through which we are able to treat the patients with the use of cutting-edge technology.

Keywords: foot and ankle surgery; treatment; joints; cutting edge technology; imaging modalities; FAAM; FAOS; metatarsus adductus; PROMs; FFI

Summary

Foot and Ankle has in itself developed as a subspecialty of importance. With growing advance in technology and imaging modalities the treatment and surgery has grown enormously in the field of foot and ankle joints. From simple bony fractures to ankle replacement to arthroscopy it's a fast pace through which we are able to treat the patients with the use of cutting-edge technology.

Patient-Reported Outcome Measures

The Foot and Ankle Outcome Score (FAOS) is the most used PROMs in foot and ankle evaluation. At present this is used widely and has been translated into several languages. It is applicable in many conditions and has proven validation across multiple foot and ankle diseases hence FAOS is the preferred PROM by many specialists.

In a study by Hansen et al. [1] took into consideration about various scores later on summarized them. Totally there were 17 patient reported outcome measures in the setting of ankle instability, including commonly used measures such as the Foot and Ankle Ability Measure (FAAM), Foot Function Index (FFI), and the Foot and Ankle Outcome Score (FAOS). Their analysis among these for ankle instability showed only 3 scales to be valid enough. Of these scales, only the Foot and Ankle Ability Measure (FAAM) had been rigorously validated for construct validity using modern psychometric methods.

Hallux Valgus

Hallux valgus surgery in patients already having metatarsus adductus is a challenging problem in foot. In a study by Choi et al. [2] they took 173 feet sample with hallux valgus and compared those with metatarsus adductus (42 feet) and those without metatarsus adductus (131 feet) after proximal chevron and Akin osteotomies. At a median follow-up of 32 months, no difference in the improvement of the hallux valgus angle and the intermetatarsal angle between groups. Recurrence rate was significantly higher in the metatarsus adductus group (28.6% compared with 6.1%) [3].

Trauma

In RCT done by White et al. [4] in which fibula fracture fixation by different methods in young patients (mean age, 42 years) either by fibular nail (63 patients) or by plate osteosynthesis (62 patients). They found no difference in the Olerud and Molander Score ($p = 0.621$) between the groups at 1 year and no significant difference in the quality of radiographic reduction, rate of complications, and rate of reoperations between the groups.

Another prospective study by Del Balso et al. was done in which radiographic and functional outcomes were compared. These were patients who underwent either closed reduction - syndesmosis with screw fixation (29 patients) or open reduction syndesmosis with repair of the

anterior inferior tibiofibular ligament (AiTFL) and screw fixation (21 patients). They reported better syndesmotom reduction in the AiTFL group on postoperative CT scans. But no difference was there between groups in overall functional scores. Nonetheless, 12 months, the AiTFL group had better Maryland Foot Shoe subscore and FAOS Quality of Life subscore. They opined that it might be a product of improved reduction.

Arthroscopy

Ankle arthroscopy (AA) is now a well-established treatment method and important armamentarium of the foot and ankle surgeon. It can be used both as a diagnostic tool and also has therapeutic applications in the same sitting using a minimally invasive technique. Many conditions such as high incidence of chondral lesions and other intra-articular abnormalities following ankle fractures can be treated well with arthroscopy. Almost 80% were treated in this manner by study done on 288 patients [5].

Arthroscopy allows for clear visualization and assessment of associated syndesmotom and ligamentous injuries. Chronic ankle instability is such a condition where pathology likely due to contribution of medial ankle ligament damage, which lead to better understanding & refinement of our treatment strategies. Further progress in ligament injury treatment has led to the development of arthroscopic reconstructive procedures of the anterior tibiotalar ligament portion of the deltoid ligament [6,7].

References

- Hansen CF, Obionu KC, Comins JD, Krogsgaard MR. Patient reported outcome measures for ankle instability. An analysis of 17 existing questionnaires. *Foot Ankle Surg.* 2021 May 1; S1268-7731(21):00087-00094.
- Choi SM, Lee JS, Lim JW, ImJM, Kho DH, Jung HG. Effect of metatarsus adductus on hallux valgus treated with proximal reverse chevron metatarsal osteotomy. *Foot Ankle Int.* 2021 Jul; 42(7):886-893.
- Conti MS, Caolo KC, Ellis SJ, Cody EA. Radiographic and clinical outcomes of hallux valgus and metatarsus adductus treated with a modified Lapidus procedure. *Foot Ankle Int.* 2021 Jan; 42(1):38-45.
- White TO, Bugler KE, Olsen L, Lundholm LH, Holck K, Madsen BL, Duckworth AD. A prospective, randomized, controlled, two-center, international trial comparing the fibular nail with open reduction and internal fixation for unstable ankle fractures in younger patients. *J Orthop Trauma.* 2022 Jan 1; 36(1):36-42.
- Hintermann B, Regazzoni P, Lampert C, Stutz G, Gächter A. Arthroscopic findings in acute fractures of the ankle. *J Bone Joint Surg Br.* 2000; 82:345-351.
- Stornebrink T, Emanuel KS, Shimozone Y, Karlsson J, Kennedy JG, Kerkhoffs GM. A change in scope: Redefining minimally invasive. *Arthrosc Tech.* 2017; 6:e1119-e1124.
- Ziai P, Benca E, Skrbensky GV, Wenzel F, Auffarth A, Krp o S, et al. The role of the medial ligaments in lateral stabilisation of the ankle joint: An in vitro study. *Knee Surg Sports Traumatol Arthrosc.* 2015; 23:1900-1906.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

Submit Manuscript

DOI: [10.31579/2694-0248/042](https://doi.org/10.31579/2694-0248/042)

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/clinical-orthopaedics-and-trauma-care>