

Body Mass Index and Complications of Diabetes Mellitus

Monoarul Haque ^{1*}, Nasrin Tamanna ², Mosharraf Hossen ³

¹ Associate Professor, Singhanian University, India.

² Department of Life Science, Kongju National University, South Korea.

³ Department of Biotechnology, Bangladesh Agricultural University, Mymensingh.

***Corresponding Author:** Md Monoarul Haque, Associate Professor, Singhanian University, India.

Received Date: April 12, 2023 | **Accepted Date:** April 26, 2023 | **Published Date:** May 08, 2023

Citation: Haque M., Tamanna N., Hossen M., (2023), Body Mass Index and Complications of Diabetes Mellitus, *Journal of Clinical and Laboratory Research*. 6(1); DOI:10.31579/2768-0487/099

Copyright: © 2023, Monoarul Haque. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Diabetes mellitus is a global public health issue which has multiple complications. In developing countries obesity associated diseases – like diabetes increasing day by day among the urban people. This study was an attempt to assess body mass index and complications of diabetes mellitus among urban people. To observe the association between BMI and complications of diabetes Chi-square test was done. It was observed that complications of diabetes were significantly higher among obese diabetic patients ($p < 0.05$) than that of non-obese diabetic patients. The effect of BMI on complications of diabetes was found statistically significant. Obesity was responsible for causing various complications among diabetic patients. Policy maker should be design and implement appropriate health education programme to prevent obesity and its associated complications among diabetic patients.

Key words: body mass index; diabetes mellitus; complications

Introduction

Diabetes is very harmful disease. It is known that a patient with diabetes develops different types of complications. Normally diabetic patients are suggested to maintain a disciplined life. Diabetic patients are also suggested to control their body weight. If a diabetic patient fails to maintain normal body weight it may develop different physical complications. Thus, it is necessary to understand properly what are the complications may be faced by the obese diabetic patients. There is very little information available regarding the risk factors of obesity associated with diabetes among selected urban people. Bangladesh is one of the developing countries in the world, which is facing rapid urbanization in recent time. [1] It was reported that prevalence of type 2 diabetes is on the rise more in urban areas compared to rural population in Bangladesh. [2-3] Urbanization was found to be associated with a sedentary lifestyle, higher calorie food intake and stressful condition, which might have contributed to the increasing prevalence of diabetes. [4]

Methods:

It was observational study. The study was carried out over a period of one year. This study was conducted in different hospitals in Dhaka city. The study area was selected purposively. I had taken total 200 diabetic patients,

divided into 100 obese diabetic and 100 non-obese diabetic patients. Non-probability purposive sampling was used to collect sample. Personal interview was taken to collect data. Diabetic book was considered for cross check. Diabetic foot was diagnosed by clinical examination. After coding and editing, data were analyzed by using Statistical Package for Social Science or SPSS (version 16.0). The study place was selected purposively. So, the respondents in this study may not be the representatives of subjects and the results cannot be generalized.

Results:

Association between nephropathy and BMI

Table reveals that 19.5% obese (case) and 8% non-obese (control) have nephropathy whereas 72.5% patients both cases and controls have no nephropathy. Association between nephropathy and BMI is statistically significant ($p < 0.05$).

Association between diabetic foot and BMI

Table indicates that 18% obese (case) and 5.5% non-obese (control) have diabetic foot and rest of the patients have no diabetic foot. Association between diabetic foot and BMI is statistically significant ($p < 0.05$).

Nephropathy	BMI		Total	χ^2	p-value
	≥ 27 (Case) %	< 27 (Control) %			
Yes	19.5	8	27.5	13.266	0.000
No	30.5	42	72.5		
Total	50	50	100		

Association between diabetic foot and BMI

Table indicates that 18% obese (case) and 5.5% non-obese (control) have diabetic foot and rest of the patients have no diabetic foot. Association between diabetic foot and BMI is statistically significant ($p < 0.05$).

Diabetic foot	BMI		Total	χ^2	p-value
	≥ 27 (Case) %	< 27 (Control) %			
Yes	18	5.5	23.5	17.383	0.000
No	32	44.5	76.5		
Total	50	50	100		

Association between retinopathy and BMI

Table reveals that 27% obese (case) and 7% non-obese (control) have retinopathy and rest of the patients have no retinopathy. Association between retinopathy and BMI is statistically significant ($p < 0.05$).

Retinopathy	BMI		Total	χ^2	p-value
	≥ 27 (Case) %	< 27 (Control) %			
Yes	27	7	30	35.651	0.000
No	23	43	70		
Total	50	50	100		

Discussion:

Overweight and obesity continue to increase substantially worldwide, affecting all ages, sexes and races. Worldwide about 58% of diabetes mellitus and 21% of ischemic heart disease are attributable to BMI above 21 kg/m². In a study that examined ethnic differences in the strength of association between BMI and hypertension, higher prevalence of hypertension was associated with higher BMI levels in different ethnic groups. [6] Significant associations between BMI and BP have also been documented in lean Chinese populations. [7] In a prospective cohort, increasing BMI was associated with a steady increase in the risks of total, ischemic, and hemorrhagic stroke. Although concomitant hypertension and diabetes accounted for much of the increase in total and ischemic stroke, a significant increase remained after adjustment for these potential biological mediators. Although the risk of ischemic stroke was highest among smokers and individuals with hypertension, these factors did not substantially modify the relationship between BMI and stroke. Body mass index was not associated with the severity of total and ischemic stroke, but the data suggest that it might be inversely associated with severity of fatal hemorrhagic stroke, particularly subarachnoid hemorrhage. The association between excess weight and stroke risk has been controversial. Among men, few prospective studies have investigated this relationship. Some of these had small sample sizes [8] and others did not classify stroke subtypes. [9] To my knowledge, the association between BMI and stroke severity has not been examined prospectively before. Association between stroke and BMI is

statistically significant in this study. Individuals with diabetes mellitus are 17 times more likely to have an amputation as a result of peripheral vascular disease and are at an increased risk of developing nephropathy, retinopathy, and coronary heart disease, among other adverse health outcomes. [10] The present study showed similar findings. Increasing evidence suggests that obesity is a risk factor for diabetes and chronic kidney diseases. As a marker of obesity, high body mass index has been reported to be related with diabetic nephropathy and end-stage renal disease. [11]

Conclusion

Chance of suffering from nephropathy, retinopathy and diabetic foot was higher among obese diabetic patients. The effect of BMI on complications of diabetes was found statistically significant. Appropriate awareness programme can be commenced immediately.

References

1. Laskar SI. (1996). Urbanization in Bangladesh: Some contemporary observations. *Bangladesh Dev Stud.*; 24: 207-216.
2. Sayeed MA, Mahtab H, Khanum PA, Latif ZA, Ali SMK, et al. (2003). Diabetes and Impaired fasting glycaemia in rural population of Bangladesh. *Diabetes Care*; 26: 1034-1035.
3. Hussain A, Rahim MA, Azad Khan AK, Ali SMK. (2005). Type 2 diabetes in rural and urban population: Diverse

- prevalence and associated risk factors in Bangladesh. *Diabet Med.* 22: 931-937.
4. Ramachandran A, Snehalatha C, Latha E, Vijay V, Vishanathan M. (1999). Impacts of urbanization on life style and on the prevalence of diabetes in native Asian Indian population. *Diabet Res Clin Pract.*; 44:207-213.
 5. World Health Organization. World Health Report 2002: Reducing risks, Promoting Healthy Life. World Health Organization: Geneva,
 6. Colin AB, Linda SA, Popkin BM. (2002). Ethnic differences in the association between body mass index and hypertension. *Am J Epidemiol*; 155: 346–353.
 7. Hu FB, Wang B, Chen C, Jin Y, Yang J, et al. (2000). Body mass index and cardiovascular risk factors in a rural Chinese population. *Am J Epidemiol*;151: 88–97.
 8. Abbott RD, Behrens GR, Sharp DS et al. (1994). Body mass index and thromboembolic stroke in nonsmoking men in older middle age: the Honolulu Heart Program. *Stroke.* ;25:2370- 252376
 9. Rhoads GG, Kagan A. (1983). The relation of coronary disease, stroke, and mortality to weight in youth and in middle age. *Lancet* ;1492- 1495
 10. Gray N et al. (2015). Relation between BMI and Diabetes Mellitus and Its Complications among US Older Adults. *South Med J.*108(1):29-36.
 11. Hsu CY, McCulloch CE, Iribarren C, Darbinian J, Go AS (2006). Body mass index and risk for end-stage renal disease. *Ann Intern Med*, 144(1):21-28.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

[Submit Manuscript](#)

DOI:10.31579/2768-0487/099

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/journal-of-clinical-and-laboratory-research>