

Acute Rheumatic Fever: The Rare Association with Vitamin B12 Deficiency Induced Neutropenia

Aamir Al-Mosawi

Advisor doctor and expert trainer, The National Center of Training and Development and Baghdad Medical City, Baghdad, Iraq.

***Corresponding Author:** Aamir Al-Mosawi, Advisor doctor and expert trainer, The National Center of Training and Development and Baghdad Medical City, Baghdad, Iraq.

Received date: November 10, 2023; **Accepted date:** November 20, 2023; **Published date:** November 30, 2023

Citation: Aamir Al-Mosawi, (2023), Acute Rheumatic Fever: The Rare Association with Vitamin B12 Deficiency Induced Neutropenia, *J Clinical Orthopaedics and Trauma Care*, 5(5); DOI: [10.31579/2694-0248/073](https://doi.org/10.31579/2694-0248/073)

Copyright: © 2023, Aamir Al-Mosawi. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Acute rheumatic fever is an autoimmune disorder that develops commonly few weeks following a group A beta hemolytic Streptococcal tonsillopharyngitis. The autoimmune response can affect joints, skin, heart, and the brain. One of the important serious consequences of acute rheumatic fever that have to be prevented is the the long-term heart disease resulting from valves damage which may occur after one severe illness or following multiple recurrence of the illness. The diagnosis of acute rheumatic fever is generally based on the presence of either two Major diagnostic criteria (multiple joint arthritis, carditis, erythema marginatum, chorea, and subcutaneous nodules) or one Major diagnostic criteria and two Minor criteria (Fever, arthralgia, first degree heart block, elevated inflammatory markers [ESR and C-reactive protein]) plus appropriate evidence of preceding streptococcal infection.

The association of acute rheumatic fever with vitamin B12 deficiency-induced neutropenia has not been reported previously in the medical literature.

Patients and methods: The case of a ten-year old girl who developed acute febrile illness associated with arthralgia, arthritis, skin rash and neutropenia is described.

Results: The patient developed fever and arthritis of one elbow (Painful and swollen) followed shortly by the development of macular rash consisting of pink rings mostly on the extensor surfaces and arthritis of the second elbow. The patient had a previous febrile illness associated with sore throat which occurred during the previous 2 to 3 weeks. ASO titer was 150 iu/ml and CRP was positive 300 mg/dL. The patient had a high ESR, leukopenia, and blood film showed severe neutropenia, poikilocytosis and, frequent elliptocytosis. Serum vitamin B₁₂ level was low, 152 pg/mL. Echocardiography showed low mitral valve prolapse.

Conclusion: The novel association of acute rheumatic fever with vitamin B12 deficiency-induced neutropenia is reported.

Keywords: acute rheumatic fever; vitamin B12 deficiency; induced neutropenia; acute febrile illness

Introduction

Acute rheumatic fever is an autoimmune disorder that develops commonly few weeks following a group A beta hemolytic Streptococcal tonsillopharyngitis. The autoimmune response can affect joints, skin, heart, and the brain. One of the important serious consequences of acute rheumatic fever that have to be prevented is the the long-term heart disease resulting from valves damage which may occur after one severe illness or following multiple recurrence of the illness.

Some of the important features of acute rheumatic fever have been known as early as the 1700s.

In 1797, Matthew Baillie quoted David Pitcairn (Figure-1A) who suggested that rheumatism is one of the causes of a morbid growth of the heart.

Matthew Baillie (Figure-1B) reported the presence of a thickening of some valves of heart in autopsies of patients who had acute rheumatism [1].



(1A)



(1B)

Figure (1A): David Pitcairn (1749-1809), a Scottish physician; (1B): Matthew Baillie (1761-1823), a British physician and pathologist

In 1803, Wagstaffe of Southwark reported a fatal case of rheumatic pericarditis with enlargement of the heart [2], and Crowfoot of Beccles reported two cases of rheumatic carditis in 1809 [3].

In 1809, David Dundas reported nine cases of heart disease associated with rheumatism that had been observed since 1770. The cases reported by Dundas were considered typical cases of acute rheumatic carditis. The heart disease started after, or sometimes during the same time of rheumatism. The cases of Dundas were mostly young patients who developed chest pain cough, dyspnea, and palpitations. There was a high death rate with most patients died because of congestive cardiac failure [4].

In 1812, William Charles Wells was still thinking that the association of rheumatism with heart disease was a concurrence rather than a

specific association as he quoted Morgagni and Ferriar who had regarded the association as apparently an interesting coincidence [5, 6].

Richard Bright (Figure-1C) was probably the first to recognize the association of neurological manifestations of rheumatic fever (Sydenham's chorea) with joint manifestations (Rheumatism) in 1831 [7].

Walter Butler Cheadle (Figure-1D) was most probably the first to link rheumatism and heart disease with tonsillitis in 1889. He reported his observations during twenty years on rheumatic fever at the Hospital for Sick Children, Great Ormond Street [8].



(1C)



(1D)

Figure (1C): Richard Bright (1789-1858), an English physician; (1D): Walter Butler Cheadle (October 1836-March 1910), an English pediatrician.

Freeric John Poynton (Figure-1E) and his colleague Paine were most probably the first to identify the diplococcus which they isolated from patients with acute rheumatic fever as the causative bacteria for the

disease [9]. In 1944, Thomas Duckett Jones (Figure-1F) suggested diagnostic criteria for its diagnosis [10].



(1E)



(1F)

Figure (1E): Frederic John Poynton (June 26, 1869-October, 29 1943), an English physician; **(1F): Thomas Duckett Jones** (February 2, 1899-November 22, 1954) was an American physician, cardiologist.

Jones diagnostic criteria have been revised and modified in 1965, 1984, 1992, and 2002. The diagnosis of acute rheumatic fever is generally based on the presence of either two Major criteria (multiple joint arthritis, carditis, erythema marginatum, chorea, and subcutaneous nodules) or one Major criteria and two Minor criteria (Fever, arthralgia, first degree heart block, elevated inflammatory markers [ESR and C-reactive protein]) plus appropriate evidence of preceding streptococcal infection [11-14].

The association of acute rheumatic fever with vitamin B12 deficiency-induced neutropenia has not been reported previously in the medical literature.

Patients and methods

The case of a ten-year old girl who developed acute febrile illness associated with arthralgia, arthritis, skin rash and neutropenia is described.

Results

The patient developed fever and arthritis of one elbow (Painful and swollen) on the 22nd of August, 2023 followed shortly by the development of macular rash consisting of pink rings mostly on the extensor surfaces and arthritis of the second elbow. The face was not involved by the rash. She was treated with oral antibiotics and analgesics without obvious improvement. The patient was brought to us on the 26th of August, and she still had painful swollen elbows and rash (Figure-2). The patient had a previous febrile illness associated with sore throat which occurred during the previous 2 to 3 weeks.



Figure (2A): Macular rash consisting of pink rings mostly on the extensor surfaces.



Figure (2B): Macular rash consisting of pink rings mostly on the extensor surfaces



Figure (2C): Macular rash consisting of pink rings mostly on the extensor surfaces

ASO titer was 150 iu/ml and CRP was positive 300 mg/dL.
 Erythrocyte Sedimentation Rate (ESR) was 100 cubic mm/hour.
 Hemoglobin was 8.9 g/dL.
 White blood cell count was $1.58 \times 10^9/L$ (Normal: 4.0 to $11.0 \times 10^9/L$).
 Granulocyte (Including neutrophils) count was $0.9 \times 10^9/L$ (Normal: 2.0 to $7.0 \times 10^9/L$).
 Blood film showed:

White blood cells: Severe neutropenia, no immature or abnormal cells.
 Red blood cells: Poikilocytosis (10% or more of blood cells have abnormal shape), frequent elliptocytosis, occasional polychromasia (Excessive immature blood cells).
 Platelets: Normal.
 Serum folic acid was within normal range.
 Serum vitamin B12 level was 152 pg/mL (Normal: 200-900 pg/mL).

Radiographs of the left elbow showed no significant abnormalities (Figure-3).

Ultrasound of the elbows showed fluid collection suggesting joint effusion (Figure-4).




Figure 3 (A): Radiographs of the left elbow; **(B):** Radiographs of the left elbow; **(C):** Radiographs of the left elbow.



Figure 4: Ultrasound of the elbows showed fluid collection suggesting joint effusion

Erythrocyte Sedimentation Rate (ESR) was 19 cubic mm/hour.

Hemoglobin was 10.1 g/dL.

White blood cell count was $2.7 \times 10^9/L$ (Normal: 4.0 to $11.0 \times 10^9/L$).

Granulocyte (Including neutrophils) was 29.6 %.

Serum vitamin B₁₂ level was 210 pg/mL (Normal: 200-900 pg/mL).

Discussion

Zeynep Canan Ozdemir from Turkey and her research group reported the etiological causes of neutropenia in 94 children. 55 children (58.5%) had post-infectious neutropenia with 38.4% occurring following upper respiratory airway. 23 (24.6%) children had idiopathic neutropenia. Six children (6.3%) had drug-induced neutropenia. Five children (5.3%) had vitamin B₁₂ deficiency-induced. Two children (2%) had chronic benign neutropenia. One child (1.1%) had immune deficiency, one child (1.1%) had autoimmune lymphoproliferative syndrome, and one child (1.1%) had post-infection hemophagocytic lymphohistiocytosis.

Neutropenia improved in ninety-one children (96.8%).

Neutropenia persisted in 3 children (3.2%).

One died because of infection [24].

Although, the patient in this report had lab-confirmed vitamin B₁₂ deficiency, the blood film showed no macrocytosis, but showed poikilocytosis and frequent elliptocytosis.

As early as 1982, Spivak emphasized that megaloblastic anemia can be masked by the absence of macrocytosis. Spivak also emphasized that poikilocytosis may indicate a vitamin deficiency [25].

As early as 1982, Schoemaker and colleagues, emphasized that vitamin B deficiency may result in an abnormal red blood cells' membrane causing elliptocytosis [26].

In this patient, echocardiography showed low mitral valve prolapse.

As early as 1980, Storozhakov et al suggested the potential of echocardiographic diagnosis of rheumatic valvulitis and mitral valve prolapse in children [27].

In 1988, Lembo et al emphasized that acute rheumatic fever commonly results in mitral regurgitation without associated mitral valve stenosis. They reported that the surgical findings have showed a high frequency of mitral prolapse in rheumatic valvular heart disease suggesting that acute rheumatic fever can cause mitral valve prolapse.

Lembo et al reported thirty patients who had previously acute rheumatic fever. 25 patients (84%) were found to have mitral regurgitation on doppler examination. Echocardiography showed mitral prolapse in 24 patients (80%). Only one patient had mitral stenosis when examined by echocardiography.

Therefore, Lembo et al suggested that acute rheumatic fever can cause mitral valve prolapse [28].

The use of ibuprofen in acute rheumatic fever was reported as early as the 1970s [15].

In 2021, Derya Altay from Turkey and her research team reported 286 patients with acute rheumatic fever. 53 patients (18.5%) [Mean age: 10.7 ± 2.5 years] were treated initially with aspirin for arthritis.

Aspirin-associated hepatotoxicity occurred in 9 (17%) of the 53 patients, and therefore ibuprofen or naproxen was given as an

alternative to aspirin. Both ibuprofen and naproxen were not associated with side effects.

30% of 53 patients had anemia, and aspirin was used for longer period in the hepatotoxic anemic patients than non-anemic patients (p=0.02) [16].

Münevver Yılmaz and her colleagues emphasized that the use of Aspirin (Acetylsalicylic acid) in acute rheumatic fever can be associated with severe side effects [17].

The use of single injections of benzathine penicillin given every month in the treatment of acute rheumatic fever was reported as early as the 1950s [18-21].

In 2013, Nigel Wilson from New Zealand reviewed the literature and emphasized that randomized controlled studies showed four weekly intramuscular long-acting benzathine penicillin G was found to be more effective than penicillin given orally to prevent recurrence of rheumatic fever [22].

In 2022, Anna P Ralph and Bart J Currie from Australia emphasized that the aims of the treatment of acute rheumatic fever include relieving symptoms, lessening heart valve damage and eliminating streptococcal infection. They recommended the use of monthly intramuscular injections benzathine benzyl penicillin G for the prevention of recurrences [23].

Conclusion

The novel association of acute rheumatic fever with vitamin B₁₂ deficiency-induced neutropenia is reported.

Acknowledgement

The author has the copy right of all figures and sketches included in this paper.

Conflict of interest

None.

References

1. Baillie M. The Morbid Anatomy of Some of the Most Important Parts of the Human Body, 2nd ed. J. Johnson and G. Nicol, London, 1797.
2. Wagstaffe MF. A Case of Thoracic Adhesions. *Med Phys J* 1803 Apr 1; 9(50):359-362.
3. Crowfoot WH. Two cases of carditis. *Edinburgh Medical and Surgical Journal* 1809; 5:298-300.
4. Dundas D. An account of a peculiar Disease of the Heart. *Med Chir Trans* 1809; 1:37-46.
5. Wells WC. On rheumatism of the heart. *Transactions of the Society for Improving Medical and Chirurgical Knowledge* 1812; 3:373-424.
6. Morgagni GB. *De sedibus, et causis morborum* (1761). Translated by B Alexander. A. Millar and T. Cadell, London, 1769.
7. Bright R. Reports of medical cases. Vol 2 Disease of the brain and nervous system: selected with a view of illustrating the symptoms and cure of diseases by reference to morbid anatomy. London: Longman, 1831:493.
8. Cheadle WB. Harveian lectures on the various

- manifestations of the rheumatic state as exemplified in childhood and early life. *Lancet* 1889; 133:821-827.
9. Poynton FJ, Paine A. The etiology of rheumatic fever. *Lancet* 1900; 156:861-869.
 10. Jones TD. Diagnosis of rheumatic fever. *JAMA* 1944; 126(8):481-484.
 11. Jones criteria (revised) for guidance in the diagnosis of rheumatic fever. *Circulation* 1965 Oct; 32(4):664-668.
 12. Jones Criteria (revised) for guidance in the diagnosis of rheumatic fever. *Circulation* 1984 Jan; 69(1):204A-208A.
 13. Guidelines for the diagnosis of rheumatic fever. Jones Criteria, 1992 update. Special Writing Group of the Committee on Rheumatic Fever, Endocarditis, and Kawasaki Disease of the Council on Cardiovascular Disease in the Young of the American Heart Association. *JAMA* 1992 Oct 21; 268(15):2069-2073.
 14. Ferrieri P. Jones Criteria Working Group. Proceedings of the Jones Criteria workshop. *Circulation*. 2002 Nov 5; 106(19):2521-2523.
 15. Morino F, Possati F, Santarelli P, Paolini E, Calafiore AM. L'impiego dell'ibuprofen nel controllo dell'attività reumatica [Use of ibuprofen in the control of rheumatic activity]. *Minerva Med* 1973 Jun 23; 64(47):2475-8 [Article in Italian].
 16. Altay D, Pamukçu Ö, Baykan A, Üzümlü K, Arslan D. Aspirin-induced hepatotoxicity and anemia in children with acute rheumatic fever. *Turk J Pediatr*. 2021; 63(2):193-199. Doi: 10.24953/turkjped.2021.02.002.
 17. Yılmaz M, Gürses D, Tükenmez G. The effectiveness and safety of ibuprofen and acetylsalicylic acid in acute rheumatic fever. *Pediatr Int* 2022 Jan; 64(1):e15133.
 18. Chamovitz R, Catanzaro FJ, Stetson CA, Rammelkamp CH Jr. Prevention of rheumatic fever by treatment of previous streptococcal infections. I. Evaluation of benzathine penicillin G. *N Engl J Med* 1954 Sep 16; 251(12):466-471.
 19. Stollerman GH. Repository benzathine penicillin for the control of rheumatic fever. *Bull Rheum Dis* 1954 Dec; 5(4):79-80.
 20. Markowitz M, Hemphill W. A comparison of oral benzathine penicillin G and sulfonamides for the prevention of streptococcal infections and recurrences of rheumatic fever. *Pediatrics* 1955 May; 15(5):509-515.
 21. Stollerman GH, Rusoff JH, Hirschfeld I. Prophylaxis against group A streptococci in rheumatic fever; the use of single monthly injections of benzathine penicillin G. *N Engl J Med* 1955 May 12; 252(19):787-792.
 22. Wilson N. Secondary prophylaxis for rheumatic fever: simple concepts, difficult delivery. *World J Pediatr Congenit Heart Surg* 2013 Oct; 4(4):380-384.
 23. Ralph AP, Currie BJ. Therapeutics for rheumatic fever and rheumatic heart disease. *Aust Prescr* 2022 Aug; 45(4):104-112.
 24. Ozdemir ZC, Kar YD, Kasaci B, Bor O. Etiological causes and prognosis in children with neutropenia. *North Clin Istanbul* 2021 May 24; 8(3):236-242.
 25. Spivak JL. Masked megaloblastic anemia. *Arch Intern Med* 1982 Nov; 142(12):2111-2114.
 26. Schoomaker EB, Butler WM, Diehl LF. Increased heat sensitivity of red blood cells in hereditary elliptocytosis with acquired cobalamin (vitamin B12) deficiency. *Blood* 1982 Jun; 59(6):1213-1219.
 27. Storožhakov GI, Kuz'mina NN, Shaikov AV, Voronina NM, Malysheva NV. Vozmozhnosti ékhokardiograficheskoi diagnostiki revmaticeskogo val'vulita i prolapsa mitral'nogo klapana u detei [Potentials of echocardiographic diagnosis of rheumatic valvulitis and mitral valve prolapse in children]. *Vopr Revm* 1980 Jan-Mar ;(1):18-22 [Article in Russian].
 28. Lembo NJ, Dell'Italia LJ, Crawford MH, Miller JF, Richards KL, O'Rourke RA. Mitral valve prolapse in patients with prior rheumatic fever. *Circulation* 1988 Apr; 77(4):830-836.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

[Submit Manuscript](#)

DOI:10.31579/2694-0248/073

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more at: <https://www.auctoresonline.org/journals/clinical-orthopaedics-and-trauma-care->