

Peripheral Sensory Stimulation for Neurological Disorders. A Novel, Non-invasive Therapeutic Option. Review Article

Heather A. Nussbaum, Eric S. Nussbaum *

National Brain Aneurysm & Tumor Center, Twin Cities, MN, USA.

*Corresponding Author: Eric S. Nussbaum, National Brain Aneurysm & Tumor Center, Twin Cities, MN, USA.

Received date: **May 29, 2023**; Accepted date: **June 06, 2023**; Published date: **June 13, 2023**

Citation: Heather A. Nussbaum, Eric S. Nussbaum (2023), Peripheral Sensory Stimulation for Neurological Disorders. A Novel, Non-invasive Therapeutic Option. Review Article, *J. Neuroscience and Neurological Surgery*. 13(3); DOI:10.31579/2578-8868/267

Copyrights: © 2023, Eric S. Nussbaum. This is an open-access article distributed under the terms of The Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Abstract

Peripheral sensory stimulation (PSS) represents a new area of therapy for patients with neurological disorders. Grounded in strong basic science research, PSS carries potential promise in the management of multiple diseases. We reviewed the literature regarding those conditions that represent reasonable targets for treatment with PSS including stroke, Parkinson's disease, traumatic brain injury, and auto-immune, inflammatory illness. Potential mechanisms of action are discussed, and future avenues of investigation are described.

Key words : neurological disorders; peripheral sensory stimulation; traumatic brain injury

Introduction

Over the past decade, there has been a growing interest in the field of neuromodulation including the use of peripheral sensory stimulation (PSS) in the management of a variety of neurological disorders [1-5]. Such PSS treatment is extremely attractive because it may enhance neurological recovery from a variety of prevalent conditions with essentially no risk of serious complications [3-6]. This paper summarizes the rationale and potential uses of PSS as a method of improving neurological conditions in the areas of stroke, Parkinson's disease, and other common neurological ailments.

Discussion

Stroke

Ischemic stroke impacts roughly 700,000 individuals per year in the United States resulting in significant short-term and long-term disability and tremendous financial loss [7,8]. Although the past 25 years have seen a revolution in the management of acute ischemic stroke based on the introduction of intravenous and intra-arterial thrombolytic therapy and mechanical thrombectomy, there has been little change in the therapeutic options available to patients who have suffered a completed stroke and are left with neurological deficit [9-13].

In rodent models, PSS has been shown to significantly and reproducibly improve neurological outcomes following ischemic injury [14-16]. Small clinical trials in humans have suggested that such PSS can improve recovery after stroke as well (3,4,17-24). In general, stimulation has been directed at the affected hand (contralateral to the injured hemisphere) and may be delivered using vibrational, electrical, or pneumatic input through the delivery of air puffs [4, 17-20]. Whether the duration, intensity, or frequency of treatment will impact the effectiveness of therapy remains an ongoing area of investigation [4].

The mechanism by which PSS may improve outcome after stroke has not been elucidated, although it appears that somatosensory input modulation of corticomotor excitability may be involved [25]. It has been suggested that somatosensory therapy may enhance neuronal reorganization after stroke or traumatic brain injury, possibly utilizing intact neural circuits either in the ipsilateral or contralateral hemispheres [26-32]. Alternatively, peripheral stimulation may improve regional cerebral perfusion, encouraging recruitment of pial collateral channels and recovery of under-perfused neurons [4]. Our previous work suggests that intermittent PSS using puffs of air may provide benefit in the setting of acute stroke, and improvement in CBF as measured by perfusion MR imaging suggests that recruitment of collateral supply may play at least a partial role in this process (Figure 1).




Figure 1: Example of a peripheral sensory stimulation device that uses air puffs delivered to the hand as a mechanism of stimulation. (courtesy, Neuro Glove, Inc.)

Parkinson's Disease

It is estimated that Parkinson's disease affects more than one million individuals in the United States, and the incidence of the disease appears to be increasing as the population ages [33-4]. Although symptoms can be controlled and improved using pharmacological therapy, the disease tends to be progressive and results in substantial disability and financial loss [33-4]. The motor deficits seen in patients with Parkinson's disease have been linked to abnormally synchronized neuronal activity in dopaminergic circuits [5]. In these patients, coordinated reset based on tactile feedback has been suggested as a mechanism to overcome excessive neuronal synchronization resulting in sustained "unlearning" of pathological synaptic connectivity and synchrony [35]. Perhaps based on such a "coordinated reset" mechanism, peripheral somatosensory stimulation has been shown to benefit patients with Parkinson's disease [5,35-38].

Early small clinical studies using PSS have shown dramatic and promising improvement in Parkinson's patients with long-standing and severe motor symptoms [5,35-38]. The potential to improve quality of life in a significant manner using a non-invasive alternative to medical therapy has encouraged further evaluation of this option for patients with Parkinson's, and larger clinical trials are ongoing. It is anticipated that these trials will help to elucidate the mechanisms by which PSS may benefit patients with movement disorders such as Parkinson's disease and will also help to define those patients and symptoms most amenable to such therapy.

Other Conditions

Because of the potential for enhancing neuronal "learning" through reorganization as well as the other mechanisms previously described, PSS has been suggested as a possible mechanism to help patients with other neurological conditions including post-traumatic brain injury and autoimmune, inflammatory conditions such as multiple sclerosis. [4, 39-41] Although the mechanism by which PSS may be beneficial in such instances remains unclear, it is possible that recruitment of regional collateral blood supply, enhancement of neuronal recovery, development of new neuronal connections, and local "coordinated reset mechanisms" may all play a role in these disorders as in stroke and Parkinson's Disease [44-45].

At this point, there is inadequate evidence to prove the effectiveness of PSS in these conditions, but as trials evaluating the usefulness of this technique in stroke and Parkinson's disease expand, we will learn more about the effectiveness and mechanisms by which peripheral stimulation can affect brain functioning, and potential indications for its use may expand as well.

Because of the extremely low-risk nature of PSS treatment, it would be safe and straightforward to evaluate its effectiveness in such conditions.

Conclusions

The use of PSS as a neuromodulation therapy represents a novel, non-invasive, low-risk treatment option for patients with various neurological disorders. Preliminary and small clinical studies have shown promise using this technique in patients with stroke and Parkinson's disease. At this point, larger studies are underway and will hopefully answer questions of efficacy in a more rigorous fashion. Questions that remain to be addressed include the importance of the mechanism, frequency, and duration of stimulus delivery on patient outcomes. In addition, the possible usefulness of such therapy in patients with other conditions such as post-traumatic brain injury remains to be evaluated.

References

1. Glenn B, Francio VT, Westerhaus BD, Goree J, Strand NH, Sparks D, Petersen E: Accessibility and Ease of Use in Neuromodulation Devices. *Neuromodulation: Technology at the Neural Interface* 2023; 1094, 7159, doi.org/10.1016/
2. Ting W, Fadul F, Fecleu S, Ethier C: Neurostimulation for stroke rehabilitation. *Front Neurosci* 2021;15:649459, doi: 10.3389/fnins.2021.649459
3. Celnik P, Hummel F, Harris-Love M, Wolk R, Cohen LG. Somatosensory stimulation enhances the effects of training functional hand tasks in patients with chronic stroke. *Archives of Physical Medicine and Rehabilitation*, 2007;88(11), 1369–1376.
4. Nussbaum ES, Janjua TM, Lowary J, Defillo A, Myers M, Nussbaum LA: Peripheral sensory stimulation of the hand in the treatment of stroke. A preliminary study of safety and effectiveness. *J Neurol Disord* 11(1):2023, in press
5. Kristina JP, Kromer JA, Cook AJ, Hornbeck T, Lim EA, Mortimer BJ, Fogarty A, Han SS, Dhall R, Halpern C, Tass PA.: Coordinated reset vibrotactile stimulation induces sustained cumulative benefits in parkinson's disease. *Front Physiol* 2021; 12: 624317
6. Lucente G, Valls-Sole J, Murillo N, Rothwell J, Coll J, Davalos A, Kumru H: Noninvasive Brain Stimulation and Noninvasive

- Peripheral Stimulation for Neglect Syndrome Following Acquired Brain Injury. *Neuromodulation: Technology at the Neural Interface* 23(3)2020, 312-323, <https://doi.org/10.1111/ner.13062>
7. GBD 2019 Stroke Collaborators: Global, regional, and national burden of stroke and its risk factors, 1990-2019: a systematic analysis for the Global Burden Disease Study 2019. *Lancet Neurology*, 20(10):795-820, 2021, [doi.org/10.1016/S1474-4422\(21\)00252-0](https://doi.org/10.1016/S1474-4422(21)00252-0)
 8. Jackson G, Chari K. National Hospital Care Survey Demonstration Projects: Stroke Inpatient Hospitalizations. *Natl Health Stat Report*. 2019 Nov;(132):1-11.
 9. Bekhmer O, Beumer F, Berg V, Lingsma H, Schonewile Y, Nederkoom V. The randomized trial of intra-arterial treatment for acute ischemic stroke (Mr. CLEAN). *N Engl J Med* 2015; 372
 10. Ogawa A, Mori E, Minematsu K, Taki W, Takahashi A, Nemoto S, Miyamoto S, Sasaki M, Inoue T; MELT Japan Study Group. Randomized trial of intraarterial infusion of urokinase within 6 hours of middle cerebral artery stroke: the middle cerebral artery embolism local fibrinolytic intervention trial (MELT) Japan. *Stroke*. 2007;38(10):2633-9. doi: 10.1161/STROKEAHA.107.488551. Epub 2007 Aug 16. PMID: 17702958.
 11. Broderick JP, Palesch YY, Demchuk AM, et al. Interventional Management of Stroke (IMS) III Investigators. Endovascular therapy after intravenous t-PA versus t-PA alone for stroke. *N Engl J Med*. 2013; 368(10):893-903
 12. Ciccone A, Valvassori L, Nichelatti M, et al. SYNTHESIS Expansion Investigators. Endovascular treatment for acute ischemic stroke. *N Engl J Med*. 2013; 368(10):904-913
 13. Jovin T, Chamorro A, Cobo E, Miquel M, Molina C, Rovira A. Thrombectomy within 8 hours after symptom onset in ischemic stroke (REVASCAT). *N Engl J Med*. 2015
 14. Lay CC, Jacobs N, Hancock AM, Zhou Y, Frostig RD. Early stimulation treatment provides complete sensory-induced protection from ischemic stroke under isoflurane anesthesia. *The European Journal of Neuroscience*. 2013;38: 2445-52. PMID 23586641 DOI: 10.1111/Ejn.12217
 15. Frostig RD, Lay CC, Davis MF: A rat's whiskers point the way toward a novel stimulus-dependent protective stroke therapy. *The Neuroscientist*, 2012. DOI.10.1177/107385841246260701.
 16. Hancock AM, Lay CC, Davis MF, Frostig RD. Sensory Stimulation-Based Complete Protection from Ischemic Stroke Remains Stable at 4 Months Post-Occlusion of MCA. *Journal of Neurological Disorders* 2013; 1: 135. PMID 24634892 DOI: 10.4172/2329-6895.1000135
 17. Conforto AB, dos Anjos SM, Bernardo WM, Silva AA, d Conti J, Machado AG, Cohen LG. Repetitive peripheral sensory stimulation and upper limb performance in stroke: A systematic review and meta-analysis. *Neurorehabilitation and Neural Repair* 2018; 32(10), 863-871. DOI. 10.1177/1545968318798943
 18. Conforto AB, Cohen LG, Dos Santos RL, Scaff M, Marie SKN. Effects of somatosensory stimulation on motor function in chronic cortico-subcortical strokes. *Journal of Neurology*, 2007;254(3), 333-339.
 19. Conrad MO, Scheidt RA, Schmidt BD. Effects of wrist tendon vibration on targeted upper-arm movements in poststroke hemiparesis. *Neurorehab Neurol Repair* 2011; 25:61-70. <https://doi.org/10.1177/1545968310378507>
 20. Maeda M, Mutai H, Toya Y, Maekawa Y, Hitai T, Katai S. Effects of peripheral nerve stimulation on paralysed upper limb functional recovery in chronic stroke patients undergoing low-frequency repetitive transcranial magnetic stimulation and occupational therapy: A pilot study. *HK J Occ Ther* 2020;33:31-11. <https://pubmed.ncbi.nlm.nih.gov/33815018/>
 21. Nasrallah FA, Mohamed AZ, Hong KY, Hwa SL, Yeow C, Lim JH. Effect of proprioceptive stimulation using a soft robotic glove on motor activation and brain connectivity in stroke survivors. *J Neural Eng* 2021;18:1. <https://doi.org/10.1088/1741-2552/ac456c>
 22. Seo NJ, Enders LR, Fortune A, Cain S, Vatinno AA, Schuster E, Ramakrishnan V, Feng W. Phase I safety trial: Extended daily peripheral sensory stimulation using a wrist-worn vibrator in stroke survivors. *Translational Stroke Research*, 2019;11(2), 204-213.
 23. Seo NJ, Woodbury ML, Bonilha L, Ramakrishnan V, Kautz SA, Downey RJ, Dellenbach BHS, Lauer AW, Roark CM, Landers LE, Phillips SK, Vatinno AA. Therabracelet stimulation during task-practice therapy to improve upper extremity function after stroke: A pilot randomized controlled study. *Physical Therapy*, 2019; 99(3), 319-328.
 24. Vatinno AA, Hall L, Cox H, et al: Using subthreshold vibratory stimulation during poststroke rehabilitation therapy. A case series. *OTJR* 2021; 42: <https://doi.org/10.1177/15394492211042275>
 25. Kaelin-Lang A, Luft AR, Sawaki L, Burstein AH, Sohn YH, Cohen LG. Modulation of human corticomotor excitability by somatosensory input. *The Journal of Physiology* 2002; 540(2), 623-633.
 26. Carrico C, Chelette KC, Westgate PM, Salmon-Powell E, Nichols L, Sawaki L. A randomized trial of peripheral nerve stimulation to enhance modified constraint-induced therapy after stroke. *Am J Phys Med Rehabil* 2016;95:397-406. doi: 10.1097/PHM.0000000000000476.
 27. Cordo P, Wolf S, Lou JS, Bogey R, Stevenson M, Hayes J, Roth E Treatment of severe hand impairment following stroke by combining assisted movement, muscle vibration, and biofeedback. *J Neurol Phys Ther* 2013 37:194-203. doi: 10.1097/NPT.0000000000000023.
 28. Lakshminarayanan K, Lauer AW, Ramakrishnan V, Webster JG, Seo NJ. Application of vibration to wrist and hand skin affects fingertip tactile sensation. *Physiological Reports* 2015; 3 2015. Article e12465.
 29. Carrico C, Chelette KC, Westgate PM, Salmon-Powell E, Nichols L, Fleischer A, Sawaki L. Nerve stimulation enhances task-oriented training in chronic, severe motor deficit after stroke: A randomized trial. *Stroke* 2016; 47, 1879-1884.
 30. Serrada I, Hordacre B, Hillier SL. Does Sensory Retraining Improve Sensation and Sensorimotor Function Following Stroke: A Systematic Review and Meta-analysis. *Front Neurosci* 2019; 13:402-15. <https://pubmed.ncbi.nlm.nih.gov/31114472/>
 31. Zhou SA, Huang Y, Jiao J, Hu J, Hsing C, Lai Z, Yang Y, Hu X. Impairments of cortico-cortical connectivity in fine tactile sensation after stroke. *J Neuroeng Rehab* 2021; 18:34. doi.org/10.1186/s12984-021-00821-7
 32. Bornheim, Stephen Transcranial direct current stimulation associated with physical-therapy in acute stroke patients - A randomized, triple blind, sham-controlled study. <https://doi.org/10.1016/j.brs.2019.10.019>. 2020
 33. Bustamante C, Brevis F, Canales S, Millón S, Pascual R Effect of functional electrical stimulation on the proprioception, motor function of the paretic upper limb, and patient quality of life: A case report. *J Hand Ther* 2016;29:507-514. doi: 10.1016/j.jht.2016.06.012. Epub 2016 Sep 21. PMID: 27665394 Review.
 34. Brown J. A., Lutsep H. L., Weinand M., Cramer S. C. (2006). Motor cortex stimulation for the enhancement of recovery from

- stroke: a prospective, multicenter safety study. *Neurosurgery* 58:464–473. 10.1227/01.Neu.0000197100.63931.04
35. Willis, A.W., Roberts, E., Beck, J.C. et al. Incidence of Parkinson disease in North America. *npj Parkinsons Dis.* 8, 170 (2022). <https://doi.org/10.1038/s41531-022-00410-y>
 36. De Lau LM, Breteler MM: Epidemiology of Parkinson's disease. *Lancet Neurol.* 5, 525–535 (2006).
 37. K.J. Pfeifer, A.J. Cook, J.K. Yankulova, B.J.P. Mortimer, E. Erickson-DiRenzo, R. Dhall, L. Montaser-Kouhsari, P.A. Tass: Clinical efficacy and dosing of vibrotactile coordinated reset stimulation in motor and non-motor symptoms of Parkinson's disease: a study protocol. *Frontiers in Neurology* 12:758481 (2021)
 1. 36.Tass PA: Vibrotactile Coordinated Reset Stimulation for the Treatment of Parkinson's Disease. *Neural Regeneration Research.* 17(7):1495-1497 (2021)
 38. Tramontano M, Morone G, De Angelis S, Conti L, Galeoto G, Grasso MG. Sensor-based technology for upper limb rehabilitation in patients with multiple sclerosis: A randomized controlled trial. *Restor Neurol Neurosci* 2020;38:333-41. <https://pubmed.ncbi.nlm.nih.gov/32925119/>
 39. Hammond, C., Bergman, H., and Brown, P. (2007). Pathological synchronization in Parkinson's disease: networks, models and treatments. *Trends Neurosci.* 2007;30, 357–364. doi: 10.1016/j.tins.2007.05.004
 40. Hall KD, Lifschitz J. Diffuse traumatic brain injury initially attenuates and later expands activation of the rat somatosensory whisker circuit concomitant with neuroplastic responses. *Brain Res* 2010; 1323:161-173. <https://doi.org/10.1016/j.brainres.2010.01.067>
 41. Lew HL, Weihing J, Myers PJ, Pogoda TK, Goodrich GL. Dual sensory impairment (DSI) in traumatic brain injury (TBI)—An emerging interdisciplinary challenge. *Neurorehabilitation* 2010; 26:213-222. <https://doi.org/10.3233/nre-2010-0557>
 42. Zheng Z, Dong X, Li Y, Gao W, Jiang R, Yue S, Zhou Z, Zhang J: Electrical stimulation improved cognitive deficits associated with traumatic brain injury in rats. *Brain and Behavior* 2017; 7:e00667. <https://doi.org/10.1002/brb3.667>
 43. Mamoei S, Hvid L, Jensen HB, Zijdwind I, Stenager E, Dalgas U: Neurophysiological impairments in multiple sclerosis—Central and peripheral motor pathways. *Acta Neurologica Scand* 2020, <https://doi.org/10.1111/ane.13289>
 44. Willingham TB, McCully K, Backus D: Skeletal Muscle Dysfunction in People With Multiple Sclerosis: A Physiological Target for Improving Physical Function and Mobility, *Archives of Physical Medicine and Rehabilitation*, 10.1016/j.apmr.2022.10.009, 104, 4, (694-706), (2023). Maffioletti NA, Dirks ML, Stevens-Lapsley J, McNeil CJ: Electrical stimulation for investigating and improving neuromuscular function in vivo: Historical perspective and major advances, *Journal of Biomechanics*, 10.1016/j.jbiomech.2023.111582, 152, (111582), (2023).
 45. Adkins DL, Hsu JE, Jones TA: Motor cortical stimulation promotes synaptic plasticity and behavioral improvements following sensorimotor cortex lesions. *Exp. Neurol.* 212: 14–28. 10.1016/j.expneurol.2008.01.031



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

Submit Manuscript

DOI:10.31579/2578-8868/267

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/neuroscience-and-neurological-surgery>