

Utilizing Amniotic Membrane Allograft as regenerative therapy for Treating Burns

Nishant Visvas Dumont¹, Ravi Kumar Chittoria^{2*}, Kanav Gupta³, Jacob Antony Chakiath³, Padmalakshmi Bharathi Mohan⁴ and Bharath Prakash reddy⁵

¹Junior Resident, Department of Oral & Maxillofacial Surgery, MGPGIDS, Puducherry.

²Professor & Registrar (Academic), Head of IT Wing and Telemedicine, Department of Plastic Surgery & Telemedicine, JIPMER, Pondicherry.

³General Surgery, Senior Resident, Department of Plastic Surgery, JIPMER, Puducherry.

⁴General Surgery, MCh Plastic Surgery, Assistant Professor, Department of Plastic Surgery, JIPMER, Puducherry.

⁵General Surgery, Senior Resident, Department of Plastic Surgery, JIPMER, Puducherry.

***Corresponding Author:** Ravi Kumar Chittoria, Senior Professor and Associate Dean (Academic), Head of IT Wing and Telemedicine, Department of Plastic, Surgery and Telemedicine, JIPMER. Pondicherry, India.

Received date: July 26, 2024; **Accepted date:** August 05, 2024; **Published date:** August 12, 2024

Citation: Nishant V. Dumont, Ravi K. Chittoria, K. Gupta, Jacob A. Chakiath and Padmalakshmi B. Mohan, et al. (2024), Utilizing Amniotic Membrane Allograft as regenerative therapy for Treating Burns, *J. Clinical Case Reports and Studies*, 5(6); DOI:10.31579/2690-8808/208

Copyright: ©, 2024, Ravi Kumar Chittoria. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Burn injuries are prevalent among all age groups and can result from thermal, scald, or electrical sources. Currently, various scaffolds are employed to enhance the healing process and minimize scar formation. Collagen serves as a scaffold, facilitating tissue regeneration and promoting the formation of new blood vessels. Additionally, scaffolds such as amniotic membrane aid in proper epithelialization and scar reduction, boasting unique anti-inflammatory and bacteriostatic properties. In our study, we utilized amniotic membrane allograft as a biological dressing for burn wounds on an adult patient's right thigh.

Key Words: amniotic membrane allograft; wound healing; regenerative therapy

Introduction

The process of wound healing is a biological response to various forms of injury, encompassing physical, chemical, mechanical, or thermal damage. This process typically involves several sequential phases: homeostasis, inflammation, proliferation/granulation, and remodeling/maturation [1]. However, deviations from the normal healing trajectory often result in stagnation during the inflammatory phase, particularly evident in cases of burns where standard healing mechanisms may be compromised [2]. In contemporary medicine, the utilization of scaffolds, whether natural or synthetic, has gained traction and recognition. An optimal scaffold should possess specific characteristics: appropriate physical and mechanical properties, a physiological background conducive to cellular adhesion, proliferation, and differentiation, high porosity, a favorable surface area to volume ratio, flexibility to conform to wound shapes, and ideally, biocompatibility and biodegradability [3]. Collagen, whether synthetic or natural, serves as a surrogate for the dermal matrix, facilitating epithelialization during wound healing [4]. Additionally, collagen degradation during this process contributes to neovascularization, thereby promoting angiogenesis. Amniotic membrane, a natural scaffold, exhibits distinctive properties such as anti-inflammatory, bacteriostatic, anti-fibrotic,

anti-scarring, and epithelialization-promoting qualities [6]. Its low immunogenicity and presence of progenitor cells make it an appealing choice for scaffold applications in wound healing. Silicone functions as a barrier, reducing mechanical friction and transepidermal water loss, factors associated with infection severity. In vitro studies suggest that silicone may modulate inflammatory growth factors implicated in fibrosis and support acute wound healing, including key inflammatory markers like TNF- α , TGF- β , IL-1, and IL-6.

Materials and Methods

This research was carried out at a Tertiary Care Centre within the Department of Plastic Surgery following approval from the department's ethical committee. Informed consent was duly obtained. The subject of this study was an eighty-two-year-old female who sustained accidental second-degree thermal burns affecting the right thigh. Despite being promptly taken to a nearby hospital, initial resuscitation was inadequate within 30 minutes. She subsequently developed blistering and swelling around the gluteal region the following day and presented to our center with a delay of 12 hours. Upon admission to the tertiary burn care unit, she received initial resuscitation involving intravenous fluids, analgesics, and prophylactic antibiotics. On the

fourth day post-burn, a three-layered scaffold dressing was prepared and applied to the deeper burn areas following dermabrasion-assisted tangential excision. This scaffold comprised sterile amniotic membrane, dry collagen sheet, and silicone sheet, with the amniotic membrane layer directly contacting the wound. The amniotic membrane (fig 1) used was obtained

from a freshly delivered placenta, thoroughly washed, and stored in an antibiotic solution in a refrigerator. The dressing remained in place for seven days, during which time the amniotic membrane was completely absorbed, and the silicone sheet layer was also removed (fig 2).



Figure 1: Figure showing amniotic membrane which was harvested from human placenta



Figure 2: Figure showing amniotic membrane which was harvested from human placenta being applied over the wound over the gluteal region

Results

Throughout both the intraoperative and postoperative phases, the patient experienced no notable events. Upon opening the dressing on the seventh day post-operation, substantial areas of re-epithelialization and healing were

observed. Complete healing of all second-degree superficial burn wounds was achieved without any complications or adverse effects noted throughout the entire procedure. BJWAT wound score improved from 32 at presentation to 25 after utilization of amniotic membrane regenerative therapy (fig 3 & 4).



Figure 3: Figure showing condition of the wound at time of presentation (BJWAT wound score - 32)



Figure 4: Figure showing condition of the wound after utilization of amniotic membrane regenerative therapy (BJWAT wound score -25)

Discussion

Partial-thickness burn injuries have the potential to heal spontaneously, whereas full-thickness burns typically necessitate skin grafting to achieve complete wound closure. Historically, split-thickness skin autograft has been the preferred method for closing excised full-thickness burn wounds [5]. However, patients with extensive burn injuries may face challenges due to limited availability of donor sites for autograft harvesting, prompting consideration of alternative options such as skin substitutes. Engineered skin substitutes offer temporary coverage for wounds until donor sites are

available for autografting or, if containing autologous cells, can provide permanent closure [7]. While the availability of permanent skin substitutes remains limited, advancements in tissue engineering are anticipated to yield improved models, thus enhancing the options for managing burn wounds. For instance, the commercially available Dermal Regeneration Template comprises a two-layered system. The outer layer, composed of a thin silicone film, serves as a protective barrier against infection and regulates heat and moisture loss. The inner layer, consisting of a collagen glycosaminoglycan (GAG) thermal layer, acts as a biodegradable template facilitating dermal tissue regeneration. The porous nature of the template aids in skin

regeneration by serving as a scaffold for tissue growth. Following regeneration of the dermal layer, the outer layer of the template is replaced with a thin epidermal skin graft, allowing for flexible, permanent skin regeneration and promoting faster wound healing with minimal scarring. Seeking to emulate this mechanism, we have developed an indigenous dermal regeneration scaffold using silicone sheet, dry collagen sheets, and amnion³. This cost-effective scaffold can be readily prepared and utilized, particularly in hospital settings within developing countries where the affordability of commercial regeneration templates may be a concern.

Conclusion

The efficacy of employing a cost-effective regenerative scaffold dressing based on amniotic membrane in treating second-degree scald burns has been demonstrated in this investigation. It accelerates the overall healing process of both superficial and deep second-degree wounds to less than a week, thereby reducing hospitalization duration and infection rates. Nevertheless, further extensive multicenter, double-blinded controlled research incorporating statistical analysis is warranted.

References

1. Haghpanah S, Bogie K, Wang X, Banks PG, Ho CH. (2006). Reliability of electronic versus manual measurement techniques. Arch Phys Med Rehabil. 87: 1396-1402.
2. Chitoria RK, Kumar P, Baljaj SP, Singh AK, Gupta DK. (2014). General clinical guidelines for wound management: Redefining acronym SWCR. Journal of SWCR.;7(1): 2-7.
3. Shetty R, Sreekar H, Lamba S, Gupta AK. (2012). A novel and accurate technique of photographic wound measurement. Indian J Plast Surg.; 45(2): 425- 429.
4. Majeske C. (1992). Reliability of wound surface area measurement. PhyTher.; 72: 138-141.
5. Mayrovitz HN, Soontupe LB. (2009). Wound area by computerized planimetry of digital images. Adv Skin Wound Care.; 22: 222-229.
6. Pandey S, Chitoria RK, Mohapatra DP, et al. (2016). Application of digital planimetry: A novel technique of wound measurement in diabetic foot ulcer. Dermatology International.; 21: 343-331.
7. Pires IM, Garcia NM. (2015). Wound area assessment using mobile application. Proceedings of the International Conference on Biomedical Electronics and Devices - SmartMedDev, International Conference on Biomedical Electronics and Devices; 12- 15; pp. 271-282.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here: [Submit Manuscript](#)

DOI:10.31579/2690-8808/208

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/journal-of-clinical-case-reports-and-studies>