

Biosilk bilayered scaffold Application in Wound bed Preparation of Thermal Burns Wound

Nithin D ¹, Ravi Kumar Chittoria ^{2*}, Kanav Gupta ³, Nagarjun S Ghatti ⁴

¹MBBS, MS General Surgery Junior resident Department of General Surgery JIPMER Pondicherry.

²MNAMS, FRCS (Edin), DSc, PhD (Plastic Surgery) Professor & Registrar (Academic) Head of IT Wing and Telemedicine Department of Plastic Surgery & Telemedicine JIPMER Pondicherry.

³MBBS, MS DNB General Surgery Senior Resident Department of Plastic Surgery JIPMER Puducherry.

⁴MBBS Junior Resident Department of Orthopaedics Aarupudai Veedu Medical College Puducherry.

***Corresponding Author:** Ravi Kumar Chittoria, MBBS, MS General Surgery Junior resident Department of General Surgery JIPMER Pondicherry.

Received date: June 19, 2024; Accepted date: July 15, 2024; Published date: September 16, 2024

Citation: Nithin D, Ravi Kumar Chittoria, Kanav Gupta, Nagarjun S Ghatti, (2024), Biosilk bilayered scaffold Application in Wound bed Preparation of Thermal Burns Wound, *Clinical Research and Clinical Trials*, 10(5); DOI:10.31579/2693-4779/214

Copyright: © 2024, Ravi Kumar Chittoria. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract:

Haemostasis, inflammation, proliferation, and remodelling are the steps that comprise the complex biological process of wound healing. Chronic wounds often remain static in the inflammatory stage of wound healing beyond the expected duration and resist healing. Biosilk or biomodified silk dressing is believed to aid in accelerating the wound healing process. The combination bio functionalised silk matrix with silver oxide and *Centella asiatica*, a medicinal plant with wound healing properties, has been found to be effective in managing chronic non-healing wounds.

Keywords: biosilk; thermal burns; scaffold; wound healing; wound management

Introduction

Wounds healing is a complex biological process which comprises of haemostasis, inflammation, proliferation, and remodelling. Cells that are involved in this process include neutrophils, macrophages, lymphocytes, keratinocytes, fibroblasts, and endothelial cells.[1] A chronic wound is one that does not heal in the expected sequence and in a predictable length of time, or one that does not heal within three months.[2] Chronic wounds often remain static in the inflammatory stage of wound healing beyond the expected duration. The predominant issue that patients with chronic ulcers face is persistent pain. In the recent years, biological dressings have gained popularity and have been found to benefit patients with chronic wounds. One such dressing substance being explored is biosilk containing silver oxide and

Centella asiatica, a medicinal plant. Its use is thought to enhance the wound healing process.

Materials and Methods

An 82 years old gentleman, with no comorbidities, presented with complaints of thermal burns of 35% BSA by cylinder blast. On examination of the left upper limb: 2nd degree superficial to deep burns on the forearm on the dorsal aspect with multiple blisters.

Wound at admission (Figure 1)



Figure 1: Wound at presentation with BJWAT score- 36

and on Day 3 of admission when Biosilk was applied (Figure 2).



Figure 2: Wound on Day 3 of admission on the left upper arm.

Wound on 10th day of admission on the left upper arm (Figure 3 and 4).



Figure 3 and 4: Wound with BJWAT 24 score on 10th day of admission on the left upper arm.

Result

At the end biosilk dressing, there was accelerated wound healing noted of the raw area of right thigh with healthy granulation tissue.

Discussion

Wound healing to be adequate requires rapid haemostasis, appropriate inflammation, mesenchymal cell differentiation, proliferation, and migration to the wound site, angiogenesis, prompt re-epithelialization and proper synthesis, cross-linking, and alignment of collagen to provide strength to the healing tissue.[4] Growth factors such as transforming growth factor (TGF)-platelet-derived growth factor (GF), fibroblast growth factor (FGF), and epidermal growth factor (EGF) play a crucial role in facilitating the above-mentioned steps of wound healing

The aim of this study was to explore the effects of use of biosilk in wound healing. The combination used here included silk, Centella asiatica extract, and silver oxide, each of which carry important properties that affect wound healing. Due to the unique properties of silk, such as, great mechanical strength, outstanding biocompatibility, and the capacity to modify the structural and morphological aspects of silk proteins, they represent a new class of sophisticated biomaterials. It has proven to be a beneficial material in biomedical engineering applications such as skin, bone, and vascular grafts. [5-9] Schneider et al conducted a study regarding use of silk mats incorporated with epidermal growth factor (EGF), for the promotion of wound healing processes and was concluded that there was an increase in wound closure by the epidermal tongue by 90%.[10]

Centella asiatica facilitate the wound healing process in both incision and burn wounds. Amongst a variety of extracts obtained for this plant, ascitic acid in the ethyl acetate extract seemed to be the most active component for healing the wound.[11] It has been reported that 1% C asiatica extract cream improves wound healing of chronic ulcer.[12] It works by inhibiting inflammation, inducing collagen synthesis, promoting angiogenesis inducing vasodilation, reducing wound oxidative stress in addition to promoting cellular growth and proliferation in injured tissues which may be related to

growth factors such as endothelial growth factor, fibroblast growth factor and vascular endothelial growth factor.[11]

The third component of the ointment was silver oxide which has great antimicrobial and bactericidal properties which is due to reaction of the highly charged silver ion (Ag⁺) to the negatively charged particles such as proteins, DNA, RNA, and chloride ions.[13]

Conclusion

In this study, we have understood to role of biosilk or biomodified silk in enhancement of wound healing and the properties of various properties of biosilk that facilitate the same. To confirm the effectivity of biosilk in healing of wound, it is required to conduct larger randomised control trials.

References

1. Guo, & and L a Dipatro. (2010). "Factors affecting wound healing" Journal of dental research vol. 89,3: 219-29
2. Mustoe T. (2006). Dermal ulcer healing Advances in understanding Tissue repair and ulcer/wound healing molecular mechanisms, therapeutic targets and future directions Paris, France EUROCONFERENCES; Archived from the original (PDF) on October 27,
3. Snyder, Robert J. (2005). Treatment of nonhealing ulcers with allografts. Clin Dermatol.:23:388-395.
4. Mathies D, Linke J-C, Wattel F. (2006), non-healing wounds. In: Handbook on hyperbaric medicine, Mathieu DE, editor. Netherlands: Springer, 401-427
5. Dal Pya I, Chharini A, Boschi A. Frodd G. Armato U (2006). Novel dermoepidermal equivalents on silk fibroin based formic acid-crosslinked three-dimensional nonwoven devices with prospective applications in human tissue engineering/regeneration/repair. Int J Mol Med:18-241-247.
6. Kim HJ, Kim UJ, Leisk GG, Bayan C, Georgakoudi I, Kaplan Dt. (2007). Bone regeneration on macroporous aqueous-derived silk threedimensional scaffolds. MacromolBiosci,7:643-655.

7. Kim KH, Jeong L, Park HN, Shin SY, Park WH, Lee SC, et al. (2005). biological efficacy of silk fibroin nanofiber membranes for guided bone regeneration. *J Biotechnol*,120-327-339.
8. Motta A, Migliaresi C, Faccioni F, Torricelli P, Fini M, Giardino R. (2004). Fibroin hydrogels for biomedical applications: preparation, characterization and in vitro cell culture studies. *J BiomatSciPolym*,15.851-864.
9. Jin HJ, Fridrikh SV, Rutledge GC, Kaplan DL. (2002). Electrospinning Bombyx mori silk with poly (ethylene oxide). *Biomacromolecules*: 3:1213-1219.
10. Schneider A, Wang XY, Kaplan DL, Garlick JA, Egles C. (2009). Biofunctionalized electrospun silk mats as a topical bioactive dressing for accelerated wound healing *Actabiomater*. 5(7):2570-2578
11. Somboonwong, J, Kankaisre, M., Tantisira, B. & Tantisira, M. H. (2012). Wound healing activities of different extracts of *Centella asiatica* in incision and burn wound models: an experimental animal study. *BMC complementary and alternative medicine*, 12,
12. Kosalwatna S, Shaipanich C, Bhanganada K. (1988). The effect of one percent *Centella asiatica* chronic ulcers. *Siriraj Hosp Gaz. cram* 40-435-460
13. Murphy, Patrick S, and Gregory RD Evans. (2012). "Advances in wound healing a review of current wound healing products. *Plastic surgery international*: 190436.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

[Submit Manuscript](#)

DOI:10.31579/2693-4779/214

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/clinical-research-and-clinical-trials>