

## Hurdle Concept and The Role of Lactic Acid Bacteria

Zakari David Adeiza <sup>1,2\*</sup>, Kareem Sarafadeen Olateju <sup>2</sup>, Bello Kizito Eneye <sup>1,3</sup>, Audu Godwin Amoka <sup>1</sup>, Aliyu Abdulbasit Anoze <sup>4</sup>, Muhammed Abdulsamad Adeiza <sup>1</sup>, Osazuwa Christopher Olojamarin <sup>5</sup>, Emurotu Marvelous Olubunmi <sup>1</sup>

<sup>1</sup>Department of Microbiology, Kogi State University (Prince Abubakar Audu University), PMB 1008, Anyigba, Kogi State, Nigeria

<sup>2</sup>Department of Microbiology, College of Biosciences, Federal University of Agriculture, PMB 2240, Abeokuta, Ogun State, Nigeria

<sup>3</sup>Department of Medical Microbiology and Parasitology, School of Medical Sciences, Universiti Sains, Malaysia

<sup>4</sup>Department of Chemistry, Prince Abubakar Audu University, PMB1008, Anyigba, Kogi State, Nigeria

<sup>5</sup>Department of Microbiology, Adekunle Ajasin University, Akungba, Ondo State, Nigeria

**Corresponding author:** Zakari David Adeiza, 1Department of Microbiology, Kogi State University (Prince Abubakar Audu University), PMB 1008, Anyigba, Kogi State, Nigeria.

**Received date:** July 08, 2024; **Accepted date:** August 13, 2024; **Published date:** August 22, 2024

**Citation:** Zakari D. Adeiza, Kareem S. Olateju, Bello K. Eneye, Audu G. Amoka, Aliyu A. Anoze, et al, (2024), Hurdle Concept and The Role of Lactic Acid Bacteria, *J. Nutrition and Food Processing*, 7(10); DOI:10.31579/2637-8914/248

**Copyright:** © 2024, Zakari David Adeiza. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract:

The hurdle idea is a food preservation technique that includes building up a number of barriers to prevent microbes from growing and prolong the shelf life of perishable goods. The idea of using lactic acid bacteria (LAB), which are essential for food preservation, is one that is significant. Through a variety of processes, including fermentation, competitive exclusion, metabolite generation, biofilm formation, nutritional competition, and metabolic activity, LAB support the hurdle concept. Lactic acid bacteria (LAB) ferment glucose to produce lactic acid, which results in an acidic environment that prevents infections and spoiling germs from growing. Additionally, LAB generates antimicrobial substances such as hydrogen peroxide, organic acids, and bacteriocins, which work by competitively excluding harmful germs from growing. Furthermore, metabolites of lactic acid bacteria (LAB) such as acetic acid, diacetyl, and carbon dioxide enhance the flavor of fermented foods and serve as organic preservatives. Moreover, LAB have the ability to create biofilms that shield them from antimicrobial agents and environmental stressors, increasing their survival and persistence in food matrices. By means of nutritional competition, LAB outcompetes pathogenic or spoilage bacteria in the consumption of available resources, hence restricting their growth. In addition, LAB produces enzymes and antibacterial substances, among other metabolic activities that further impede the growth of infections.

**Key words:** lactic acid bacteria; diacetyl; biofilms; acetic acid; food matrices

### Introduction

The hurdle idea is a method of food preservation that combines the use of several non-lethal barriers to stop pathogenic and spoilage bacteria from growing and surviving in food. Physical, chemical, and biological variables are some of the obstacles. Because it minimizes the use of preservatives and other potentially dangerous chemicals while preserving the nutritional value, flavor, and texture of food, this method of food preservation has gained popularity in recent years (Panel, 2022).

The hurdle concept and the function of lactic acid bacteria (LAB) are important factors in food preservation since they guarantee the durability and safety of perishable food items. The hurdle concept is using several barriers or hurdles to prevent microbes from growing and prolong food shelf life. Important players in this notion are lactic acid bacteria, a varied group of bacteria recognized for their capacity to create lactic acid through fermentation.

Fundamentally, the hurdle concept seeks to establish a setting in which microbes find it difficult to endure and multiply. The total microbial load in food is reduced by combining several barriers, such as temperature regulation, pH adjustment, water activity reduction, and preservatives, increasing the stability and safety of food products. Because of their special qualities and contributions to preventing the proliferation of pathogens and spoilage bacteria, LAB play a major role in this framework (Nura *et al.*, 2016).

LAB, which include a variety of species like *Lactobacillus*, *Lactococcus*, *Leuconostoc*, and *Streptococcus*, are ubiquitous in nature. These microorganisms are frequently present in fermented foods and give them their unique tastes, textures, and scents. Food products have a lower pH because LAB convert carbohydrates into lactic acid through metabolic processes. This environment's acidity hinders the development of spoiling microorganisms as well as the spread of harmful bacteria (Malheiros *et al.*, 2015).

Furthermore, the many metabolites and chemicals that LAB create further impede the growth of unwanted microbes. For example, they create natural preservatives such as hydrogen peroxide, organic acids, and bacteriocins, which are antibacterial substances. These substances show inhibitory actions against pathogens and spoilage bacteria, adding another line of defense against microbial contamination (Nura *et al.*, 2016).

LAB also support the hurdle notion by outcompeting other microbes for resources and available space through a process known as competitive exclusion. Because of their better nutrient utilization skills, LAB are able to inhibit the growth of harmful or spoilage bacteria and efficiently absorb available resources. LAB can also create biofilms, which increase their survival and persistence in food matrices by providing defense against environmental stressors and antimicrobial agents (Kaya *et al.*, 2019).

Food manufacturers are able to increase the safety, sensory qualities, and shelf life of perishable foods by utilizing LAB and the hurdle concept. The application of LAB in a range of culinary items, including cheese, yogurt, sauerkraut, kimchi, and sourdough bread, demonstrates their adaptability and significance in the maintenance and evolution of unique flavors in these goods (Leal-Sánchez *et al.*, 2002).

### History and Evolution:

Van Valen initially presented the idea of hurdle technology in the early 1960s. He suggested using a number of low-level physical and chemical barriers to stop microorganisms from growing in food. Subsequently, the hurdle theory has been improved and broadened to encompass more biological and physical variables. Over time, the hurdle notion has grown and changed to encompass a wider variety of physical, chemical, and biological elements that are involved in food preservation. Van Valen first offered the idea in the early 1960s, suggesting the use of several low-level chemical and physical barriers to stop microorganisms from growing in food. Since then, other components and their combinations have been added to the hurdle concept, which has been improved and broadened (Panel, 2022).

### Scientific Principles:

The hurdle concept is founded on the scientific idea that a variety of low-level stressors working together can limit or completely stop microorganisms in food from growing and surviving. These stress variables, which collectively prevent the development and survival of bacteria, include temperature, pH, water activity, preservatives, and antimicrobial compounds. Because of their combined effect, less preservative and antibacterial agent is needed than would be the case if these components were used independently (Parada *et al.*, 2007).

### Practical Applications:

A wide range of food products, including meats, dairy products, fruits, and vegetables, have been effectively preserved using the hurdle idea. The hurdle idea is frequently used in the meat industry to produce cooked and cured meats, which require the use of heat, nitrite or nitrate, low pH, and low water activity. Similar to this, lactic acid bacteria and low pH are utilized in the dairy industry to create fermented foods like kefir, yogurt, and cheese using the hurdle idea. The idea of the hurdle is also applied to the preservation of fruits and vegetables, wherein low pH, heat, and refrigeration are combined to keep the produce fresh and unspoiled (Panel, 2022).

In order to stop the growth of microorganisms and lengthen the shelf life of food, the hurdle idea in food preservation refers to a technique that combines several barriers or elements. The steps involved in applying the hurdle principle to food preservation are as follows:

### Steps involved in the use of hurdle concept in food preservation

1. Hurdle Selection: The first stage is to determine which hurdles will be utilized to keep the food fresh. A number of variables, including

temperature, pH, water activity, chemical preservatives, packaging, and altered environment, might operate as hurdles (Malheiros *et al.*, 2015).

2. Analysis of Microbial Targets: It's critical to know which particular microbes are most likely to proliferate in the food that is being preserved. The right barriers can be chosen with the aid of this analysis (Kaya *et al.*, 2019).

3. Determining obstacle Parameters: Each obstacle requires a set of parameters that are determined based on the study. For example, the type of food and the microorganisms present can be used to establish the target temperature, pH level, water activity, or concentration of preservatives (Mahrous *et al.*, 2013).

4. Combination of Hurdles: Combining several hurdles is a crucial component of the hurdle concept. Various obstacles combine to prevent microbial development. Finding the ideal combination to optimize the preservation effect is crucial (Kaya *et al.*, 2019).

5. Application of Barriers: After defining the barriers and their limitations, the meal is subjected to them. This may entail procedures such as heat treatment (such as pasteurization or sterilization), pH level adjustments, preservative additions, packing under modified atmospheres, or high-pressure processing (Oladapo *et al.*, 2014).

6. Monitoring and Control: To make sure that the barriers are successful in stopping microbial development, the preservation process must be watched over. The effectiveness of the hurdle system can be confirmed through routine testing, such as microbiological analysis can be performed to verify the efficacy of the hurdle system (Nura *et al.*, 2016).

7. Storage and Distribution: Following the preservation procedure, the food can be kept and shared in suitable settings, keeping in mind the particular obstacles that were employed. To keep the barriers functional, handling, packing, and storage temperatures must all be followed (Leistner, 2000).

8. Determining Shelf Life: Over time, quality and safety factors are assessed to ascertain how long a preserved product will last. This entails assessing any deterioration or spoiling through chemical analysis, microbiological testing, and sensory evaluation (Panel, 2022).

### Factors used in food preservation

**Temperature:** A key element in the preservation of food is temperature. Food microbes are frequently destroyed or rendered inactive by heat treatment, and food microbiological quality is greatly dependent on temperature control. In order to attain the required degree of preservation, hurdle technology employs combinations of heat treatments at various temperatures (Oladapo *et al.*, 2014).

**Acidity/pH:** Another significant variable that can be controlled for food preservation is acidity or pH. Because some microbes are pH sensitive, reducing a food product's pH can prevent them from growing and surviving. The pH levels of food products have been adjusted by the use of organic acids as well as other acids including citric, lactic, and acetic acids (Nura *et al.*, 2016).

**Water Activity:** A food product's water molecule availability is gauged by its water activity (aw), with a lower aw signifying a smaller amount of available water molecules. Because certain microbes require high aw levels to thrive, microbial development and spoiling can be prevented by decreasing aw through drying, salting, or the use of other chemicals (Oladapo *et al.*, 2014).

**Preservatives:** To stop microbes from growing, preservatives are frequently employed in food preservation. With hurdle technology, the same amount of microbial inhibition that would normally need larger doses of single preservatives is achieved by using many preservatives, frequently in low quantities (Leal-Sánchez *et al.*, 2002).

**Antimicrobial Agents:** Materials that can stop the growth of bacteria include synthetic and natural antimicrobial agents like bacteriocins. The food business makes extensive use of bacteriocins, which are proteins made by specific bacteria that either kill or stop the growth of other bacteria (Kaya *et al.*, 2019).

**Biological Processes:** Dairy products, meats, and vegetables are frequently preserved using biological processes like fermentation and the use of starter cultures. During fermentation, carbohydrates are broken down by microbes, most often lactic acid bacteria, which result in the production of organic acids and other substances that aid in the regulation of microbial development and enhance the quality of food (Malheiros *et al.*, 2015).

### Various food products that employs hurdle technology

1. **Cooked and Cured Meats:** Haze technique is frequently used to preserve foods including sausages, deli meats, and cooked ham. This calls for the application of several techniques, including low pH (from acids or fermentation), low water activity (from drying or salt addition), heat treatment (cooking or smoking), and the use of nitrite or nitrate as preservatives (Nura *et al.*, 2016).

2. **Fermented Dairy Products:** Using hurdle technology, fermented dairy products include kefir, cheese, and yogurt. A mixture of lactic acid bacteria ferment lactose to produce lactic acid, which in turn creates an acidic environment that prevents the growth of pathogenic microbes and spoiling (Kaya *et al.*, 2019).

3. **Fruit and Vegetable Preservation:** To retain quality and increase shelf life, fresh fruits and vegetables can be preserved using hurdle technology. To reduce microbial and enzymatic activity, methods like refrigeration, modified atmosphere packaging (MAP), and temperature and humidity management are used (Panel, 2022).

4. **Pickles and Sauerkraut:** The hurdle principle is used to preserve pickled vegetables, such as cucumbers and sauerkraut. The flavors and textures are enhanced and rotting is prevented by the combination of high salt concentrations, low pH (from fermentation or vinegar), and chilling (Oladapo *et al.*, 2014).

5. **Shelf-Stable Ready-to-Eat Meals:** The manufacturing of shelf-stable, ready-to-eat meals frequently uses hurdle technology. To preserve safety and increase shelf life, a variety of preservation procedures are used, including low pH, low water activity, vacuum packaging, thermal processing (canning), and oxygen removal methods (Nura *et al.*, 2016).

6. **Bakery Products:** Hinderneath technology can also be used to preserve baked items such as bread, cakes, and pastries. Preservatives, temperature control, low water activity (from sugar or reduced water content), and altered packaging methods are some of the strategies used to keep product quality and stop microbiological deterioration (Malheiros *et al.*, 2015).

### How are the factors of food preservation employed in hurdle concept

In order to stop microbiological development and lengthen the shelf life of food, the hurdle concept in food preservation combines a number of hurdles, such as temperature, pH, water activity, moisture content, chemical preservatives, packaging, and changed environment. Each of these obstacles is used as follows:

1. **Temperature:** One of the biggest obstacles to food preservation is temperature. Microorganisms can be killed by high temperatures or their growth might be slowed down by low temperatures (Leistner, 2000). The type of food, the microorganisms present, and the preservation method all play a role in determining the temperature. For instance, pasteurization entails heating the food to a precise temperature (often between 72°C and 75°C) in order to kill the majority of the bacteria, whereas sterilization entails heating the food to a higher temperature (sometimes beyond 100°C) in order to eradicate all germs (Kaya *et al.*, 2019).

2. **pH:** Another significant barrier to food preservation is pH. Many bacteria are inhibited in their growth by acidity. Food can be made more acidic and less conducive to microbial development by adjusting its pH level using organic acids such as citric acid, lactic acid, or acetic acid (Walter, 2008).

3. **Water activity:** The quantity of water that is available for microbial growth in a food product is known as water activity (aw). Since microorganisms require water to develop, lowering the water activity may prevent them from doing so. Drying, adding sugar or salt, or utilizing other dehydrating agents can all help achieve this (Yi *et al.*, 2013).

4. **Moisture content:** Moisture content is a significant barrier to food preservation and is correlated with water activity. It is possible to lower the food's moisture content by drying, dehydration, or desiccation processes, which prevent microbial growth (Zúñiga *et al.*, 1993).

5. **Chemical preservatives:** To stop or slow down the growth of germs in food, chemicals such organic acids, salt, nitrites, and sulfites are frequently utilized as barriers. The kind of microorganisms present and the food product being preserved determine the chemical preservative's type and concentration (Ulloa *et al.*, 2008).

6. **Packaging:** Food preservation is greatly aided by packaging. By establishing a barrier between the food and the environment, it can stop microbiological growth. Using materials for airtight packaging will help achieve this. Additionally, some packaging methods can lower the oxygen content and produce an anaerobic environment that prevents the growth of bacteria (Ulloa *et al.*, 2008).

7. **Modified atmosphere:** It is possible to alter the environment around food in order to prevent the growth of microorganisms. This can be accomplished by changing the air's composition in the packaging material to include a mix of nitrogen, oxygen, and carbon dioxide. Packaging with a modified environment can be helpful for fresh items like meat products and fruits and vegetables (Van Geel-Schutten *et al.*, 1998).

In order to stop microbes from growing, food preservation experts use a combination of barriers, such as temperature, pH, water activity, moisture level, chemical preservatives, packaging, and a modified atmosphere. The application of each of these barriers depends on the particular food product, the kind of microorganisms present, and the preservation method being employed (Ulloa *et al.*, 2008).

### How does hurdle concept help in extending the shelf life of perishable foods

1. **Microbial Growth Inhibition:** The hurdle idea establishes an environment that prevents the growth of microbes by using a variety of hurdles, including temperature, pH, water activity, moisture content, chemical preservatives, packing, and changed atmosphere. This prolongs the shelf life by preventing deterioration and the growth of foodborne germs (Walter, 2008).

2. **Combination Effect:** The barriers in the hurdle idea cooperate with one another to increase their efficacy against microbes. When combined, for instance, low temperature, packaging in a modified environment, and chemical preservatives can significantly suppress microbial development more than when any one of these barriers is used alone (Yi *et al.*, 2013).

3. **Reduction of Water Activity:** Since microorganisms need water to thrive, the barrier idea produces an environment that is adverse to microbial growth by reducing the water activity by methods including drying, dehydration, or the addition of salt or sugar. This slows down the rate of spoiling, extending the shelf life of perishable items (Zúñiga *et al.*, 1993).

4. **Preservation of Quality Attributes:** The hurdle idea aids in maintaining the quality characteristics of perishable foods in addition to preventing microbial development (Kaban and Kaya, 2008). For example, keeping



the food at the right temperature, utilizing packaging with a modified environment, and managing moisture content can all help to efficiently preserve the meal's sensory qualities by reducing oxidative reactions, delaying physical deterioration, and slowing down enzymatic activity (Jang *et al.*, 2006).

5. **Safety Assurance:** By preventing the formation of foodborne germs, the hurdle concept not only increases the shelf life of perishable goods but also guarantees their safety. By controlling harmful germs, a safer product is provided for consumers through the use of proper chemical preservatives and pH modification, among other challenges (Kaban and Kaya, 2008).

6. **Shelf Life Prediction and Extension:** The hurdle idea makes it possible to better understand the shelf life of perishable foods through regular monitoring, shelf life prediction models, and quality assessments (Kadariya *et al.*, 2014). The items' shelf life can be increased by modifying and optimizing the hurdle characteristics, which will decrease food waste and boost profitability.

Perishable goods can be kept safe from infection and deterioration by using the hurdle idea in food preservation, and ensuring an extended shelf life while maintaining their quality and safety (Kaya *et al.*, 2019).

#### Chemical preservatives commonly used in hurdle concept

1. **Benzoates:** Common preservatives include potassium and sodium benzoate. They work well against some bacteria, molds, and yeasts. Pickles, fruit juices, salad dressings, fizzy drinks, and pickles all frequently include benzoates (KP *et al.*, 2016).

2. **Sorbates:** Common preservatives include calcium and potassium sorbate. They work well against some bacteria, yeasts, and molds. Sorbates are frequently found in wine, cheese, baked products, and dried fruits (Leal-Sánchez *et al.*, 2002).

3. **Sulfites:** Strong preservatives with antibacterial and antioxidant qualities, sulfites include sulfur dioxide, sodium sulfite, and potassium metabisulfite. They are frequently found in wine, dried fruits, canned vegetables, and seafood and are efficient against a variety of bacteria (Leistner, 2000).

4. **Nitrites and Nitrates:** To stop the growth of *Clostridium botulinum* and other dangerous bacteria, processed meat products like bacon, ham, and sausages frequently contain potassium and sodium nitrate. Additionally, nitrates and nitrites contribute to the distinct flavor, color, and shelf life of cured meats (Mahrous *et al.*, 2013).

5. **Organic Acids:** Preservatives such as lactic acid, malic acid, acetic acid, and citric acid work well. They produce an acidic environment that prevents microbial growth by lowering the pH of the meal. Organic acids are frequently found in drinks, dressings, sauces, and dairy products (Malheiros *et al.*, 2015).

6. **Propionates:** To prevent the formation of mold and germs, calcium and sodium propionate are frequently added to bread and other baked foods. They aid in extending the shelf life of baked goods and are efficient against rotting microbes (Mandal *et al.*, 2008).

7. **Parabens:** Two common food preservatives found in baked goods, confections, and beverages are methyl and propylparaben. They work well against germs, molds, and yeasts (McDonald *et al.*, 1987).

8. **EDTA, or ethylenediaminetetraacetic acid,** is a chelating chemical that helps maintain the quality and freshness of some foods, particularly those that are processed and canned. It binds to metal ions that can catalyze unwanted reactions, preventing the deterioration of texture, flavor, and color (Mead *et al.*, 1999).

#### Role of Lactic Acid Bacteria in the preservation of fermented dairy products through hurdle technology.

They contribute to the safety, flavor, and shelf life of these products by performing the following functions in preservation process:

1. **Acidification:** Lactose, the naturally occurring sugar in milk, is fermented by LAB to produce lactic acid (Mokoena *et al.*, 2016). The dairy product's pH is lowered via acidification, which also produces an acidic environment that prevents the growth of harmful bacteria and spoiling. The product's flavor and texture are also preserved by the higher acidity (Naidu *et al.*, 1999).

2. **Competitive Exclusion:** Because LAB can grow in the low pH environment they produce, they have an edge over many other microbes (Nura *et al.*, 2016). LAB quickly acidify the product, making it uninhabitable for more potentially hazardous bacteria, yeasts, and molds. By limiting the proliferation of spoilage germs, this competitive exclusion reduces the likelihood of foodborne diseases (Oladapo *et al.*, 2014).

3. **Production of Preservative Compounds:** During fermentation, LAB create a variety of organic compounds, including lactic acid, hydrogen peroxide, and bacteriocins. By preventing the growth of harmful microbes and deterioration, these substances function as organic preservatives (Onwuakor *et al.*, 2014). LAB specifically produces proteinaceous substances called bacteriocins, which have antibacterial effect against bacteria that are closely related to one another or compete with it (Oppegård *et al.*, 2010).

4. **Development of Flavor and Aroma:** LAB gives fermented dairy products their distinct flavors and fragrances. Lactic acid and other chemicals such as acetic acid, diacetyl, and volatiles are produced during fermentation by lactose breakdown by lactic acid bacteria (Panel, 2022). These substances improve the product's sensory qualities by imparting to it the characteristically acidic, sour, or creamy flavors of fermented dairy products (Parada *et al.*, 2007).

5. **Shelf Life Extension:** Fermented dairy products have a longer shelf life due to the interaction of acidity, competition, and antibacterial chemicals produced by LAB (Perez *et al.*, 2014). The risk of spoiling and waste is decreased because the controlled growth of LAB prevents the growth of bacteria that cause spoiling and helps maintain product quality for a longer amount of time (Rajilić-Stojanović and De Vos, 2014).

#### How different strains of lactic acid bacteria impact the flavor and texture of fermented dairy products

1. ***Streptococcus thermophilus:*** Yogurt is typically made using this common culture of bacteria. It produces lactic acid by fermenting lactose, which lowers pH and coagulates milk proteins to give yogurt its sour flavor and creamy texture (Saxena *et al.*, 2009). Exopolysaccharides, which are also produced by *S. thermophilus*, enhance the end product's viscosity and texture (Sidhu and Nehra, 2019).

2. ***Lactobacillus bulgaricus:*** This common yogurt culture is also essential in giving yogurt its characteristically sour taste (Hati *et al.*, 2013). *L. bulgaricus* coagulates milk proteins and helps give yogurt its creamy texture, working in tandem with *S. thermophilus*. But *L. bulgaricus* is also capable of producing volatile substances like diacetyl and acetaldehyde, which give yogurt its distinct flavor (Herbel *et al.*, 2013).

3. ***Lactobacillus delbrueckii subsp. bulgaricus:*** Because of the particular bacterial culture utilized, this strain of bacteria is utilized to produce Bulgarian yogurt, which differs slightly from regular yogurt (Alkema *et al.*, 2016). Bulgarian yogurt's smooth, creamy texture is produced by the fermentation of lactose into lactic acid by *Lactobacillus delbrueckii subsp. bulgaricus*, which also lowers pH. It also yields a unique flavor that is sometimes characterized as being somewhat acidic and tasting like fresh cheese (Aditya and Nida, 2015).

4. ***Lactococcus lactis:*** This bacteria is used to make a variety of cheeses, such as feta, Gouda, and cheddar. Cheese gets its distinctive pores and

texture from the fermentation of lactose into lactic acid and carbon dioxide. The creation of the cheese flavor is also aided by the bacteria's breakdown of the milk proteins (Alakomi *et al.*, 2000).

5. Bifidobacteria: Some yogurts and other fermented dairy products are made with the help of bifidobacteria. They have a reputation for having possible health benefits, including as boosting immunity and enhancing digestive health (Aljewicz and Cichosz, 2017). These goods might have a smooth, creamy texture and a mild, somewhat sweet flavor thanks to the impact of bifidobacteria (Alkema *et al.*, 2016).

These are just a few examples, but the specific bacterial strains used, as well as the fermentation conditions and processing techniques, can significantly impact the final product characteristics (Anurag Chaturvedi *et al.*, 2013). The choice of bacterial culture used in the fermentation process can greatly influence the sensory attributes, texture, and nutritional value of fermented dairy products (Arora *et al.*, 2009).

### Some specific food products that employs lactic acid bacteria in its hurdle concept

Lactic acid bacteria (LAB) can be employed in several specific food products using the hurdle concept to inhibit the growth of spoilage microorganisms and pathogens (Bazhal *et al.*, 2003). Here are some examples:

1) Cheese: The fermentation process in the making of cheese is started with LAB. Lactic acid is produced by LAB during this process, which acidifies the cheese curd and makes it uninhabitable for pathogens and bacteria that cause spoiling (Bourdichon *et al.*, 2019). The main application of some LAB strains, such as *Lactobacillus helveticus* and *Lactococcus lactis*, is in the making of cheese, which gives different types of cheese their distinct flavor, texture, and scent (De Vuyst and Leroy, 2007).

2) Yogurt: Lactic acid bacteria (LAB) are essential for the fermentation of lactose, a milk sugar, into lactic acid, which lowers pH and imparts a tangy flavor to yogurt (Devi *et al.*, 2013). *Lactobacillus bulgaricus* and *Streptococcus thermophilus* are two particular LAB strains that are frequently employed in the manufacturing of yogurt; during fermentation, they also create vital vitamins and nutrients (Diep *et al.*, 2009).

3) Sauerkraut: LAB use lactic acid fermentation to turn cabbage into sauerkraut, which has a sour, tart flavor that is typical of fermented foods (Dimov *et al.*, 2005). The LAB *Lactobacillus plantarum* and *Leuconostoc mesenteroides* are in charge of fermenting sauerkraut. Because lactic acid production lowers the pH of sauerkraut, spoilage bacteria cannot thrive there, extending the product's shelf life (Djadouni and Kihal, 2012).

4) Kimchi: A fermented vegetable food that goes through lactic acid fermentation, kimchi is similar to sauerkraut (Elayaraja *et al.*, 2014). *Leuconostoc mesenteroides*, *Lactobacillus kimchii*, and *Lactobacillus plantarum* are LAB species that aid in the fermentation of kimchi. In addition to extending the product's shelf life, LAB give kimchi its distinct flavor and color (Field *et al.*, 2018).

5) Sourdough bread: To produce sourdough bread, LAB are used in the fermentation process, which gives the dough a sour taste. *Lactobacillus sanfranciscensis* and *Lactobacillus brevis* are the main LAB strains used in sourdough fermentation; they enhance the bread's shelf life and aid in the formation of taste and fragrance molecules (Ganguly, 2013).

### Challenges and solutions to hurdle concept

Combining a number of barriers or hurdles to stop the growth of microbes and increase the shelf life of food items is known as the hurdle concept in food preservation. Although it provides practical preservation solutions, there are certain implementation-related difficulties.

The following are some typical obstacles and possible ways to overcome them in relation to the hurdle concept:

1. Microbial Adaptation: The efficiency of individual barriers can be decreased when microorganisms learn to adapt and become resistant to one or more obstacles found in the food environment. Reduced shelf life and impaired safety may result from this (Goh and Philip, 2015).

Solution: Evaluate and modify the set of obstacles utilized in the food preservation procedure on a regular basis. The effectiveness of barriers can be maintained and microbial adaptation can be inhibited by rotating between different preservation procedures (Goldstein *et al.*, 2015).

2. Synergistic Effects: When several obstacles are employed together, they may have a synergistic effect that produces a preservation outcome that is greater than when they are utilized alone. It can be difficult to figure out the best combination and dosage of obstacles, though (Gordon, 2016).

Solution: To determine the best combination and concentration of obstacles for a given food product, conduct extensive research and trial. Extensive testing and collaboration with food scientists can yield important insights on optimization and synergy (Güllüce *et al.*, 2013).

3. Quality and Sensory Changes: During the preservation process, some obstacles like intense heat or potent preservatives might change the food products' sensory characteristics and quality. Consumer acceptability may be impacted by this (Gunathilake, 2005).

Solution: Look into different barriers or altered processing methods that can minimize detrimental effects on sensory attributes while still achieving microbial control. For instance, replacing chemical preservatives with natural antimicrobials or applying a gentler heat treatment (Hati *et al.*, 2013).

4. Cost and Implementation: The barrier concept may need to be implemented with additional testing, equipment, and monitoring, which could be costly and logistically difficult, particularly for smaller food enterprises (Herbel *et al.*, 2013).

Solution: Make the best use of the tools and resources already in place while looking at more affordable options for implementing the barrier. Partnering with academic institutions or business associations can facilitate access to resources, shared knowledge, and funding.

5. Regulatory Compliance: Regulations and rules pertaining to the installation of hurdle technology, including the concentrations of permissible hurdles, may vary based on the geographical region (Hu *et al.*, 2010).

### Recommendation

Maintain awareness of the particular rules and policies in the intended market. Consult a professional, interact with regulatory bodies, and make sure local food safety laws are being followed.

### Conclusion

The hurdle idea is a flexible and successful method of food preservation that combines a number of non-lethal elements to stop microbes from growing and surviving in food. Improved food safety, longer shelf life, and a decrease in the need of preservatives and other potentially dangerous chemicals are just a few advantages of using this strategy. Several food products, including as meats, dairy products, fruits, and vegetables, have been produced effectively using the hurdle idea. The obstacle idea is probably going to become a more crucial tactic in the food business as consumer desire for naturally occurring, minimally processed goods grows. The hurdle notion has evolved over time to encompass a variety of physical, chemical, and biological elements that combine to produce successful food preservation techniques. Combining these variables enables the use of preservative and antibacterial agent dosage

reductions with the same degree of microbial inhibition. The barrier concept will keep changing as new methods of preservation, as well as new components and procedures, are created to assist satisfy the increasing consumer desire for naturally occurring, minimally processed foods.

## References

- Aditya Pundhir, A. P. and Nida Murtaza, N. M. (2015). Hurdle technology-an approach towards food preservation.
- Alakomi, H. L., Skytta, E., Saarela, M., Mattila-Sandholm, T., Latva-Kala, K. and Helander, I. M. (2000). Lactic acid permeabilizes gram-negative bacteria by disrupting the outer membrane. *Applied and environmental microbiology*, 66(5), 2001-2005.
- Aljewicz, M. and Cichosz, G. (2017). Influence of probiotic (*Lactobacillus acidophilus* NCFM, *L. paracasei* LPC37, and *L. rhamnosus* HN001) strains on starter cultures and secondary microflora in Swiss-and Dutch-type cheeses. *Journal of food processing and preservation*, 41(6), e13253.
- Alkema, W., Boekhorst, J., Wels, M. and van Hijum, S. A. (2016). Microbial bioinformatics for food safety and production. *Briefings in Bioinformatics*, 17(2), 283-292.
- Anurag Chaturvedi, A. C., Sujatha, V., Ramesh, C. and Babu, J. D. (2013). Development of shelf stable intermediate moisture carrot (*Daucus carota*) shreds using radiation as hurdle technology.
- Arora, A. K., Mahajan, S., Gupta, P. and Kapoor, S. S. (2009). Is chemical safety to food hazardous?—Dangers of food preservatives. *Journal of Indian Academy of Forensic Medicine*, 31(4), 419-422.
- Bazhal, M. I., Ngadi, M. O., Raghavan, G. S. V. and Smith, J. P. (2003). Minimal processing of foods using hurdle technologies. In *CSAE/SCGR meeting*, Montreal, Quebec.
- Bourdichon, F., Laulund, S. and Tenning, P. (2019). Inventory of microbial species with a rationale: a comparison of the IDF/EFFCA inventory of microbial food cultures with the EFSA Biohazard Panel qualified presumption of safety. *FEMS microbiology letters*, 366(5), fnz048.
- De Vuyst, L. and Leroy, F. (2007). Bacteriocins from lactic acid bacteria: production, purification, and food applications. *Journal of molecular microbiology and biotechnology*, 13(4), 194-199.
- Devi, M., Rebecca, L. J. and Sumathy, S. (2013). Bactericidal activity of the lactic acid bacteria *Lactobacillus delbreuckii*. *Journal of Chemical and Pharmaceutical Research*, 5(2), 176-180.
- Diep, D. B., Mathiesen, G., Eijsink, V. G. H. and Nes, I. F. (2009). Use of lactobacilli and their pheromone-based regulatory mechanism in gene expression and drug delivery. *Current pharmaceutical biotechnology*, 10(1), 62-73.
- Dimov, S., Ivanova, P. and Harizanova, N. (2005). Genetics of bacteriocins biosynthesis by lactic acid bacteria. *Biotechnology & Biotechnological Equipment*, 19(sup2), 4-10.
- Djadouni, F. and Kihal, M. (2012). Antimicrobial activity of lactic acid bacteria and the spectrum of their biopeptides against spoiling germs in foods. *Brazilian Archives of Biology and Technology*, 55, 435-444.
- Elayaraja, S., Annamalai, N., Mayavu, P. and Balasubramanian, T. (2014). Production, purification and characterization of bacteriocin from *Lactobacillus murinus* AU06 and its broad antibacterial spectrum. *Asian Pacific journal of tropical biomedicine*, 4, S305-S311.
- Field, D., Ross, R. P. and Hill, C. (2018). Developing bacteriocins of lactic acid bacteria into next generation biopreservatives. *Current Opinion in Food Science*, 20, 1-6.
- Ganguly, S. (2013). Biologically viable methods for food preservation: A review. *Res J Chem Environ*, 1, 01-02.
- Goh, H. F. and Philip, K. (2015). Purification and characterization of bacteriocin produced by *Weissella confusa* A3 of dairy origin. *PLoS one*, 10(10), e0140434.
- Goldstein, E. J., Tyrrell, K. L. and Citron, D. M. (2015). *Lactobacillus* species: taxonomic complexity and controversial susceptibilities. *Clinical Infectious Diseases*, 60(suppl\_2), S98-S107.
- Gordon, A. (Ed.). (2016). *Food Safety and Quality Systems in Developing Countries: Volume II: Case Studies of Effective Implementation*. Academic Press.
- Güllüce, M., Karadayı, M. and Barış, Ö. (2013). Bacteriocins: promising natural antimicrobials. *Local Environ*, 3(6), 1016-1027.
- Gunathilake, K. D. P. P. (2005). Application of hurdle technique to preserve fresh scraped coconut at ambient and refrigerated storage.
- Hati, S., Mandal, S. and Prajapati, J. B. (2013). Novel starters for value added fermented dairy products. *Current Research in Nutrition and Food Science Journal*, 1(1), 83-91.
- Herbel, S. R., Vahjen, W., Wieler, L. H. and Guenther, S. (2013). Timely approaches to identify probiotic species of the genus *Lactobacillus*. *Gut Pathogens*, 5, 1-13.
- Hu, C. B., Malaphan, W., Zendo, T., Nakayama, J. and Sonomoto, K. (2010). Enterocin X, a novel two-peptide bacteriocin from *Enterococcus faecium* KU-B5, has an antibacterial spectrum entirely different from those of its component peptides. *Applied and environmental microbiology*, 76(13), 4542-4545.
- Jang, J. D., Seo, G. H., Lyu, E. S., Yam, K. L. and Lee, D. S. (2006). Hurdle effect of vinegar and sake on Korean seasoned beef preserved by sous vide packaging. *Food control*, 17(3), 171-175.
- Kaban, G. and Kaya, M. (2008). Identification of lactic acid bacteria and Gram-positive catalase-positive cocci isolated from naturally fermented sausage (sucuk). *Journal of food science*, 73(8), M385-M388.
- Kadariya, J., Smith, T. C. and Thapaliya, D. (2014). *Staphylococcus aureus* and staphylococcal food-borne disease: an ongoing challenge in public health. *BioMed research international*, 2014.
- Kaya, H. İ., Özel, B. and Şimşek, Ö. (2019). A natural way of food preservation: Bacteriocins and their applications. *Health and safety aspects of food processing technologies*, 633-659.
- KP, S., PV, K., PK, B., RC, R., PB, D. and AK, S. (2016). Production and characterization of bacteriocin produced by *Lactobacillus viridescence* (NICM 2167). *Brazilian Archives of Biology and Technology*, 59, e16150518.



30. Leal-Sánchez, M. V., Jiménez-Díaz, R., Maldonado-Barragán, A., Garrido-Fernández, A. and Ruiz-Barba, J. L. (2002). Optimization of bacteriocin production by batch fermentation of *Lactobacillus plantarum* LPCO10. *Applied and Environmental Microbiology*, 68(9), 4465-4471.
31. Leistner, L. (2000). Basic aspects of food preservation by hurdle technology. *International journal of food microbiology*, 55(1-3), 181-186.
32. Mahrous, H., Mohamed, A., Abd El-Mongy, M., El-Batal, A. I. and Hamza, H. A. (2013). Study bacteriocin production and optimization using new isolates of *Lactobacillus* spp. isolated from some dairy products under different culture conditions. *Food and Nutrition Sciences*, 4(03), 342.
33. Malheiros, P. S., Sant'Anna, V., Todorov, S. D. and Franco, B. D. (2015). Optimization of growth and bacteriocin production by *Lactobacillus sakei* subsp. *sakei*2a. *Brazilian Journal of Microbiology*, 46, 825-834.
34. Mandal, V., Sen, S. K. and Mandal, N. C. (2008). Optimized culture conditions for bacteriocin production by *Pediococcus acidilactici* LAB 5 and its characterization.
35. McDonald, L. C., McFeeters, R. F., Daeschel, M. A. and Fleming, H. (1987). A differential medium for the enumeration of homofermentative and heterofermentative lactic acid bacteria. *Applied and environmental microbiology*, 53(6), 1382-1384.
36. Mead, P. S., Slutsker, L., Dietz, V., McCaig, L. F., Bresee, J. S., Shapiro, C. and Tauxe, R. V. (1999). Food-related illness and death in the United States. *Emerging infectious diseases*, 5(5), 607.
37. Mokoena, M. P., Mutanda, T. and Olaniran, A. O. (2016). Perspectives on the probiotic potential of lactic acid bacteria from African traditional fermented foods and beverages. *Food & Nutrition Research*, 60(1), 29630.
38. Naidu, A. S., Bidlack, W. R. and Clemens, R. A. (1999). Probiotic spectra of lactic acid bacteria (LAB). *Critical reviews in food science and nutrition*, 39(1), 13-126.
39. Nura, A., Araihi, C. C. and Abu, J. O. (2016). Effects of chemical hurdles and packing materials on microbial load and bacterial distribution in kilishi under ambient storage. *Int J Res Technol*, 5, 306-315.
40. Nura, A., Chukwuma, A. C. and Abu, J. O. (2016). CRITICAL REVIEW ON PRINCIPLES AND APPLICATIONS OF HURDLE TECHNOLOGY IN FOOD PRESERVATION. *Annals: Food Science & Technology*, 17(2).
41. Oladapo, A. S., Akinyosoye, F. A. and Abiodun, O. A. (2014). The inhibitory effect of different chemical food preservatives on the growth of selected food borne pathogenic bacteria.
42. Onwuakor, C. E., Nwaugo, V. O., Nnadi, C. J. and Emetole, J. M. (2014). Effect of varied culture conditions on crude supernatant (bacteriocin) production from four *Lactobacillus* species isolated from locally fermented maize (ogi). *American Journal of Microbiological Research*, 2(5), 125-130.
43. Oppegård, C., Rogne, P., Kristiansen, P. E. and Nissen-Meyer, J. (2010). Structure analysis of the two-peptide bacteriocin lactococcin G by introducing D-amino acid residues. *Microbiology*, 156(6), 1883-1889.
44. Panel, E. B. (2022). Update of the list of QPS-recommended biological agents intentionally added to food or feed as notified to EFSA 15: suitability of taxonomic units notified to EFSA until September 2021. *EFSA Journal*, 20(1), e07045.
45. Parada, J. L., Caron, C. R., Medeiros, A. B. P. and Soccol, C. R. (2007). Bacteriocins from lactic acid bacteria: purification, properties and use as biopreservatives. *Brazilian archives of Biology and Technology*, 50, 512-542.
46. Perez, R. H., Zendo, T. and Sonomoto, K. (2014). Novel bacteriocins from lactic acid bacteria (LAB): various structures and applications. *Microbial cell factories*, 13, 1-13.
47. Rajilić-Stojanović, M. and De Vos, W. M. (2014). The first 1000 cultured species of the human gastrointestinal microbiota. *FEMS microbiology reviews*, 38(5), 996-1047.
48. Saxena, S., Mishra, B. B., Chander, R. and Sharma, A. (2009). Shelf stable intermediate moisture pineapple (*Ananas comosus*) slices using hurdle technology. *LWT-Food Science and Technology*, 42(10), 1681-1687.
49. Sidhu, P. K. and Nehra, K. (2019). Bacteriocin-nanoconjugates as emerging compounds for enhancing antimicrobial activity of bacteriocins. *Journal of King Saud University-Science*, 31(4), 758-767.
50. Ulloa, J. A., Escalona, H. and Díaz, L. (2008). Colour behaviour on mango (*Mangifera indica*) slices self-stabilized in glass jars by hurdle technology during storage. *African Journal of Biotechnology*, 7(4).
51. Van Geel-Schutten, G. H., Flesch, F., Ten Brink, B., Smith, M. R. and Dijkhuizen, L. J. A. M. (1998). Screening and characterization of *Lactobacillus* strains producing large amounts of exopolysaccharides. *Applied Microbiology and Biotechnology*, 50, 697-703.
52. Walter, J. (2008). Ecological role of lactobacilli in the gastrointestinal tract: implications for fundamental and biomedical research. *Applied and environmental microbiology*, 74(16), 4985-4996.
53. Yi, H., Han, X., Yang, Y., Liu, W., Liu, H., Zhang, Y. and Ma, F. (2013). Effect of exogenous factors on bacteriocin production from *Lactobacillus paracasei* J23 by using a resting cell system. *International journal of molecular sciences*, 14(12), 24355-24365.
54. Zacharof, M. P. and Lovitt, R. W. (2012). Bacteriocins produced by lactic acid bacteria a review article. *Apcbee Procedia*, 2, 50-56.
55. Zúñiga, M., Pardo, I. and Ferrer, S. (1993). An improved medium for distinguishing between homofermentative and heterofermentative lactic acid bacteria. *International journal of food microbiology*, 18(1), 37-42.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

**Submit Manuscript**

**DOI:10.31579/2637-8914/248**

**Ready to submit your research? Choose Auctores and benefit from:**

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/nutrition-and-food-processing>