

Adenomyoepithelioma of the Breast: A Rare Case Report from B.P Koirala Memorial Cancer Hospital

Deepak Singh Mouni*, Sulochana Shrestha, Ankita Adhikari

Breast Onco Surgery Unit, Sugical Oncology Depatment, B.P. Koirala Memorial Cancer Hospital, Bharatpur, Nepal.

***Corresponding Author:** Deepak Singh Mouni., Breast Onco Surgery Unit, Sugical Oncology Depatment, B.P. Koirala Memorial Cancer Hospital, Bharatpur, Nepal.

Received Date: 08 July, 2024 **Accepted Date:** 22 July, 2024 | **Published Date:** 16 August, 2024

Citation: Deepak S Mouni., Shrestha S., Adhikari A., (2024), Adenomyoepithelioma of the Breast: A Rare Case Report from B.P Koirala Memorial Cancer Hospital., Journal of Clinical Surgery and Research, 5(6) DOI:10.31579/2768-2757/130

Copyright: © 2024, Deepak S Mouni. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract:

Adenomyoepithelioma is a rare, benign breast neoplasm characterized by the proliferation of both epithelial and myoepithelial cells within the breast lobules and ducts. Although it is most commonly seen in salivary glands and skin, Hamper reported the first case in the breast in the 1970 (1). While this benign tumor is predominantly found in older females, rare cases have been identified in young women as well. Adenomyoepithelioma is notorious for local recurrence, and malignant metastasis has been reported (2). Adenomyoepithelioma is very rare case and in 10 years period, out of 19000 cases treated in OPD, we have encountered only 2 cases of adenomyoepithelioma, one was referred case & another one was diagnosed at our center.

keywords: adenomyoepithelioma; asymptomatic; benign mass; case report; surgical excision

Introduction

A 35 years old female presented to the out patient department with the chief complaint of painless lump in the upper outer quadrant of the right breast that had been present for the past 1 and half months. The patient had no history of pain, nipple discharge, skin changes or increase in the size of the mass. The patient had no personal history of oral contraceptive pills (OCP) intake, is non smoker, non alcoholic and no family history of breast tumors. On examination, a 2 x 1.5 cm, firm, well defined, round, partially mobile lump with a smooth margin was palpated over the upper outer quadrant of the breast, with no skin changes. Ultrasonography revealed an ill-defined, heterogeneous lesion measuring 20 x 14 x 13mm in the fibroglandular tissue of the right breast at the 7 to 8 o'clock position, with minimal vasculature noted within the lesion. A fine-needle aspiration cytology (FNAC) was performed and showed features consistent with a benign proliferative disease of the breast, specifically a fibroadenoma. The patient underwent wide local excision under intravenous anesthesia and was discharged on the same day with no postoperative complications. The histopathology report revealed adenomyoepithelioma, with immunohistochemistry markers showing P63 diffusely immunoreactive in the myoepithelial component, S-100 diffusely immunoreactive in the myoepithelial component, and HMWCK immunoreactive in the epithelial component. The proliferation index Ki-67 was 1-5%. ER was patchy immunoreactive in 10-15% of neoplastic cells. The patient has been advised to undergo strict six-monthly follow-up with ultrasound

of the breast and axilla to monitor for local recurrence and metastatic spread.

Discussion

Adenomyoepithelioma can be classified into three types: tubular, lobulated, and spindle cell. Tubular adenomyoepithelioma has an ill-defined margin, similar to tubular adenoma. Lobulated

adenomyoepithelioma presents as nests of myoepithelial cells surrounding compressed epithelial-lined spaces. Spindle cell adenomyoepithelioma shows sparse epithelial-lined spaces that resemble leiomyomalike cells. (3)

Diagnosis:


Adenomyoepitheliomas typically presents as a single, palpable, well-circumscribed, firm mass, with dimensions of up to 8 cm. Microscopic satellite lesions may be present in the periphery of the tumor at times, but it is mostly located in the central portion. Rarely, adenomyoepitheliomas may also present in phyllodes tumor or fibroadenoma. Pain and nipple discharge are rare symptoms associated with adenomyoepitheliomas. (15)

Radiology evaluation:


Radiology imaging is not specific and cannot distinguish between benign and malignant lesions (2). Ultrasonography (USG) shows that

adenomyoepitheliomas are typically solid or have a combined solid and cystic appearance. On mammography, they present as a lobulated dense mass with a partially indistinct margin, with or without calcifications. Malignant adenomyoepitheliomas show irregular shapes with spiculated margins on mammography. (14)

Pathology evaluation:



**Myoepithelial
cells
surrounding
glands**



**Compressed
tubules**

Microscopic description of adenomyoepithelioma: (10)

Immuno Histo chemistry (IHC):

IHC is an important tool in the diagnosis of adenomyoepithelioma. The panel of markers, including p63, S-100, and HMWCK.P63 is a transcription factor that is expressed in the nuclei of myoepithelial cells and is a very sensitive and specific marker for identifying these cells. In adenomyoepithelioma, P63 staining is typically positive and shows a consistent rim of staining around the epithelial cells, which can help to distinguish it from other breast tumors. Ki-67 proliferation index is also useful in predicting the recurrence rate of the tumor. ER, PR, and HER-2 are usually negative in malignant adenomyoepithelioma. (6) (7) (9)(12)

Differential diagnosis:

- **Sclerosing adenosis:** Sclerosing adenosis is a condition where there is an increase in the number of epithelial glands along with hardening of the surrounding stroma, which can lead to an abnormal shape of the glands. It is usually not associated with the formation of a well-defined mass, and there is no significant presence of myoepithelial cells.
- **Intraductal papilloma:** Refers to a papillary growth consisting of an increase in epithelial cells along with fibrovascular cores, without a significant presence of myoepithelial cells.
- **Invasive carcinoma:** Malignant growth of epithelial cells with clear evidence of invasion.
- Absence of myoepithelial markers such as p63/p40, SMMHC, calponin, and CK5.
- **Nipple adenoma:** Growth of epithelial cells within the collecting ducts of the nipple. There is no significant presence of myoepithelial cells.
- **Tubular adenoma:** well-defined border and the proliferation of tubules with a single layer of myoepithelial cells surrounding the epithelial component without expansion of the myoepithelial component characterize the tubular adenoma. (16)

Treatment

No specific guidelines have been established for the treatment of adenomyoepithelioma. However, wide local excision with negative margins is recommended due to the local recurrence nature of the tumor. Some cases of malignant adenomyoepithelioma have also been

FNAC and core needle biopsy are also rarely diagnostic due to the heterogeneity of the tumor, with many cases falsely diagnosed as fibroadenoma in the literature. However, the presence of tightly aggregated glands arranged in compact nodules and prominent clear cell or spindle cell myoepithelium can be clues to the diagnosis. (4) (8) (13)

reported in the literature, where mastectomy and sentinel lymph node biopsy are recommended. (10) (11)

Prognosis

Benign adenomyoepitheliomas generally have a good prognosis but require close monitoring and followup. In cases of local recurrence, radiotherapy has been used with positive results. However, malignant adenomyoepithelioma has a poor prognosis due to its low-grade invasiveness, high recurrence rate, and resistance to chemotherapy. Malignant tumors are more likely to metastasize via hematogenous spread to distant organs such as the brain, liver, and lungs rather than through the lymphatic system. (10) (11)

Conclusion

Adenomyoepitheliomas (AMEs) are rare benign breast tumors that should be considered in the differential diagnosis of solid breast lumps. While imaging features are not specific, suspicion of benign or malignant nature can be raised through radiologic examination. FNAB is often not diagnostic.

Currently, there are no established guidelines for treating either benign or malignant AME. Surgical excision with negative margins is recommended for both types of AME due to the high recurrence rate for benign tumors and the aggressiveness of malignant ones. Close monitoring and follow-up are necessary for benign cases, while malignant cases have a poor prognosis and are characterized by low grade, invasiveness, and high recurrence rate. Chemotherapy is not effective for treating malignant AME, and metastasis occurs through hematogenous spread to organs such as the brain, liver, and lungs.

References

1. Hamperl H. (1970). The myoepithelium (myoepithelial cells). Normal state; regressive changes; hyperplasia; tumors. *Curr Top Pathol.*; 53:161-220.
2. El-Helou E, Terro JJ, Kansoun A, Neaime GR, Mochairefa H, Ismail N, Naccour J, Zaarour M, Alam H. Breast adenomyoepithelioma, a case report. *Int J Surg Case.*
3. Intagliata E, Gangi S, Trovato C, Vecchio R, Strazzanti A. Benign adenomyoepithelioma of the breast: Presentation of two rare cases and review of literature. *International Journal of Surgery Case*

4. Reports. Pradhan S, Yadav RP, (2011). Adenomyoepithelioma of breast, case report, *Journal of pathology of Nepal*,
5. Krystel-Whittemore M, Wen HY. Adenomyoepithelioma. *PathologyOutlines.com*.
6. McLaren BK, Smith J, Schuyler PA, Dupont WD, Page DL. (2005). Adenomyoepithelioma: Clinical, Histologic, and Immunohistologic Evaluation of a Series of Related Lesions. *Am J Surg Pathol*; 29:1294-1299.
7. Malcolm MH. (2011). Adenomyoepithelioma of the breast: a review stressing its propensity for malignant transformation. *J Clin Pathol*; 64:477-484
8. Tamura G, Monma N, Suzuki Y, et al. (1993). Adenomyoepithelioma (myoepithelioma) of the breast in a male. *Hum Pathol*; 24:678-681
9. Tavassoli F, Soares J. (2003). Myoepithelial lesions. In: Tavassoli FA, Stratton MR, editors. WHO Classification of Tumors: Pathology and Genetics, Tumors of the Breast and Female Genital Organs. Lyon: IARC, pp86-7
10. Rakha E, Tan PH, Ellis I, Quinn C. (2021). Adenomyoepithelioma of the breast: a proposal for classification. *Histopathology* ;79(4):465-479.
11. Parikh P, Jameel Z, Falcon S, Rosa M, Kiluk J, et al. (2021). Adenomyoepithelioma of the breast: Case series and literature review. *Clin Imaging*; 75:157-164.
12. Kulka J, Madaras L, Floris G, Lax SF. (2022). Papillary lesions of the breast. *Virchows Arch* ;480(1):65-84.
13. Tay TKY, Tan PH. (2021). Papillary neoplasms of the breast- reviewing the spectrum. *Mod Pathol*. 34(6):1044-1061.
14. AlQurashi M, Abdel Hadi M, Binammar AA, Al Muhanna A, Kussaibi H, et al. (2022). Adenomyoepithelioma of the Breast: A Report of 3 Cases. *Am J Case Rep.*;23: e936070.
15. Xu J, Tang X, Iida Y, Fuchinoue F, Kusumi T, et al. (2016). Masuda S. Adenomyoepithelioma with carcinoma of the breast: A report of two cases and a review of the literature. *Pathol Res Pract.*;212(2):130-134.
16. Hayes, M. M. (2011). Adenomyoepithelioma of the breast: a review stressing its propensity for malignant transformation. *Journal of Clinical Pathology*, 64(6), 477-484.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

Submit Manuscript

DOI: [10.31579/2768-2757/130](https://doi.org/10.31579/2768-2757/130)

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://www.auctoresonline.org/journals/journal-of-clinical-surgery-and-research>