

Prenatal Diagnosis of Idiopathic Infantile Arterial Calcinosis – A Case Report on A Rare Fatal Condition

Sukanya Karunanidhi^{1*}, Sindhujha Sekar², Nirmala Jaget³

Department of Obstetrics and Gynecology Sri Venkateshwaraa Institute of Medical sciences and Research Centre, Puducherry.

***Corresponding Author:** Sukanya Karunanidhi, Department of Obstetrics and Gynecology Sri Venkateshwaraa Institute of Medical sciences and Research Centre, Puducherry.

Received Date: 12 December 2024; Accepted Date: 18 December 2024; Published Date: 26 December 2024

Citation: Sukanya K, Sindhujha Sekar, Nirmala Jaget, (2024), Prenatal Diagnosis of Idiopathic Infantile Arterial Calcinosis A Case Report on A Rare Fatal Condition. J. Obstetrics Gynecology and Reproductive Sciences, 8(9) DOI:10.31579/2578-8965/246

Copyright: © 2024 Sukanya K. This is an open-access article distributed under the terms of The Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction

Idiopathic infantile arterial calcinosis also known as Generalised arterial calcinosis of infancy (GACI) is a rare genetic disorder causing extensive premature calcification of major vessels. About 200 cases have been reported till date as per literature (1) and only less than 10 cases have been reported to be diagnosed during prenatal period by ultrasound (6). Hyperechogenicity of large vessels in the antenatal ultrasound helps in the prenatal diagnosis of this disease. Patient with positive history of IIAC in index pregnancy, unexplained death during neonatal or early infancy period can undergo molecular testing for diagnosis in early gestation. Genetic counselling and pre-implantation genetic diagnosis can be offered to these parents for their future pregnancies.

Case presentation

A 31-year-old Gravida 3, Para 2, living 1 and abortion 1 presented at 17 weeks of gestation with her chorionic villous sampling report showing mutation in ENPP1 gene at exon 9 in homozygous condition. She is a known case of hypothyroidism and her NT scan was normal. She had previous one intrauterine death at 34 weeks of gestation and the autopsy showed mutation in ENPP1 gene following which both the parents went for genetic counselling and underwent karyotyping. Both the parents had mutation in ENPP1 gene at exon 9 on chromosome 6q in heterozygous state. Hence chorionic villous sampling was done for this current pregnancy. It was reported as mutation in the similar variant in homozygous state. Patient was advised for medical termination of pregnancy. In spite of explaining the poor prognosis and very fewer evidence of success with bisphosphonate therapy, both the parents want to continue the pregnancy. Anomaly scan showed multiple echogenic lesions in the left ventricle.



Figure 1: Hyperechogenicity of the vessel walls seen in the anomaly scan

Fetal echocardiography showed diffuse calcification in papillary muscle tip, TV annulus, Left ventricle, crux, descending aorta, abdominal aorta extending upto its bifurcation.




Figure 2: Calcifications seen in the descending aorta extending upto abdominal aorta

After fetal medicine expert and neonatologist opinion, patient opted for medical termination of pregnancy.

Discussion

Idiopathic infantile arterial calcinosis is an autosomal recessive disorder with fatal outcome. It was first described by Bryant and white in 1901 (1, 2). Approximately 85% of cases are caused by loss of function mutation in ectonucleotide pyrophosphatase/ phosphodiesterase -1 (ENPP1) gene in chromosome 6q23 and causes death within the first 6 months of life (1,5). The association between IIAC and ENPP1 has been explained by Rutsch et al in 2003(2). ENPP1 is expressed on fibroblasts, osteoblasts and hepatocytes and is responsible for pyrophosphohydrolase activity. It produces inorganic pyrophosphate (Ppi) which prevents the deposition of calcium hydroxyapatite crystals in the arteries (7). The mutation of this gene leads to deposition of calcium along the internal elastic membrane of medium and large arteries of cardiac and renal system, accompanied by fibrous thickening of the intima which causes luminal narrowing.

During the insertion of an intrauterine device (IUD), stretching of the cervical internal os is the most painful step. The second most painful steps are the placement of the vulsellum, uterine sounding, and IUD insertion [16]. The classic approach to IUD insertion involves stretching the cervical os twice by introducing the uterine sound and then the IUD inserter, which increases the pain. However, in our study, we introduced the IUD inserter into the uterine cavity only once, which may result in less pain. Additionally, the plastic nature of the IUD inserter, when compared to the metal sound, exerts lesser tissue trauma, which could also cause less or no pain.

Diagnosis:



Figure 3: Diffuse calcifications of the left ventricle seen in the fetal echocardiography

Conclusion

IIAC is a rare disorder and has poor fetal outcome. Hence, careful and detailed history taking of risk factors and serial fetal echocardiography during antenatal period should be carried out. If IIAC is suspected, molecular

Most of the diagnosis is made during autopsy (8). Prenatal diagnosis can be made if risk factors like consanguinity, positive family history or history of IIAC in the index pregnancy along with serial fetal echocardiogram. Few cases have reported association of polyhydramios and fetal hydrops with IIAC (5, 7). Even though arterial biopsy is the gold standard for diagnosis of IIAC, DNA analysis for ENPP1 is a good antenatal screening test. IIAC should be suspected in patient with risk factors like consanguinity, positive family history, previous IIAC in the index pregnancy, unexplained early neonatal death and hyperechogenicity of the vessel wall (5). Once the suspicion is made, serial ultrasound examinations and amniotic fluid testing to detect DNA mutation should be advised.

Treatment:

Very few studies have reported successful treatment with bisphosphonate therapy, an analogue of PPI (2, 8). It alters the calcium balance by interfering with hydroxyapatite crystal formation and inhibits the calcium deposition in the vessels. Postnatal treatment with bisphosphonate has poor prognosis and the most common side effect of prolonged use of bisphosphonates leads to severe skeletal dysplasia similar to rickets (4). More studies should be done to study the outcome of medical therapy with bisphosphonates. Therefore, once the diagnosis of IIAC is made, couples should be offered genetic counselling and preimplantation genetic diagnosis should be advised.

testing of DNA for mutation will help in early diagnosis of the disease. Genetic counselling and preimplantation genetic diagnosis may be helpful in the future pregnancies.



Figure 4: Fetus along with the placenta expelled after termination of pregnancy

References

1. Mulcahy, C.H., Mone, F., McAuliffe, F.M. et al. (2019). Antenatal diagnosis of idiopathic infantile arterial calcification (IIAC): a single centre experience and review of the literature. *J Congenit Heart Dis* 3, 1 <https://doi.org/10.1186/s40949-018-0022-1>
2. Rutsch F, Böyer P, Nitschke Y, (2008). the GACI Study Group, et al. Hypophosphatemia, hyperphosphaturia, and bisphosphonate treatment are associated with survival beyond infancy in generalized arterial calcification of infancy. *Circ Cardiovasc Genet*. 1:133–140.
3. Votava-Smith JK, Pitukcheewanont P, Randolph LM, Chmait RH. (2017). Generalized arterial calcification in a recipient twin: discordant fetal hemodynamics result in differing phenotypes in monozygotic twins with an ABCC6 mutation. *Fetal Diagn Ther*. 41:234–236.
4. Yunfeng L, Tongyan H, Jing W and Xiaomei T (2021). Case Report: A Novel Genetic Mutation Causes Idiopathic Infantile Arterial Calcification in Preterm Infants. *Front. Genet*. 12:763916. doi: 10.3389/fgene.2021.763916
5. Swati B. Mune, P.A. Gadgil, Gulfam B. Shaikh, et al. (2023). Fugare, Idiopathic infantile arterial calcification: An autopsy evaluation, *IJH Cardiovascular Case Reports (CVCR)*, 10.1016/j.ihjccr.2023.01.005, 7, 1, (15-18),
6. Mamta gowda, *J. Fetal Med.* (2016). 3:143–146 DOI 10.1007/s40556-016-0086-y
7. Attia TH, Abd Alhamed MM, Selim MF, Haggag MS, Fathalla D. (2015). Idiopathic Arterial Calcification of Infancy: case report. *J Radiol Case Rep*. 30;9(11): 32-40. doi: 10.3941/jrcr.v9i11.2622.
8. Önal H, Topal N, Yıldız M, Aydın B, Dindar A. (2019). Idiopathic Infantile Arterial Calcification: Echocardiography Diagnosis And Successful Treatment. *Medical Research Reports* 2(1):28-29.
9. Kanchan C, Viv P, Maloriya M. (2017). Familial Idiopathic Infantile Arterial Calcification. *Pediatr Oncall J*. 14: 17-18. doi: 10.7199/ped.oncall.2017.8
10. Nada Djokanovic, Chagit Klieger-Grossmann, Gideon Koren, Does Treatment With Bisphosphonates Endanger the Human Pregnancy?
11. *Journal of Obstetrics and Gynaecology Canada*, Volume 30, Issue 12, 2008, Pages 1146-1148, ISSN 1701-2163



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

Submit Manuscript

DOI:[10.31579/2578-8965/157](https://doi.org/10.31579/2578-8965/157)

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://www.auctoresonline.org/journals/obstetrics-gynecology-and-reproductive-sciences>