

The Potential Role of Peripheral Somatosensory Stimulation in the Treatment of Mental Health Symptoms following Traumatic Brain Injury

LA Nussbaum, TM Janjua, ES Nussbaum *

National Brain Aneurysm & Tumor Center.

*Corresponding Author: ES Nussbaum, National Brain Aneurysm & Tumor Center.

Received date: **November 14, 2023**; Accepted date: **December 01, 2023**; Published date: **December 08, 2023**

Citation: LA Nussbaum, TM Janjua, ES Nussbaum (2023), The Potential Role of Peripheral Somatosensory Stimulation in the Treatment of Mental Health Symptoms following Traumatic Brain Injury, *J. Neuroscience and Neurological Surgery*, 14(2); DOI:[10.31579/2578-8868/282](https://doi.org/10.31579/2578-8868/282)

Copyrights: © 2023, ES Nussbaum. This is an open-access article distributed under the terms of The Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Abstract

Background

Patients with traumatic brain injury (TBI) may develop symptoms of post-traumatic stress disorder (PTSD), depression, and anxiety. Despite decades of research investigating treatment options, these mental health symptoms remain a major cause of quality-of-life impairment. In a series of patients with PTSD being treated with peripheral somatosensory stimulation (PSS), we further evaluated 3 patients who had suffered from TBI to detail their clinical response to therapy.

Methods

Three patients with TBI and PTSD underwent daily PSS therapy over a 4-week period. Patients were evaluated using a Veterans Affairs TBI scale of cognitive impairment and subjective mental health symptoms at baseline and then at the conclusion of the treatments. All data were obtained through a self-reported survey.

Results

Three men with TBI and PTSD completed one month of daily PSS therapy. No adverse events were described by the patients. All patients demonstrated improvement in terms of their scores assessing cognitive impairment and subjective symptoms. The Patients beginning with greater impairment appeared to derive more benefit from therapy.

Conclusions

PSS stimulation appeared to improve symptoms in three patients with TBI and PTSD. Symptoms related to both anxiety and depression were improved. We suggest that further investigation into the potential usefulness of PSS therapy in patients with TBI and mental health symptoms is warranted.

Key words: anxiety ; depression ; neuromodulation ;PTSD ; somatosensory stimulation ; TBI ; trauma

Introduction

A significant percentage of patients with TBI will develop symptoms of PTSD, particularly following severe trauma [1-5]. These individuals may struggle with anxiety, depression, and even suicidal ideation in extreme cases. The mental health symptoms associated with PTSD in the setting of TBI result in a tremendous societal burden in terms of impaired quality of life, functional disability, medical costs, and unemployment [3-14]. It is estimated that more than 7 million Americans are diagnosed with PTSD every year. [2-5] In addition to pharmacotherapy and psychological interventions, treatment for PTSD has included various methods of relaxation including yoga and deep breathing exercises. [7-10,15]

Peripheral somatosensory stimulation (PSS) therapy is a non-invasive technique which may be beneficial to patients with a variety of neurological disorders. [16-19] We recently evaluated the impact of PSS on symptoms in patients with PTSD. [20] Three of these patients had a history of TBI, and this report focuses on our experience with this small subset of TBI patients.

Methods

We treated 6 patients who had been diagnosed with and suffered from active PTSD symptoms with PSS using the NeuroGlove device. NeuroGlove is a non-invasive device that provides PSS stimulation in the form of pneumatic puffs of air directed at the volar surface of the distal forearm, the palm, and the fingers. Subjects used the device at home for 1 hour of therapy per day (30 minutes using each hand) for 4 weeks and were directed to synchronize their breathing to the firing (on/off cycle) of the machine to encourage relaxation during device use. The results of this trial have been previously reported. [20]

Of the six patients in the trial, 3 had a prior history of significant TBI. All felt that the TBI had at least in part contributed to their PTSD. We interviewed these patients to obtain detailed information regarding their response to treatment and analyzed the results obtained in this small subgroup of patients with TBI, focusing on their mental health symptoms of anxiety, depression, and general well-being to evaluate the potential usefulness of PSS in this subgroup of patients. We utilized the Veterans Affairs TBI Evaluation of Cognitive Impairment and Subjective Symptoms Scoring System [21]. This scale surveys patients in terms of their ability to function in 9 areas: memory and attention, judgement, social interaction, orientation, motor activity, visual spatial orientation, subjective symptoms, neurobehavioral symptoms (irritability, anxiety),

and communication ability. Each of the 9 areas was graded based on an ordinal response as: 0 for normal, 1 for mild impairment, 2 for moderate impairment, or 3 for severe impairment.

Results

Three men with a history of TBI and active PTSD completed the trial. Mean age was 45 years; age range was 40-54. Compliance with the treatment protocol was excellent based on patient reporting and our ability to query the devices for a log of treatment times. All three patients enjoyed the treatments and stated that they looked forward to using the device. All patients were interested in continuing to use the device after the conclusion of the 4-week treatment regimen. No adverse events were reported related to device use.

The results of the evaluation based on the Veterans Affairs TBI Evaluation are shown in **Table 1**. All patients demonstrated overall improvement in their scores with the greatest effects noted in patients 1 and 3. Although patient 2 improved, his baseline total score was low suggesting that his symptoms related to TBI were minimal even at baseline. The greatest improvements were seen in patient’s subjective symptoms and orientation. The total scores at baseline compared to the conclusion of the trial are depicted in **Figure 1**.

Category	Patient 1 Baseline - Conclusion	Patient 2 Baseline - Conclusion	Patient 3 Baseline - Conclusion
Memory	3 - 1	2 - 2	3 - 2
Judgement	2 - 1	0 - 0	3 - 2
Social Interaction	2 - 1	0 - 0	3 - 2
Orientation	3 - 1	1 - 0	3 - 1
Motor Activity	2 - 1	0 - 0	3 - 2
Visual Spatial Orientation	3 - 1	0 - 0	3 - 2
Subjective Symptoms	3 - 1	2 - 1	2 - 1
Neurobehavioral Effects	3 - 1	0 - 0	3 - 2
Communication	2 - 1	0 - 0	3 - 2
Total Score	21 - 9	5 - 3	26 - 16

Table 1: Scores Obtained at Baseline and Conclusion of Treatments in 3 Patients with TBI and PTSD




Figure 1: Bar graph comparing total scores at baseline and at trial conclusion

Discussion

The relationship between TBI and symptoms of PTSD, depression, and anxiety has been well-described. [1-5] PTSD is a potentially disabling condition that appears to represent a form of anxiety disorder associated with hypervigilance and increased sympathetic output. [6-12] Although the mainstay of treatment for PTSD was originally pharmacologic, an increasing emphasis has been placed on psychological interventions and various relaxation techniques including yoga and deep breathing exercises. [8-12,15] Similarly, depression and anxiety are prevalent conditions with tremendous societal and personal costs [12-14].

Our previous work demonstrated a statistically significant improvement in symptoms of PTSD, anxiety, and depression with the use of PSS. [20,22] The scientific bases for such improvement may be related to the role played by sensory processing in anxiety, depression, and PTSD. PTSD has been associated with a variety of physiological alterations within the brain, and increasing evidence suggests that impaired sensory processing may play a critical role in the development and pathophysiology of this disorder. [23] Haricharichan et al postulated that alterations in the neural pathways important for processing sensory input have a cascading effect on the ability to perform higher cognitive functions implying that abnormal sensory processing may be contributory and associated with PTSD [24]. In addition, it has been shown that individuals with PTSD have decreased prefrontal cortex activation resulting in impaired sensory integration and emotional regulation [25-7]. Engel-Yeger et al identified reproducible patterns of sensory hypersensitivity in patients suffering from PTSD [28].

Interestingly, a variety of physiological alterations within the brain have been associated with anxiety and depression, and increasing evidence suggests that impaired sensory processing may play a critical role in these disorders [29-34]. Of note, various sensory inputs have been shown to impact symptoms in patients with depression and anxiety; auditory stimulation with music, olfactory stimulation with particular odors, and gustatory stimulation with particular foods have all been shown to improve mood and relieve anxiety [35-38]. Transcranial magnetic stimulation and vagal nerve stimulation have also been used to ameliorate anxiety and depressive disorders [39-40]. Based on these findings, we hypothesized that tactile PSS might be beneficial in patients with the mental health symptoms related to TBI.

It should be noted that PSS has been shown to be of therapeutic benefit in a variety of neurological disorders. PSS has shown promise in humans after stroke and also in rodent models of cerebral ischemia where PSS can actually prevent ischemic injury if applied early enough. [41-43]. The mechanism for this protection remains unclear but may include enhanced neuronal reorganization encouraging neuronal recovery and/or improved regional cerebral perfusion through the recruitment of local collateral blood supply [44]. Interestingly, similar benefits have been demonstrated in patients with Parkinson's disease, and PSS has also shown promise following traumatic brain injury and in inflammatory, auto-immune conditions such as multiple sclerosis [45-49].

Based on this, we analyzed a small subgroup of 3 patients with PTSD and TBI who were treated with one month of PSS on a daily basis. All three patients were evaluated using the Veterans Affairs TBI Evaluation, and all showed improvement in their scores over the course of the 4-week trial. As noted above, patient 2 appeared to have minimal symptoms related to TBI and showed limited improvement. It is notable that the other two patients had significant impairment at baseline, and both showed meaningful improvement in their scoring by the conclusion of the trial.

Conclusions

We describe our experience with 3 patients with TBI who underwent one month of PSS treatment on a daily basis. This is a very small group of patients precluding significant conclusions regarding efficacy of PSS on patients with TBI. Nevertheless, this study was meant to evaluate in very

preliminary fashion the potential usefulness of PSS in patients with TBI, potentially forming the basis for a larger study. All patients showed improvement in their symptoms suggesting that further investigation into the potential use of PSS in the treatment of patients with TBI is warranted.

Institutional Review Board Statement: The study was approved by the Western (WCG) Institutional Review Board (protocol code: 20233103, date of approval: 7/19/2023).

Trial Registration: This study was registered with clinicaltrials.gov prior to initiation any work or patient enrollment. The trial number is NCT06050590.

Disclosure: Drs. Leslie and Eric Nussbaum are shareholders in Neuro Glove, LLC

References

1. Howlett, J. R., Nelson, L. D., & Stein, M. B. (2021). Mental health consequences of traumatic brain injury. *Biological Psychiatry*, 91(5), 413-420.
2. Vasterling, J. J., Jacob, S. N., & Rasmusson, A. (2018). Traumatic brain injury and posttraumatic stress disorder: Conceptual, diagnostic, and therapeutic considerations in the context of co-occurrence. *Journal of Neuropsychiatry and Clinical Neurosciences*, 30(2), 90-100.
3. Loignon, A., Ouellet, M. C., & Belleville, G. (2020). A systematic review and meta-analysis on PTSD following TBI among military/Veteran and civilian populations. *The Journal of Head Trauma Rehabilitation*, 35(1), E21-E35.
4. Alosco, M. L., Supelana, C. J., & Vasterling, J. J. (2017). Traumatic brain injury and psychological trauma. In S.N. Gold (Ed.), *APA handbook of trauma psychology: Foundations in knowledge, Vol. 1* (pp. 333-347). American Psychological Association.
5. Albrecht, J. S., Abariga, S. A., Rao, V., & Wickwire, E. M. (2020). Incidence of new neuropsychiatric disorder diagnoses following traumatic brain injury. *The Journal of Head Trauma Rehabilitation*, 35(4), E352-E360.
6. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders DSM-IV-R*. 4. Washington DC: 2000. Revised.
7. Yehuda R, Hoge CW, McFarlane AC, Vermetten E, Lanius RA, Nievergelt CM, Hobfoll SE, Koenen KC, Neylan TC, Hyman SE. Post-traumatic stress disorder. *Nat Rev Dis Primers*. 2015 Oct 08;1:15057.
8. Sherin JE, Nemeroff CB. Post-traumatic stress disorder: the neurobiological impact of psychological trauma. *Dialogues Clin Neurosci*. 2011;13(3):263-278.
9. Bryant RA, Friedman MJ, Spiegel D, Ursano R, Strain J. A review of acute stress disorder in DSM-5. *Depress Anxiety*. 2011 Sep;28(9):802-817.
10. Qi W, Gevonden M, Shalev A. Prevention of Post-Traumatic Stress Disorder After Trauma: Current Evidence and Future Directions. *Curr Psychiatry Rep*. 2016 Feb;18(2):20.
11. Blechert J, Michael T, Grossman P, Lajtman M, Wilhelm FH. Autonomic and respiratory characteristics of posttraumatic stress disorder and panic disorder. *Psychosom Med*. 2007 Dec;69(9):935-943.
12. National Institute of Mental Health. [March 6, 2011]: *The Numbers Count: Mental Disorders in America*.
13. Johnston, K. M., Powell, L. C., Anderson, I. M., Szabo, S., & Cline, S. (2019). The burden of treatment-resistant depression: A systematic review of the economic and quality of life literature. *Journal of Affective Disorders*, 242, 195-210.

14. Rehm, J., & Shield, K. D. (2019). Global burden of disease and the impact of mental and addictive disorders. *Current Psychiatry Reports*, 21, 10.
15. Descilo T, Vedamurtachar A, Gerbarg PL, et al. Effects of a yoga breath intervention alone and in combination with an exposure therapy for post-traumatic stress disorder and depression in survivors of the 2004 South-East Asia tsunami. *Acta Psychiatr Scand*. 2010 Apr;121(4):289–300
16. Celnik P, Hummel F, Harris-Love M, Wolk R, Cohen LG. Somatosensory stimulation enhances the effects of training functional hand tasks in patients with chronic stroke. *Archives of Physical Medicine and Rehabilitation*, 2007;88(11), 1369–1376.
17. Nussbaum HA, Nussbaum ES: Peripheral Sensory Stimulation for Neurological Disorders. A Novel, Non-invasive Therapeutic Option. Review Article. *J Neurosci and Neurol Surgery* 13:267-271, 2023.
18. Kristina JP, Kromer JA, Cook AJ, Hornbeck T, Lim EA, Mortimer BJ, Fogarty A, Han SS, Dhall R, Halpern C, Tass PA.: Coordinated reset vibrotactile stimulation induces sustained cumulative benefits in parkinson's disease. *Front Physiol* 2021; 12: 624317
19. Lucente G, Valls-Sole J, Murillo N, Rothwell J, Coll J, Davalos A, Kumru H: Noninvasive Brain Stimulation and Noninvasive Peripheral Stimulation for Neglect Syndrome Following Acquired Brain Injury. *Neuromodulation: Technology at the Neural Interface* 23(3)2020, 312-323.
20. Janjua TM, Pederson J, Nussbaum ES: Peripheral somatosensory stimulation in the treatment of post-traumatic stress disorder. A clinical trial. *J Neurol Dis* 11:569-575, 2023.
21. National Academy of Sciences, Engineering, and Medicine: Committee on the Review of the Department of Veterans Affairs Examination for TBI. National Academies Press;2019.
22. Nussbaum LA, Janjua T, Pederson J, Nussbaum ES: Peripheral somatosensory stimulation in the treatment of depression and anxiety. A clinical trial. In press
23. Ayres AJ: Treatment of sensory integrative dysfunction *A J Occ Ther* 1440, 1972.
24. Harricharan S, McKinnon MC, Lanius RA: How processing sensory information from the internal and external worlds shape the perception and engagement with the world in the aftermath of trauma: Implications for PTSD. *Front Neurosci* 15:2021.
25. Etkin A, Wager TD: Functional neuroimaging of anxiety. A metanalysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *Am J Psychiatr* 164: 1476-1488, 2007.
26. Nicholson AA, Friston KJ, Zeidman P, Harricharan S, McKinnon MC, Densmore M, et al: Dynamic causal modeling in PTSD and its dissociative subtype: bottom-up vs top-down processing within fear and emotional regulation circuitry. *Human Brain Mapp* 38: 5551-5561, 2017.
27. Shin LM, Rauch SL, Pittman RK: Amygdala, medial prefrontal cortex, and hippocampal function in PTSD. *Ann NY Acad Sci* 1071:67-79, 2006.
28. Engel-Yeger B, Palgy-Levin D, Lev-Wiesel R: The sensory profile of people with PTSD. *Occup Ther Ment Health* 29:266-78, 2013.
29. Engel-Yeger B, & Dunn W. (2011). The relationship between sensory processing difficulties and anxiety level of healthy adults. *British Journal of Occupational Therapy*, 74, 210-216.
30. Liljas AEM, Jones A, Cadar D, Steptoe A, Lassale C. (2020). Association of multisensory impairment with quality of life and depression in English older adults. *JAMA Otolaryngology. Head & Neck Surgery*, 146, 278-285.
31. Fitzgerald P J. (2013). Gray colored glasses: Is major depression partially a sensory perceptual disorder? *Journal of Affective Disorders*, 151, 418-422.
32. Acevedo BP, Aron EN, Aron A, Sangster MD, Collins N, Brown L (2014). The highly sensitive brain: An fMRI study of sensory processing sensitivity and response to other's emotions. *Brain and Behavior: A Cognitive Neuroscience Perspective*, 4, 580-594.
33. Canbeyli R. (2010). Sensorimotor modulation of mood and depression: An integrative review. *Behavioural Brain Research*, 207, 249-264.
34. Canbeyli R: Sensory stimulation via the visual, auditory, olfactory and gustatory systems can modulate mood and depression. *Eur J Neurosci* 55(1):244-263, 2022.
35. Aalbers S, Fusar-Poli L, Freeman RE, Spreen M, Ket JC, Vink AC, Maratos A, Crawford M, Chen XJ, Gold C. (2017). Music therapy for depression. *Cochrane Database of Systematic Reviews*, 11 CD004517.
36. Al-Karawi D, Jubair L. (2016). Bright light therapy for nonseasonal depression: Meta-analysis of clinical trials. *Journal of Affective Disorders*, 198, 64-71.
37. Matsubara E, Ohira T. (2018). Inhalation of Japanese cedar (*Cryptomeria japonica*) wood odor causes psychological relaxation after monotonous work among female participants. *Biomedical Research*, 39, 241-249.
38. Meier BP, Noll SW, Molokwu OJ. (2017). The sweet life: The effect of mindful chocolate consumption on mood. *Appetite*, 108, 21-27.
39. Andrade P, Noblesse LHM, Temel Y, AckermansL., LimLW, Steinbusch HWM, Visser-Vandewalle V. (2010). Neurostimulatory and ablative treatment in major depressive disorder: A systematic review. *Acta Neurochirurgica*, 152, 565-577.
40. Askalsky P, Iosifescu DV (2019). Transcranial photobiomodulation for the management of depression: Current perspectives. *Neuropsychiatric Disease and Treatment*, 15, 3255-3272.
41. Lay CC, Jacobs N, Hancock AM, Zhou Y, Frostig RD. Early stimulation treatment provides complete sensory-induced protection from ischemic stroke under isoflurane anesthesia. *The European Journal of Neuroscience*. 2013;38: 2445-52.
42. Hancock AM, Lay CC, Davis MF, Frostig RD. Sensory Stimulation-Based Complete Protection from Ischemic Stroke Remains Stable at 4 Months Post-Occlusion of MCA. *Journal of Neurological Disorders* 2013; 1: 135.
43. Conforto AB, dos Anjos SM, Bernardo WM, Silva AA, d Conti J, Machado AG, Cohen LG. Repetitive peripheral sensory stimulation and upper limb performance in stroke: A systematic review and meta-analysis. *Neurorehabilitation and Neural Repair* 2018; 32(10), 863–871.
44. Nussbaum ES, Janjua TM, Lowary J, Defillo A, Myers M, Nussbaum LA: Peripheral sensory stimulation of the hand in the treatment of stroke. A preliminary study of safety and effectiveness. *J Neurol Disord* 11(1):2023
45. Tass PA: Vibrotactile Coordinated Reset Stimulation for the Treatment of Parkinson's Disease. *Neural Regeneration Research*. 17(7):1495-1497 (2021)
46. Tramontano M, Morone G, De Angelis S, Conti L, Galeoto G, Grasso MG. Sensor-based technology for upper limb rehabilitation in patients with multiple sclerosis: A randomized controlled trial. *Restor Neurol Neurosci* 2020;38:333-41.
47. Hall KD, Lifschitz J. Diffuse traumatic brain injury initially attenuates and later expands activation of the rat somatosensory

- whisker circuit concomitant with neuroplastic responses. *Brain Res* 2010; 1323:161-173.
48. Lew HL, Weihing J, Myers PJ, Pogoda TK, Goodrich GL. Dual sensory impairment (DSI) in traumatic brain injury (TBI)--An emerging interdisciplinary challenge. *Neurorehabilitation* 2010; 26:213-222.
49. Zheng Z, Dong X, Li Y, Gao W, Jiang R, Yue S, Zhou Z, Zhang J: Electrical stimulation improved cognitive deficits associated with traumatic brain injury in rats. *Brain and Behavior* 2017; 7:e00667.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

Submit Manuscript

DOI: [10.31579/2578-8868/282](https://doi.org/10.31579/2578-8868/282)

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/neuroscience-and-neurological-surgery>