

# K of the Cured food Nutritional Value, Digestibility and High Quality with Special Reference to Their Role on Public Health

**Fahim Aziz Eldin Shaltout**

Food Control, Faculty of Veterinary Medicine, Benha University, Egypt.

**Corresponding author:** Fahim Aziz Eldin Shaltout, Food Control, Faculty of Veterinary Medicine, Benha University, Egypt.

**Received date:** January 22, 2025; **Accepted date:** February 24, 2025; **Published date:** March 11, 2025

**Citation:** Fahim Aziz Eldin Shaltout, (2025), K of the Cured food Nutritional Value, Digestibility and High Quality with Special Reference to Their Role on Public Health, *J. Nutrition and Food Processing*, 8(3); DOI:10.31579/2637-8914/293

**Copyright:** © 2025, Fahim Aziz Eldin Shaltout. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract:

The human beings are a unique host of heterogeneous and vivacious group of microbes, which together forms the human-microbiome superorganism types. The Human being gut serves as a home to over 100–1000 microbial species types, which primarily modulate the host internal environment and thereby, play a major role in public health types. The spectacular symbiotic relationship has attracted extensive research in the field types. More specifically, the microorganisms play key roles in defense activity, eupepsia along with catabolism and anabolism, and impact brain-gut responses types. The emergence of microbial with resistance and the tolerance to existing conventional drugs and antibiotics has decreased the drug efficacies on pathogenic microorganisms. The new biotechnology mediated nano-encapsulated multiplex supplements appear to be high cost and inconvenient. low-cost, receptive and intrinsic approach to achieve health benefits is vital in the present era types. The Supplementation with the food probiotics, the food prebiotics, and the food synbiotics has shown promising results against various enteric pathogenic microorganisms due to their unique ability to compete with pathogenic microbiale for adhesion sites, to alienate pathogenic microorganisms or to stimulate, modulate and regulate the host's immune response by initiating the activation of specific genes in and outside the host intestinal tract. The Probiotics have been shown to regulate fat storage and stimulate intestinal angiogenesis. The investigations aim to underline the possible beneficial impact of the food probiotics for the public health and the medical importance and for the good lifestyle types.

**Key words:** food probiotics; food prebiotics; food synbiotics; new biotechnology mediated nano-encapsulated multiplex supplements

## Introduction

The food Probiotic is a phrase of the new era, denotation “for life” and is in use to name bacterial association with beneficial effects on the human beings and animal health. The food probiotics as modification of the floral/microbial diversity in the human beings’ bodies and replaces the harmful microbes with useful ones. The microbial concentration of a particular type of the bacteria in the stool samples of the infected diarrhea in young children were significantly lower in comparison to healthy young children. The oral administration of the live microorganisms to the patients with diarrhea and help restore a healthy gut flora type. The new definition of food probiotic was put forward as a viable mono or mixed culture of bacteria types which, when applied to animal or man, affects the host beneficially by improving the properties of the indigenous flora types. The specific food probiotic cultures along with substantial health benefits upon consumption types. The ecological consideration of the gut flora is necessary to understand their relevance in the human being’s health, as well as the probiotic food products types concept importance. Each individual has a unique signature of more than one hundred to one thousand microbial species in the gastrointestinal tract in human. The Bacterial cells comprise half of the wet weight of colonic material and

their numbers exceed by tenfold the number of tissue cells forming the human beings body. The stomach contains 103 different bacterial species, and the total microbial population of the colon comprises of about 1011–1012 cfu/g. The Bacterial colonization of the gut begins at birth when newborns are first exposed to a non-sterile environment. It evolves and transforms over a lifetime, depending on a complex and dynamic interplay between the diet, genome, and lifestyle of the host, as well as antibiotic use. Notable age-specific compositional shifts reported in gut microbiota composition include a decrease in the Bacteroidetes/Firmicutes ratio and a marked decrease in bifidobacteria in people aged > 60 years, around the time that the immune system starts to decline. The composition of the core intestinal microflora is considered to be essentially stable throughout adulthood time. The beneficial usage of intestinal microflora, referred to as “colonization resistance” or the “barrier effect” is an important mechanism used by the indigenous gut bacteria to maintain their presence and confer niche protection against freshly ingested microorganisms, including pathogenic microorganisms. It could be assumed that manipulation of the gut microflora to improve the relative numbers of “beneficial bacteria”, which have certain impacts

on immune activity, digestion, metabolism, and brain-gut communication mechanism. The alterations in their diversity may result in several disorders and diseases, for which conventional medicines provide very limited efficacy mainly due to the emergence of antibiotic-resistant and tolerant pathogenic microbes. Attempts to avoid such critical issue by increasing the drug delivery system to the target sites using nano-encapsulated multiplex supplements have been reported to as a possible solution, although, such approaches appear to be cost-effective, and inconvenient for common use. A simple, low-cost, receptive and intrinsic means to improve host health has become a critical issue in the present era. In this context, the food probiotics tends to serve as supplement to the host microflora and provide protection against various enteric pathogenic microorganisms. The food Probiotics are known to demonstrate promising results like improved gut barrier activity; adding to the unique ability to compete with pathogenic microbiota for adhesion to the gut and improve their colonization method. The food Probiotics stimulate, modulate and regulate the host's immune response by initiating the activation of specific genes of localized host cells. The gastrointestinal hormone release and regulate brain behavior through bidirectional neuronal signaling, as part of the gut-brain axis mechanism so much. The food Probiotics plays a significant role in inducing intestinal angiogenesis by vascular endothelial growth factor receptor signaling that, in turn, regulates acute and chronic inflammation in intestinal mucosal tissue caused by the progression of the inflammatory bowel disease types. The food Probiotics have physiological activities that contribute to the health of the host environment regulating microbes and are helpful in combating overweight and obesity so much. The food probiotics have considerable potential in nutritional and clinical applications, the implementation of the food probiotics into the human beings health, nutrition and regulation of different abnormalities types. The investigations emphasize the possible benefaction of the food probiotics for improving the human being's health, nutrition optimization and regulation of common metabolic disorders or abnormalities action types [1,2,3,4,5,6 and 7].

The meanings for the food probiotics, the food prebiotics, and the food synbiotics, they can be best explained as microbe or a group of microbes that inhabits within the gut and nourishes the host body internally. They are commonly consumed as preparations with active live cultures and contain bacteria, such as lactobacilli, lactococci or bifidobacteria that has been isolated from natural environments. The food probiotics have been recognized as key health promoters to humans, investigating the culture conditions and viability of the food probiotic strains during processing and storage; sensitivity to low pH values, gastric fluid, bile, pancreatic and intestinal fluids and intestinal or respiratory mucus; adherence to isolated cells or cell cultures and interactions with other pathogenic microorganisms' types [8,9,10,11,12,13 and 14].

### The Postbiotics

The bacterial products, in the absence of life microorganisms, may have similar effects on signaling pathways and barrier activity. The bacterial products are broadly characterized as postbiotics and can be defined as dead bacterial products or metabolic byproducts from food probiotic microorganisms that have biologic activity in the host. The postbiotics include bacterial metabolic byproducts, such as bacteriocins, organic acids, ethanol, diacetyl, acetaldehydes and hydrogen peroxide, but it is found that certain heat-dead food probiotics can retain important bacterial structures that may exert biological activity in the host body. The metabolic products have a broad inhibitory property toward pathogenic microbes and can be used as an alternative to antibiotics. The Postbiotics are non-toxic, non-pathogenic and resistance to hydrolysis by mammalian enzymes, as these are non-viable bacterial products or metabolic byproducts from the food probiotics. In some instances, postbiotics can

increase barrier activity against species like *Saccharomyces boulardii*, and improve angiogenesis in vitro and in vivo in epithelial cells by activation of  $\alpha 2\beta 1$  integrin collagen receptors types. The species of *Bifidobacterium breve*, *Bifidobacterium lactis*, *Bifidobacterium infantis*, *Bacteroides fragilis*, *Lactobacillus*, *Escherichia coli* and *Faecalibacterium prausnitzii* types [15,16,17,18,19,20 and 21].

### The food Prebiotics

The exploration of the food probiotics have led to the development of the food prebiotics, which are certain nutrients that modify the gut microbial flora although not easily digested by human beings but have a selective role in stimulation of growth or activity of beneficial bacterial species in the gut. The food prebiotics includes bifidogenic properties of insulin, oligofructose, and fructo-oligosaccharides compounds synthetically produced from sucrose, as well as the galactose-containing and xylose-containing oligosaccharides compounds types [22,23,24,25,26,27 and 28]. The fermentation of carbohydrates represents a major source of energy for epithelial cells in the colon and food prebiotics can readily fulfill these requirements as a result of their fermentation by gut microbiota, as bifidobacteria bacteria. The bifidobacteria, several other gut microorganisms that play a significant role in fermenting these non-digestible oligosaccharides compounds types. The food Prebiotics can be obtained naturally from sources like vegetables, fruits, and grains consumed in our daily life span. The food Prebiotics not only serve as an energy source but have several health benefits such as reducing the prevalence and duration of diarrhea, providing relief from inflammation and other symptoms associated with intestinal bowel disorders, and exerting protective effects to prevent colon cancer types [29,30,31,32,33,34 and 35]. The food Prebiotics are implicated in enhancing the bioavailability and uptake of minerals, lowering of some risk factors for cardiovascular disease, and promoting satiety and weight loss conditions. The vast nutritional and medicinal benefits, research regarding screening new versatile food prebiotics is scarce mechanism [36,37,38,39,40,41 and 42].

### The food Synbiotics

The Development in microbial investigation has led to formation of food synbiotics which is a fusion of the food probiotics and the food prebiotics products and helps in enhancing the survival and the implantation of live microbial dietary supplements in the gut of human. The synergistic benefits are more efficiently promoted when both the food probiotic and food prebiotic work together in the living system body. The food symbiotic relationship between the food prebiotics and the food probiotics contributes significantly to health of human beings. The Commercial interest in the active foods products types containing the food synbiotics has improved due to the awareness of the benefits for gut health of human beings, disease prevention and therapy of diseases. The development of new health-promoting foods products types. The selection of new cultures demonstrating an increased ability to colonize the human being's gut, along with their ability to digest new forms of the food prebiotics. The Conventional trials and investigation have shown that the various beneficiary effects of the food probiotics, the food prebiotics, and the food synbiotics are much more effective than their unitary use known till date now [43,44,45,46,47,48 and 49].

### The Clinical significance of the food probiotics and its potential applications

The use of the food probiotics for the clinical health benefits is a fascinating area of research that the present era has yet to explore. Some of the elite properties of the probiotics types, such as the anti-pathogenicity, anti-diabetic, anti-obesity, anti-inflammatory, anti-cancer,

anti-allergic, and angiogenic activities and their effect on the brain and the central nervous system of the body. The Anti-pathogenic activity of the probiotic's types, The Anti-pathogenic activity is regarded as one of the most beneficial effects of the probiotics because unlike classic antibiotics, disturbance or alteration in the composition of the complex population of the gut microbiota is inhibited. The anti-pathogenic activity of the probiotics or a probiotic mixture. The influence of the probiotics on the survival of *Salmonella enterica*, Serovar typhimurium and *Clostridium difficile* in an in vitro model and postulated that the probiotics inhibit pathogenic microorganisms by the production of the short-chain fatty acids, as the acetic, the propionic, the butyric and the lactic acids. The short-chain fatty acids help to maintain an appropriate pH in the colonic lumen, which is imperative in the expression of numerous bacterial enzymes and in metabolism of foreign compounds and carcinogens in the gut suggested that a wide variety of anti-pathogenic compounds, like the bacteriocins, ethanol, organic acids, diacetyl, acetaldehydes, hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and peptides are produced by many probiotics' types. The peptides and the bacteriocins, in particular are mostly involved in increasing the membrane permeability of the target cells, which leads to the depolarization of the membrane potential and, ultimately, cell death of the body. The production of the H<sub>2</sub>O<sub>2</sub> by the bacterial groups causes the oxidation of sulfhydryl groups, resulting in the denaturation of several enzymes results in the peroxidation of membrane lipids, thus, increasing membrane permeability of the pathogenic microorganism and consequently, cell death of the body [50,51,52,53,54,55 and 56]. The compounds may act by lowering pH by organic acids like lactic and acetic acids also. The production of the anti-pathogenic bioactive compounds that directly affect pathogenic microorganisms, the probiotics stimulate host anti-pathogenic defense pathways, such as stimulating or activating the pathway involved in the production of defensins that are cationic anti-microbial peptides produced in several cell types including Paneth cells in the crypts of the small intestine and intestinal epithelial cells of the body. The mechanism by which the probiotics exert anti-pathogenic activity is by competing for pathogen binding and receptor sites, as well as for available nutrients and growth inside human beings [57,58,59 ,60,61,63 and 64].

### The Urogenital health care

Over one billion women around the world suffer from non-sexually transmitted urogenital infections, such as bacterial vaginosis, urinary tract infection and several other yeast infections to human beings. The species typically associated with BV include *Gardnerella vaginalis*, *Ureaplasma urealyticum*, and *Mycoplasma hominis* Sexually transmitted diseases are significant cause of morbidity in the world. The two most commonly documented bacterial STDs in some developed countries in the world are gonorrhoea and Chlamydia, which are caused by *Neisseria gonorrhoeae* and *Chlamydia trachomatis*, respectively The pathogenic microbes, among others, are concurrently becoming resistance to the present medicines in the world. The development of new medicines, developing new live supplements, like non-pathogenic microbes that act against the pathogenic microorganisms [65,66,67,68,69,70 and 71]. There is an association between abnormal vaginal microbial flora and an improved incidence of urinary tract infection. About 50 different species inhabiting the vagina, like *Lactobacillus* species, *Lactobacillus brevis*, *Lactobacillus casei*, *Lactobacillus vaginalis*, *Lactobacillus delbrueckii*, *Lactobacillus salivarius*, *Lactobacillus reuteri*, and *Lactobacillus rhamnosus* that are regarded as the main regulators of the vaginal micro-environment. Imbalance in the microbial composition greatly influences the health of the vaginal microenvironment, potentially leading to compromised state of bacterial vaginosis in the world. The compromised states in the world

can be reassured by balancing the number of the *Lactobacillus* sp. through the supplementation of the probiotics types [72,73,74,75,76,77 and 78].

### The Anti-diabetic activities of the probiotics

The International Diabetes Federation (IDF) of Southeast Asia in the world, 425 million people have diabetes disease worldwide including 78 million people in the Southeast Asian region in the world. This number is expected to rise to 629 million by 2045 if nothing is done. The management of this disorder includes multiple medications although there is no definitive cure for diabetes disease. Nonetheless, bimolecular and pharmacological researchers have made progress in understanding the importance of food synbiotics in curing the disorder. The large-scale 16 S rRNA gene sequencing, quantitative real-time PCR and fluorescent in situ hybridization, the connection between the composition of the intestinal microbiota and metabolic diseases, like obesity and diabetes disease, enhancing the beneficial microbial by the use of the probiotics is expected to play a significant role in neutralization of the disorder Gram-negative bacteroidetes and the Gram-positive firmicutes are two specific bacterial phyla that dominant the gut microenvironment [79,80,81,82,83,84,85 and 86]. Recent research has proven that obesity is associated with improved bacteroidetes over time, concurrent with a reduction in firmicutes More specifically, patients with type-2 diabetes disease have significantly reduced numbers of firmicutes species, such that the bacteroidetes/firmicutes ratio has improved, which positively correlates with plasma glucose concentration. The development of auto-immune diseases, such as type-1 diabetes disease. Alterations in the microbiome improve invasion of opportunistic pathogenic microorganisms, which are resistant to oxidative stress and simultaneously capable of reducing sulfates and inhibiting the growth of butyrate-producing bacteria [87,88,89,90,91,92 and 93].

Management of type-2 diabetes disease by modulating gut hormones, such as gastric inhibitory polypeptide and glucagon-like peptide-1, through probiotic and food prebiotic interventions is another convincing strategy. In this context, hormones play an implicated role in glucose homeostasis, which results in neutralizing the disorder caused by peripheral insulin resistance or failure of  $\beta$ -cells to produce insulin Currently, research is focused on production new food prebiotics, as arabinoxylan and arabinoxylan oligosaccharides compounds, which show promising results in counteracting related metabolic disorders, because both carbohydrates have been linked to adiposity reduction [94,95,96,97,98,99 and 100].

### The Anti-obesity activity of the probiotics

The Abnormal or excessive fat (obesity) accumulation that directly impairs health is linked to an improve in energy availability, sedentariness and a greater control of ambient temperature, leading to an imbalance in energy intake and expenditure It has been evident that transplantation of the intestinal microflora from obese mice into germ-free mice could replicate the obese phenotype and may lead to more efficient at extracting energy from food products types and stimulating lipogenesis.

The Probiotics have physiological activities that contribute to the health of host environment regulating microbes. In most instances, weight loss is facilitated by thermogenic and lipolytic responses through stimulating the sympathetic nervous system. Probiotic strains, *Lactobacillus gasseri* BNR17 have shown properties of inhibiting the improve in adipocyte tissue that are the main source of leptin and adiponectin and thereby, limiting leptin secretion. Other probiotic microbes such as *L. casei*, *Lactobacillus acidophilus* and *Bifidobacterium longum* have been mentioned to have hypocholesterolemic effects [101,102,103,104,105,106 and 107].



### The Anti-inflammatory activity of the probiotics

Crohn's disease (CD) and ulcerative colitis (UC) are among the most chronic inflammatory diseases of the gastrointestinal tract and are collectively called the inflammatory bowel disease. CD can affect any part of the gastrointestinal tract like the mucosa, submucosa, and serosa, and the inflammation can even spread to the whole gastrointestinal tract. In contrast, UC characteristically involves the large bowel; specifically the mucosa and submucosa of the colon. Research has shown that an imbalance in the gut microbiota plays an important pathophysiological role in the positive regulation of the inflammatory bowel disease. The disorder could possibly be altered by supplementation with the probiotics, the prebiotics, and the food synbiotics. The inflammatory bowel disease is being associated with impaired production of the short-chain fatty acids, particularly, acetate, butyrate, and propionate. The short-chain fatty acids have been known to play a key role in maintaining colonic homeostasis in the body [108,109,110,111,112,113 and 114]. The probiotics have anti-inflammatory effects and improve the propulsive colonic activity. It is reasonable to consider that supplementation with indigestible carbohydrates and fiber (food prebiotic) alone, or in combination with the food probiotics to improve the production of the short-chain fatty acids could be useful therapeutic approaches. Developing genetically engineered food probiotic bacterial strains that are able to produce and discharge immunomodulators, such as interleukin-10, trefoil factors (compact proteins co-expressed with mucins in the gastrointestinal tract, or lipoteichoic acid (a major constituent of the cell wall of Gram-positive bacteria that can impact the host immune system, resulting in the restoration of the level of protective commensal bacterial species. *Lactobacillus*, *Bifidobacterium*, *Enterobacter* and *E. coli* are the most widely used the probiotics in foods products types. Apart from these microorganisms, new or genetically modified should be developed to counteract the inflammatory bowel disease [115,116,117,118, 119, 120 and 121].

### The Anti-cancer activity of the probiotics

The natural sources that have anti-carcinogenic effects, as the probiotics have been receiving prime focus in recent years. Thus have attracted intense interest from clinical nutritionists, scientists, and industrialists to work in a collaborative manner to bring down the disease and develop an effective drug with minimal or no side-effects types. The probiotic strains, *The Lactobacillus fermentum*, are highly potent in suppressing colorectal cancer cells and promoting normal epithelial colon cell growth through the production of the ferulic acid type [122,123,124,125,126,127 and 128]. This ability was compared with other probiotics namely the *L. acidophilus* and the *L. rhamnosus* both of which were previously characterized with tumorigenic activity. Again, two different probiotic strains the *L. acidophilus* and the *L. casei* I have been found to show pronounced cytotoxic activities, with in vitro anti-proliferative activity against two colorectal cancer cell lines types. The probiotics could play a significant role in neutralizing cancer, the investigations are limited only to in vitro tests types. The anti-cancer potential of the probiotics must be proven in vivo models and proceed towards animal and clinical trials types [129, 130, 131, 132,133,134 and 135].

### The Anti-allergic activity of the probiotics

The increasing prevalence of the allergic diseases caused by immune disorders is a serious economic and social burden worldwide. Comprehending the fundamental molecular mechanism that contributes to the etiology of allergic diseases, as well as new treatment approaches is vital for the follow-up and prevention of these diseases. In recent times, the beneficial role of the probiotics in the protection and the management of the allergic diseases had advanced the understanding of their cause and

prevention techniques. The *Lactobacillus plantarum*, the potential to prevent the allergy-associated disorders with the production of interleukin-12 and interferon- $\gamma$  in the host types. The *L. plantarum* significantly alleviated allergic symptoms and reduced the levels of total immunoglobulin E, ovalbumin-specific immunoglobulin E, and histamine in the sera of ovalbumin-sensitized mice. In spleen cells of the mice, the *L. plantarum* is known to significantly increase the secretions of interferon- $\gamma$  and interleukin-4, which are responsible for alleviating allergic symptoms. To evaluate the anti-allergic activity of the probiotics and the mode of the action types [136,137, 138, 139,140,141 and 142].

### The Angiogenic activity of the probiotics

The Angiogenesis mechanism has been an important phenomenon and is necessary for wound recovery process mechanism through delineated cellular responses to regenerate damaged tissues in humans. The angiogenic program consists of a deliberately orchestrated series of cellular events by which new vessels arise from pre-existing ones by promoting recruitment of inflammatory cells and producing cytokines, matrix-degrading enzymes, and the chemokines. Deregulated angiogenesis has a prominent impact on the major human beings diseases, such as cancer, diabetic retinopathy, and the inflammatory bowel disease mechanism including CD and UC. The Non-pathogenic probiotic yeast types, *S. boulardii*, has been mentioned to protect against intestinal injury and inflammation. The molecular mechanisms by which the probiotics mediate these beneficial effects remain unclear. The potential mechanisms of the probiotics in angiogenesis process mechanism may include alteration of inflammatory cytokine profiles, down-regulation of pro-inflammatory cascades or induction of regulatory mechanisms in a strain-specific manner, epithelial barrier activity increased, visceral hypersensitivity reduction, spinal afferent traffic, and stress response [143,144,145,146,147,148 and 149].

### The Effect of the probiotics on the brain and the central nervous system

The colonization of the microbiota in the gastrointestinal tract is well-associated with both the gastrointestinal tract and gastrointestinal diseases of human. The investigations have been devoted towards elucidating the influence of gut microbiota on the brain and the central nervous system of human. The "microbiota-gut-brain axis" is an interactive, bi-directional communication established by the exchange of regulatory signals between the gastrointestinal tract and the brain and the central nervous system of human. The effect of the probiotics on the brain and the central nervous system has been mainly studied in clinical trials, where it has been evident that gut microbiota influences the human beings brain development activity of human. In young children with autism spectrum disorder, a daily dose of the *L. plantarum* WCFS1 led to an improvement in their school records and attitude towards food products types, reduced psychological distress in a randomized trial involving healthy peoples treated with oral administration of the *Lactobacillus helveticus* and the *B. longum* types. The decrease in anxiety symptoms by administration of the *L. casei* strain Shirota to patients suffering from chronic fatigue syndrome of human. Despite an improvement in the *Lactobacillus* and *Bifidobacteria* levels, the bowel activities were under investigations [150,151,152,153,154,155 and 156]. The reduced anxiety was due to improved bowel activity of human. Autism spectrum and attention-deficit/hyperactivity disorders in young children could be inhibited by the *L. rhamnosus* administration to the mother at four four weeks from expected delivery. It has been observed that many gut bacteria synthesize to neuroactive compounds similar to those produced in the host brain. The human beings intestinally derived strains of the *L. brevis* and *Bifidobacterium dentium* were mentioned to produce large amounts of  $\gamma$ -

aminobutyric acid, a brain neurotransmitter that helps the human beings to suppress anxiety and depression types. Uses of a multispecies probiotic containing the *L. brevis*, *B. lactis*, the *L. acidophilus*, the *Bifidobacterium bifidum*, the *L. salivarius*, the *L. casei*, and *Lactococcus lactis* to healthy human beings revealed a significant overall reduction in the cognitive reactivity to sad mood types. The Probiotic trials involving patients suffering from anxiety and clinical depression are lacking and require more time and work to validate this effect. The Oral intake of the *L. acidophilus* has been shown to assist people to regulate their mood towards rewards and addictive behavior types [157,158,159, 160,161,162 and 163].

### The Commercial significance of the probiotics

The development to avoid the barriers related to their successful administration and minimal side effects several forms of the probiotics are available commercially and are in use in large amount types [160,161,162,163,164 and 165].

### The Recent advancements and utility the food prebiotics

As food probiotics, food prebiotics is being widely explored for their utility in the various fields of applied science types, more specifically as nutrients and supplements. The food prebiotics are a collection of nutritionally enriched compounds grouped together with the efficiency to increase and support the growth and sustenance of specific beneficial gut microflora types. The food prebiotics is those compounds that are non-digestible and able to specifically modulate the sustenance of health-promoting gut bacteria.

Presently the idea on the complexity as well as the usability of these non-digestible compounds has improved to a greater extent because of the development of various 'omic' tools such as the proteomics, the genomics, the metabolomics, the transcriptomics types. The investigations based on the various mode of synthesis has become the current focus of the present era types [165,166,167,168,169,170 and 171]. The food products types industries of the present decade require simple, sustainable, cost-effective and highly efficient methods for large-scale production and application. Food prebiotic oligosaccharides compounds could be obtained from food products types; otherwise, could be synthesized chemically or enzymatically from disaccharides or other substrates as well as by hydrolysis of the polysaccharides compounds. The food prebiotics of natural origin have already been evaluated for their beneficiary role types. The Enzymes (the  $\beta$ -galactosidase, the fructosyltransferase etc.) from various sources such as microbes and plants are being utilized for their synthesis. The Enzymes are engineered to better regulate regioselectivity and the increase in the yield of reaction which increase the glycodiversification and the quality of the products attained types [172,173,174,175,176,177 and 178]. The emergence of genetically engineered microorganisms resulted in boosting the production of oligosaccharides compounds by fermentation process for large-scale industrial production Due to the tangible association of food prebiotics oligosaccharides compounds with the gut microbiome as well as maintenance and restoration of the microbial homeostasis which is again keenly associated with positive health outcome of the host, the investigations regarding food prebiotics are given much emphasis in the investigations types. The food Prebiotic compounds are food-grade substances from which beneficial short-chain fatty acid could be produced as a result of degradation by microbes such as bifidobacteria and lactobacilli within the host appeal for their utilization as nutrient supplements types. The biomedical benefaction not only covers gastrointestinal system but systems located away. Recent investigations several rat models have demonstrated calcium absorption types, retention bone density and strength is increased due to the intake of the galacto-

oligosaccharides compounds specifically types. Gut microbes influence the expression of gamma-aminobutyric acid receptor in the brain, food prebiotics such as fructo-oligosaccharides compounds and the galacto-oligosaccharides compound is likely to exploit this connection to tune the brain-derived neurotrophic factors, D-serine, and other synaptic proteins such as the synaptophysin and the N-methyl-D-aspartate receptor subunit . The food prebiotics such as the oligofructose, the  $\beta$ -fructan, the oligofructose/inulin mix have immunomodulatory benefits in the case of pathogenic attack, atopic dermatitis, allergic prevention, chronic inflammation and up-regulated responses against the vaccinations types [179,180,181,182,183,184 and 185]. Benefactions of this non-digestible compound have emerged for a variety of skin related-conditions. Improve water retention and prevention of erythema was observed in a hairless mice skin on the galacto-oligosaccharides compounds supplementation types. The investigations prove that skin's barrier properties are improved by the improved dermal expression of cell adhesion and matrix formation markers CD44, and type 1 collagen on the galacto-oligosaccharides compounds treatment Again, the Galacto-oligosaccharides compound alone or with *B. breve* is found to impede water and keratin depletion effectuated by phenolic compounds types . Food prebiotics are widely being explored for their usefulness in the treatment of various types of disorder and diseases types [186,187, 188, 189, 190,191 and 192].

### Conclusion

The Probiotics have important active attributes that could fulfill most of our basic nutritional and clinical supplementation requirements types. The microbes have shown positive responses to clinical treatment against several diseases and disorders, such as diarrhea associated with rotavirus, the IBS and food allergies types. The contribution of the probiotics in preventing and treatment of diabetes disease, obesity, cancer and diseases related to the pathogenic microbes is an exciting and rapidly advancing the investigations types. The dietary probiotic supplementation generally involves dairy products but the probiotics can be incorporated into non-dairy fermented food products types, presenting an alternative and more advantageous source in the process of evaluating the new probiotic strains types. Present clinical and nutritional evaluations have been successful in exposing some remarkable activities of particular probiotic strains. Specifically, regulation of energy in various catabolic and anabolic processes, acid and bile tolerance, ability to adhere to gut epithelial cells, to combat against pathogenic microorganisms, along with certain other properties, like their safety-enhancing property, serviceability as diet and beneficial supplements for the human being's health. The evaluation of new strains of the probiotics and their applicability in biomedical/clinical investigations, paving a new direction for exploration and exploitation of the probiotics aimed at improving the human being's health.

### References:

1. Shaltout, F.A.(2023): Viruses in Beef, Mutton, Chevon, Venison, Fish and Poultry Meat Products. Food Science & Nutrition Technology 8(4):1-10.Bartlett JG. Antibiotic-associated diarrhea. N Engl J Med. 2002;346(5):334–339.
2. Bartlett JG. Antibiotic-associated diarrhea. N Engl J Med. 2002;346(5):334–339.
3. Shaltout, F.A ., Riad,E.M ., and AbouElhassan, Asmaa , A(2017): prevalence Of Mycobacterium Tuberculosis In Imported cattle Offals And Its lymph Nodes. Veterinary Medical Journal -Giza (VMJG), 63(2): 115 – 122.
4. Szajewska H, Horvath A, Piwowarczyk A. Meta-analysis: the effects of *Saccharomyces boulardii* supplementation on *Helicobacter pylori* eradication rates and side effects during treatment. Aliment Pharmacol Ther. 2010;32(9):1069–1079.

5. Shaltout, F.A., Riad, E.M., and Asmaa Abou-Elhassan (2017): Prevalence Of Mycobacterium Spp. In Cattle Meat And Offal's Slaughtered In And Out Abattoir. Egyptian Veterinary medical Association, 77(2): 407 – 420.
6. Andrews JM, Tan M. Probiotics in luminal gastroenterology: the current state of play. Intern Med J. 2012;42(12):1287–1291.
7. Abd Elaziz, O., Fatin S. Hassanin, Fahim A. Shaltout and Othman A. Mohamed (2021): Prevalence of Some Foodborne Parasitic Affection in Slaughtered Animals in Local Egyptian Abattoir. Journal of Nutrition Food Science and Technology 2(3): 1-5.
8. Abd Elaziz, O., Fatin, S Hassanin, Fahim, A Shaltout, Othman, A Mohamed (2021): Prevalence of some zoonotic parasitic affections in sheep carcasses in a local abattoir in Cairo, Egypt. Advances in Nutrition & Food Science 6(2): 6(2): 25-31.
9. Bonnema AL, Kolberg LW, Thomas W, Slavin JL. Gastrointestinal tolerance of chicory inulin products. J Am Diet Assoc. 2010;110(6):865–868.
10. Al Shorman, A.A.M.; Shaltout, F.A. and Hilal, N (1999): Detection of certain hormone residues in meat marketed in Jordan. Jordan University of Science and Technology, 1st International Conference on Sheep and goat Diseases and Productivity, 23-25 October, 1999.
11. Ebeed Saleh, Fahim Shaltout, Essam Abd Elaal (2021); Effect of some organic acids on microbial quality of dressed cattle carcasses in Damietta abattoirs, Egypt. Damanhour Journal of Veterinary Sciences 5(2): 17-20.
12. Cencic A, Chingwaru W. (2010). The role of functional foods, nutraceuticals, and food supplements in intestinal health. Nutrients. 2(6):611–625.
13. Edris A, Hassanin, F. S; Shaltout, F.A., Azza H Elbaba and Nairoz M Adel (2017): Microbiological Evaluation of Some Heat Treated Fish Products in Egyptian Markets. EC Nutrition 12.3 (2017): 124-132.
14. Edris, A., Hassan, M.A., Shaltout, F.A. and Elhosseiny, S (2013): Chemical evaluation of cattle and camel meat. BENHA VETERINARY MEDICAL JOURNAL, 24 (2): 191-197.
15. Hempel S, Newberry SJ, Maher AR, Wang Z, Miles JN, Shanman R, Shekelle PG. Probiotics for the prevention and treatment of antibiotic-associated diarrhea: a systematic review and meta-analysis. JAMA. 2012;307(18):1959–1969.
16. Edris, A.M., Hassan, M.A., Shaltout, F.A. and Elhosseiny, S (2012): Detection of E.coli and Salmonella organisms in cattle and camel meat. BENHA VETERINARY MEDICAL JOURNAL, 24 (2): 198-204.
17. Edris A.M.; Hemmat M. I., Shaltout F.A.; Elshater M.A., Eman F.M.I. (2012): STUDY ON INCipient SPOILAGE OF CHILLED CHICKEN CUTS-UP. BENHA VETERINARY MEDICAL JOURNAL, VOL. 23, NO. 1, JUNE 2012: 81-86.
18. Pathak YV (Ed.) (2011) Handbook of nutraceuticals: ingredients, formulations, and applications (Vol. 1). CRC Press
19. Edris A.M.; Hemmat M.I.; Shaltout F.A.; Elshater M.A., Eman, F.M.I. (2012): CHEMICAL ANALYSIS OF CHICKEN MEAT WITH RELATION TO ITS QUALITY. BENHA VETERINARY MEDICAL JOURNAL, 23 (1): 87-92.
20. Hatakka K, Holma R, El-Nezami H, Suomalainen T, Kuisma M, Saxelin M, Korpela R. The influence of Lactobacillus rhamnosus LC705 together with Propionibacterium freudenreichii ssp. shermanii JS on potentially carcinogenic bacterial activity in human colon. Int J Food Microbiol. 2008;128(2):406–410.
21. Edris, A.M.; Shaltout, F.A. and Abd Allah, A.M. (2005): Incidence of Bacillus cereus in some meat products and the effect of cooking on its survival. Zag. Vet. J.33 (2):118-124.
22. Edris, A.M.; Shaltout, F.A. and Arab, W.S. (2005): Bacterial Evaluation of Quail Meat. Benha Vet. Med.J.16 (1):1-14.
23. Veerappan GR, Betteridge J, Young PE. Probiotics for the treatment of inflammatory bowel disease. Curr Gastroenterol Rep. 2012;14(4):324–333.
24. Edris, A.M.; Shaltout, F.A.; Salem, G.H. and El-Toukhy, E.I. (2011): Incidence and isolation of Salmonellae from some meat products. Benha University, Faculty of Veterinary Medicine, Fourth Scientific Conference 25-27th May 2011 Veterinary Medicine and Food Safety ) 172-179 benha, Egypt.
25. Macfarlane S, Macfarlane GT, Cummings JT. (2006). Review article: prebiotics in the gastrointestinal tract. Aliment Pharmacol Ther. 24(5):701–714.
26. Edris AA, Hassanin, F. S; Shaltout, F.A., Azza H Elbaba and Nairoz M Adel. (2017): Microbiological Evaluation of Some Heat Treated Fish Products in Egyptian Markets. EC Nutrition 12.3 (2017): 134-142.
27. Edris, A.M.; Shaltout, F.A.; Salem, G.H. and El-Toukhy, E.I. (2011): Plasmid profile analysis of Salmonellae isolated from some meat products. Benha University, Faculty of Veterinary Medicine, Fourth Scientific Conference 25-27th May 2011 Veterinary Medicine and Food Safety ) 194-201 benha, Egypt.
28. Isolauri E, Kirjavainen PV, Salminen S. Probiotics: a role in the treatment of intestinal infection and inflammation. Gut. 2002;50(suppl 3):iii54–iii59.
29. Ragab A, Abobakr M. Edris, Fahim A.E. Shaltout, Amani M. Salem (2022): Effect of titanium dioxide nanoparticles and thyme essential oil on the quality of the chicken fillet. BENHA VETERINARY MEDICAL JOURNAL 41(2): 38-40.
30. Hassan, M.A, Shaltout, F. A, Arfa M.M, Mansour A.H and Saudi, K. R (2013): BIOCHEMICAL STUDIES ON RABBIT MEAT RELATED TO SOME DISEASES. BENHA VETERINARY MEDICAL JOURNAL 25(1):88-93.
31. Roshchina VV (2010) Evolutionary considerations of neurotransmitters in microbial, plant, and animal cells. in Microbial Endocrinol 17–52, Springer New York
32. Hassan, M.A and Shaltout, F.A. (1997): Occurrence of Some Food Poisoning Microorganisms In Rabbit Carcasses Alex. J. Vet.Science, 13(1):55-61.
33. Hassan M, Shaltout FA\* and Saqur N (2020): Histamine in Some Fish Products. Archives of Animal Husbandry & Dairy Science 2(1): 1-3.
34. Hord NG. Eukaryotic-microbiota crosstalk: potential mechanisms for health benefits of prebiotics and probiotics. Annu Rev Nutr. 2008;28:215–231.
35. Hassan, M.A and Shaltout, F.A. (2004): Comparative Study on Storage Stability of Beef, Chicken meat, and Fish at Chilling Temperature. Alex.J.Vet.Science, 20(21):21-30.
36. Hassan, M.A; Shaltout, F.A.; Arafa, M.M.; Mansour, A.H. and Saudi, K.R. (2013): Biochemical studies on rabbit meat related to some diseases. Benha Vet. Med.J.25 (1):88-93.
37. McFarland LV. Meta-analysis of probiotics for the prevention of antibiotic associated diarrhea and the treatment of Clostridium difficile disease. Am J Gastroenterol. 2006;101(4):812–822.
38. Hassan, M.A; Shaltout, F.A.; Maarouf, A.A. and El-Shafey, W.S. (2014): Psychrotrophic bacteria in frozen fish with special



- reference to pseudomonas species .Benha Vet. Med.J.27 (1):78-83.
39. Hassan, M.A ; Shaltout, F.A. ; Arafa ,M.M. ; Mansour , A.H. and Saudi, K.R.(2013): Bacteriological studies on rabbit meat related to some diseases Benha Vet. Med.J.25 (1):94-99.
  40. Narayan SS, Jalgaonkar S, Shahani S, Kulkarni VN. Probiotics: current trends in the treatment of diarrhoea. Hong Kong Med J. 2010;16(3):213–218.
  41. Hassanin, F. S; Hassan,M.A., Shaltout, F.A., Nahla A. Shawqy and 2Ghada A. Abd-Elhameed (2017): Chemical criteria of chicken meat.BENHA VETERINARY MEDICAL JOURNAL, 33(2):457-464.
  42. Hassanin, F. S; Hassan,M.A.; Shaltout, F.A. and Elrais-Amina, M(2014): CLOSTRIDIUM PERFRINGENS IN VACUUM PACKAGED MEAT PRODUCTS. BENHA VETERINARY MEDICAL JOURNAL, 26(1):49-53.
  43. Hassanien, F.S.; Shaltout, F.A.; Fahmey, M.Z. and Elsukkary, H.F.(2020): Bacteriological quality guides in local and imported beef and their relation to public health. Benha Veterinary Medical Journal 39: 125-129.
  44. Cashman K. Probiotics and calcium bioavailability. Curr Issues Intest Microbiol. 2003;4(1):21–32.
  45. Hassanin, F. S; Shaltout,F.A. and , Mostafa E.M(2013): Parasitic affections in edible offal. Benha Vet. Med.J.25 (2):34-39.
  46. Hassanin, F. S; Shaltout, F.A., Lamada, H.M., Abd Allah, E.M. (2011): THE EFFECT OF PRESERVATIVE (NISIN) ON THE SURVIVAL OF LISTERIA MONOCYTOGENES. BENHA VETERINARY MEDICAL JOURNAL (2011)-SPECIAL ISSUE [I]: 141-145.
  47. Zhang MM, Cheng JQ, Lu YR, Yi ZH, Yang P, Wu XT. Use of pre-, pro-and synbiotics in patients with acute pancreatitis: a meta-analysis. World J Gastroenterol: WJG. 2010;16(31):3970.
  48. Khattab, E.,Fahim Shaltout and Islam Sabik (2021): Hepatitis A virus related to foods. BENHA VETERINARY MEDICAL JOURNAL 40(1): 174-179.
  49. Saad M. Saad, Fahim A. Shaltout, Amal A. A. Farag & Hashim F. Mohammed (2022): Organophosphorus Residues in Fish in Rural Areas. Journal of Progress in Engineering and Physical Science 1(1): 27-31.
  50. Saif,M., Saad S.M., Hassanin, F. S; Shaltout FA, Marionette Zaghoul (2019): Molecular detection of enterotoxigenic Staphylococcus aureus in ready-to-eat beef products. Benha Veterinary Medical Journal 37 (2019) 7-11.
  51. Matsumoto S, Hara T, Hori T, Mitsuyama K, Nagaoka M, Tomiyasu N, Sata M. Probiotic Lactobacillus-induced improvement in murine chronic inflammatory bowel disease is associated with the down-regulation of pro-inflammatory cytokines in lamina propria mononuclear cells. Clin Exp Immunol. 2005;140(3):417–426.
  52. Saif,M., Saad S.M., Hassanin, F. S; Shaltout, F.A., Marionette Zaghoul (2019); Prevalence of methicillin-resistant Staphylococcus aureus in some ready-to-eat meat products. Benha Veterinary Medical Journal 37 (2019) 12-15.
  53. Farag, A. A., Saad M. Saad<sup>1</sup>, Fahim A. Shaltout<sup>1</sup>, Hashim F. Mohammed (2023 a): Studies on Pesticides Residues in Fish in Menofia Governorate. Benha Journal of Applied Sciences., 8(5): 323-330.
  54. Szymański H, Pejcz J, Jawień M, Chmielarczyk A, Strus M, Heczko PB. Treatment of acute infectious diarrhoea in infants and children with a mixture of three Lactobacillus rhamnosus strains– a randomized, double-blind, placebo-controlled trial. Aliment Pharmacol Ther. 2006;23(2):247–253.
  55. Farag, A. A., Saad M. Saad<sup>1</sup>, Fahim A. Shaltout<sup>1</sup>, Hashim F. Mohammed (2023 b): Organochlorine Residues in Fish in Rural Areas. Benha Journal of Applied Sciences, 8 (5): 331-336.
  56. Kuo SM. The interplay between fiber and the intestinal microbiome in the inflammatory response. Adv Nutr: Intern Rev J. 2013;4(1):16–28.
  57. Shaltout, F.A., Mona N. Hussein, Nada Kh. Elsayed (2023): Histological Detection of Unauthorized Herbal and Animal Contents in Some Meat Products. Journal of Advanced Veterinary Research 13(2): 157-160.
  58. Schley PD, Field CJ. The immune-enhancing effects of dietary fibres and prebiotics. Br J Nutr. 2002;87(S2): S221–S230.
  59. Shaltout, F. A., Heikal, G. I., Ghanem, A. M. (2022): Mycological quality of some chicken meat cuts in Gharbiya governorate with special reference to Aspergillus flavus virulent factors. benha veteriv medical journal veterinary 42(1): 12-16.
  60. Shaltout, F.A., Ramadan M. Salem, Eman M. Eldiasty, Fatma A. Diab (2022): Seasonal Impact on the Prevalence of Yeast Contamination of Chicken Meat Products and Edible Giblets. Journal of Advanced Veterinary Research 12(5): 641-644.
  61. Liong MT, Dunshea FR, Shah NP. Effects of a synbiotic containing Lactobacillus acidophilus ATCC 4962 on plasma lipid profiles and morphology of erythrocytes in hypercholesterolaemic pigs on high-and low-fat diets. Br J Nutr. 2007;98(4):736–744.
  62. Shaltout, F.A., Abdelazez Ahmed Helmy Barr and Mohamed Elsayed Abdelaziz (2022): Pathogenic Microorganisms in Meat Products. Biomedical Journal of Scientific & Technical Research 41(4): 32836-32843.
  63. Scaldaferrri F, Gerardi V, Lopetuso LR, Del Zompo F, Mangiola F, Boškoski I, Gasbarrini, A (2013) Gut microbial flora, prebiotics, and probiotics in IBD: their current usage and utility. BioMed Res Intern 2013.
  64. Shaltout, F.A., Thabet, M.G. and Koura, H.A. (2017). Impact of Some Essential Oils on the Quality Aspect and Shelf Life of Meat. J Nutr Food Sci., 7: 647.
  65. Shaltout, F.A.,, Islam Z. Mohammed<sup>2</sup>, El -Sayed A. Afify(2020): Bacteriological profile of some raw chicken meat cuts in Ismailia city, Egypt.Benha Veterinary Medical Journal 39 (2020) 11-15.
  66. Pokusaeva K, Fitzgerald GF, van Sinderen D. Carbohydrate metabolism in Bifidobacteria. Gen Nutr. 2011;6(3):285–306.
  67. Shaltout, F.A.,Islam, Z. Mohammed<sup>2</sup>, El -Sayed A. Afify(2020): Detection of E. coli O157 and Salmonella species in some raw chicken meat cuts in Ismailia province, Egypt. Benha Veterinary Medical Journal 39 (2020) 101-104.
  68. Shaltout, F.A., E.M. El-diasty and M. A. Asmaa- Hassan (2020): HYGIENIC QUALITY OF READY TO EAT COOKED MEAT IN RESTAURANTS AT Cairo. Journal of Global Biosciences 8(12): 6627-6641.
  69. Romeo J, Nova E, Wärnberg J, Gómez-Martínez S, DíazLigia LE, Marcos A. Immunomodulatory effect of fibres, probiotics and synbiotics in different life-stages. Nutr Hosp. 2010;25(3):341–9.
  70. Shaltout, F.A., Marrionet Z. Nasief , L. M. Lotfy , Bossi T. Gamil(2019): Microbiological status of chicken cuts and its products. Benha Veterinary Medical Journal 37 (2019) 57-63.

71. Shaltout, F.A.(2019): Poultry Meat. *Scholarly Journal of Food and Nutrition* 22 1-2.
72. Niittynen L, Kajander K, Korpela R. Galacto-oligosaccharides and bowel function. *Scand J Food Nutr.* 2007;51(2):62.
73. Panda AK, Rao SVR, Raju MV, Sharma SR. Dietary supplementation of *Lactobacillus sporogenes* on performance and serum biochemico-lipid profile of broiler chickens. *J Poultr Sci.* 2006;43(3):235–240.
74. Shaltout, F.A. (2019): Food Hygiene and Control. *Food Science and Nutrition Technology* 4(5): 1-2.
75. Hassanin, F. S; Shaltout, F.A., Seham N. Homouda and Safaa M. Araakeb (2019): Natural preservatives in raw chicken meat. *Benha Veterinary Medical Journal* 37 (2019) 41-45.
76. Kelesidis T, Pothoulakis C. Efficacy and safety of the probiotic *Saccharomyces boulardii* for the prevention and therapy of gastrointestinal disorders. *Ther Adv Gastroenterol.* 2012;5(2):111–125.
77. Hazaa,W., Shaltout, F.A., Mohamed El-Shate(2019): Prevalence of some chemical hazards in some meat products. *Benha Veterinary Medical Journal* 37 (2) 32-36.
78. Hazaa,W, Shaltout, F.A., Mohamed El-Shater(2019): Identification of Some Biological Hazards in Some Meat Products. *Benha Veterinary Medical Journal* 37 (2) 27-31.
79. Parnell JA, Reimer RA. Effect of prebiotic fibre supplementation on hepatic gene expression and serum lipids: a dose–response study in JCR: LA-cp rats. *Br J Nutr.* 2010;103(11):1577–1584.
80. Gaafar,R. , Hassanin, F. S; Shaltout, F.A., Marionette Zaghoul (2019): Molecular detection of enterotoxigenic *Staphylococcus aureus* in some ready to eat meat-based sandwiches. *Benha Veterinary Medical Journal* 37 (2) 22-26.
81. Webb GP (2011) *Dietary supplements and functional foods.* John Wiley and Sons
82. Gaafar,R., Hassanin, F. S; Shaltout, F.A., Marionette Zaghoul( 2019): Hygienic profile of some ready to eat meat product sandwiches sold in Benha city, Qalubiya Governorate, Egypt. *Benha Veterinary Medical Journal* 37 (2) 16-21.
83. Artiss JD, Brogan K, Brucal M, Moghaddam M, Jen KLC. The effects of a new soluble dietary fiber on weight gain and selected blood parameters in rats. *Metabolism.* 2006;55(2):195–202.
84. Saad S.M. , Shaltout, F.A., Nahla A Abou Elroos, Saber B El-nahas( 2019): Antimicrobial Effect of Some Essential Oils on Some Pathogenic Bacteria in Minced Meat. *J Food Sci Nutr Res.* 2019; 2 (1): 012-020.
85. McFarland LV. Meta-analysis of probiotics for the prevention of traveler'sdiarrhea. *Travel Med Infect Dis.* 2007;5(2):97–105.
86. Saad S.M., Shaltout, F.A., Nahla A Abou Elroos2 and Saber B El-nahas (2019): Incidence of *Staphylococci* and *E. coli* in Meat and Some Meat Products. *EC Nutrition* 14.6.
87. Watson RR, Preedy VR (2010) *Bioactive foods in promoting health: probiotics and prebiotics.* Academic Press
88. Saad S.M., Hassanin, F. S.; Shaltout, F.A., Marionette Z Nassif, Marwa Z Seif. (2019): Prevalence of Methicillin-Resistant *Staphylococcus Aureus* in Some Ready-to-Eat Meat Products. *American Journal of Biomedical Science & Research* 4(6):460-464.
89. Kaur N, Gupta AK. (2002). Applications of inulin and oligofructose in health and nutrition. *J Biosci.* 27(7):703–714.
90. Shaltout, Fahim (2019): Pollution of Chicken Meat and Its Products by Heavy Metals. *Research and Reviews on Healthcare: Open Access Journal*, 4, 3(381-3382).
91. Sheil B, Shanahan F, O'Mahony L. Probiotic effects on inflammatory bowel disease. *J Nutr.* 2007;137(3):819S–824S.
92. Shaltout, F. A.; E.M EL-diasty; M. S. M Mohamed (2018): Effects of chitosan on quality attributes fresh meat slices stored at 4 C. *BENHA VETERINARY MEDICAL JOURNAL, VOL. 35, NO. 2: 157-168.*
93. Shaltout and Abdel-Aziz, 2004: *Salmonella enterica* serovar Enteritidis in poultry meat and their epidemiology. *Vet. Med. J. Giza*, 52 (2004), pp. 429-436.
94. Langen LV, Mirjam AC, Dieleman LA. Prebiotics in chronic intestinal inflammation. *Inflamm Bowel Dis.* 2009;15(3):454–462.
95. Shaltout, F.A., Hala F El-Shorah, Dina I El Zahaby, Lamiaa M Lotfy (2018):Bacteriological Profile of Chicken Meat Products. *SciFed Food & Dairy Technology Journal*, 2:3.
96. Swennen K, Courtin CM, Delcour JA. Non-digestible oligosaccharides with prebiotic properties. *Crit Rev Food Sci Nutr.* 2006;46(6):459–471.
97. Shaltout, F.A., Mohamed, A.H. El-Shater., Wafaa Mohamed Abd El-Aziz (2015): Bacteriological assessment of Street Vended Meat Products sandwiches in kalyobia Governorate. *BENHA VETERINARY MEDICAL JOURNAL*, 28( 2):58-66.
98. Loscalzo J. Lipid metabolism by gut microbes and atherosclerosis. *Circ Res.* 2011;109(2):127–129.
99. Shaltout, F.A., Mohamed A El shatter and Heba M Fahim(2019): Studies on Antibiotic Residues in Beef and Effect of Cooking and Freezing on Antibiotic Residues Beef Samples. *Scholarly Journal of Food and Nutritionm* 2(1) 1-4
100. Shaltout FA, Zakaria IM and Nabil ME.(2018): Incidence of Some Anaerobic Bacteria Isolated from Chicken Meat Products with Special Reference to *Clostridium perfringens*. *Nutrition and Food Toxicology* 2.5 (2018): 429-438.
101. Kruis W, Chrubasik S, Boehm S, Stange C, Schulze J. A double-blind placebo-controlled trial to study therapeutic effects of probiotic *Escherichia coli* Nissle 1917 in subgroups of patients with irritable bowel syndrome. *Int J Color Dis.* 2012;27(4):467–474.
102. Shaltout FA, Ahmed A A Maarouf and Mahmoud ES Elkhoully. (2017): Bacteriological Evaluation of Frozen Sausage. *Nutrition and Food Toxicology* 1.5; 174-185.
103. Shaltout FA, El-Toukhy EI and Abd El-Hai MM. (2019): Molecular Diagnosis of *Salmonellae* in Frozen Meat and Some Meat Products. *Nutrition and Food Technology Open Access* 5(1): 1-6.
104. Nguyen TDT, Kang JH, Lee MS. Characterization of *Lactobacillus plantarum* PH04, a potential probiotic bacterium with cholesterol-lowering effects. *Int J Food Microbiol.* 2007;113(3):358–361.
105. Shaltout, F.A., A.M.Ali and S.M.Rashad (2016): Bacterial Contamination of Fast Foods. *Benha Journal of Applied Sciences (BJAS)* 1 (2)45-51.
106. Shaltout, F.A., Zakaria. I. M., Jehan Eltanani, Asmaa. Elmelegy (2015): Microbiological status of meat and chicken received to University student hostel. *BENHA VETERINARY MEDICAL JOURNAL*, 29(2):187-192, DECEMBER, 2015.
107. Erejuwa OO, Sulaiman SA, Wahab MSA. Modulation of gut microbiota in the management of metabolic disorders: the prospects and challenges. *Int J Mol Sci.* 2014;15(3):4158–4188.



108. Saad,S.M.;Edris, A.M.; Shaltout,F.A. and Edris, Shima(2012): Isolation and identification of salmonellae and E.coli from meat and poultry cuts by using A.multiplex PCR. Benha Vet. Med.J.special issue 16-26.
109. Saad, S.M. and Shaltout, F.A. (1998):Mycological Evaluation of camel carcasses at Kalyobia Abattoirs. Vet.Med.J. Giza,46(3):223-229.
110. Lee JH, Nam SH, Seo WT, Yun HD, Hong SY, Kim MK, Cho KM. The production of surfactin during the fermentation of cheonggukjang by potential probiotic *Bacillus subtilis* CSY191 and the resultant growth suppression of MCF-7 human breast cancer cells. *Food Chem.* 2012;131(4):1347–1354.
111. Saad S.M., Shaltout, F.A., Nahla A Abou Elroos, Saber B El-nahas. 2019: Antimicrobial Effect of Some Essential Oils on Some Pathogenic Bacteria in Minced Meat. *J Food Sci Nutr Res.* 2019; 2 (1): 012-020.
112. Saad S.M., Hassanin, F. S; Shaltout, F.A., Marionette Z Nassif, Marwa Z Seif.(2019): Prevalence of Methicillin-Resistant *Staphylococcus Aureus* in Some Ready-to-Eat Meat Products. *American Journal of Biomedical Science & Research* 4(6):460-464.
113. Delcenserie V, Martel D, Lamoureux M, Amiot J, Boutin Y, Roy D. (2008). Immunomodulatory effects of probiotics in the intestinal tract. *Curr Issues Mol Biol.* 10(1/2):37.
114. Shaltout FA, Riad EM, TES Ahmed and AbouElhassan A. (2017): Studying the Effect of Gamma Irradiation on Bovine Offal's Infected with *Mycobacterium tuberculosis* Bovine Type. *Journal of Food Biotechnology Research* 1 (6): 1-5.
115. Saulnier D, Spinler JK, Gibson GR, Versalovic J. Mechanisms of probiosis and prebiosis: considerations for enhanced functional foods. *Curr Opin Biotechnol.* 2009;20(2):135–141.
116. Shaltout FA, Zakaria IM and Nabil ME. (2018): Incidence of Some Anaerobic Bacteria Isolated from Chicken Meat Products with Special Reference to *Clostridium perfringens*. *Nutrition and Food Toxicology* 2.5 (2018): 429-438.
117. Howarth GS. Inflammatory bowel disease, a dysregulated host-microbiota interaction: are probiotics a new therapeutic option. *J Gastroenterol Hepatol.* 2008;23(12):1777–1779.
118. Shaltout FA, Mohamed, A.Hassan and Hassanin, F. S(2004): THERMAL INACTIVATION OF ENTEROHAEMORRHAGIC *ESCHERICHIA COLI* O157:H7 AND ITS SENSITIVITY TO NISIN AND LACTIC ACID CULTURES. 1rst Ann. Confr. , FVM., Moshtohor, Sept, 2004.
119. Shaltout FA, El-diahy, E.M. ;Elmesalmy, M. and Elshaer, M.(2014): Study on fungal contamination of some chicken meat products with special reference to 2 the use of PCR for its identification . Conference, Veterinary Medical Journal – Giza vol. December 2014/12/17 vol.60: 1-10.
120. Both E, Gyenge L, Bodor Z, Gyorgy E, Lanyi S, Abraham B. Intensification of probiotic microorganisms viability by microencapsulation using ultrasonic atomizer. *UPB Buletin Stiintific Series B: Chem Mater Sc.* 2012;74(1):27–32.
121. Shaltout, F.A.(2002): Microbiological Aspects of Semi-cooked chicken Meat Products. *Benha Veterinary Medical Journal*13,2,: 15-26.
122. Shaltout FA, Thabet, M.G2 and Hanan, A. Koura3. (2017): Impact of some essential oils on the quality aspect and shelf life of meat.BENHA VETERINARY MEDICAL JOURNAL, 33, (2): 351-364.
123. Kim HJ, Vazquez Roque MI, Camilleri M, Stephens D, Burton DD, Baxter K, Zinsmeister AR. A randomized controlled trial of a probiotic combination VSL# 3 and placebo in irritable bowel syndrome with bloating. *Neuro-gastroenterology and Motility.* 2005;17(5):687–696.
124. Shaltout FA, Mohammed Farouk; Hosam A.A. Ibrahim and Mostafa E.M. Afifi4.2017: Incidence of Coliform and *Staphylococcus aureus* in ready to eat fast foods. *BENHA VETERINARY MEDICAL JOURNAL,* 32( 1): 13 - 17, MARCH, 2017.
125. Oberreuther-Moschner DL, Jahreis G, Rechkemmer G, Pool-Zobel BL. Dietary intervention with the probiotics *Lactobacillus acidophilus* 145 and *Bifidobacterium longum*913 modulates the potential of human faecal water to induce damage in HT29clone19A cells. *Br J Nutr.* 2004;91(06):925–932.
126. Shaltout, F.A., Zakaria, I.M., Nabil, M.E.(2017): Detection and typing of *Clostridium perfringens* in some retail chicken meat products.BENHA VETERINARY MEDICAL JOURNAL., 33(2):283-291.
127. Shaltout, F.A. (1992): Studies on Mycotoxins in Meat and Meat by Products. M.V.Sc Thesis Faculty of Veterinary Medicine, Moshtohor, Zagazig University Benha branch.
128. Jonkers D, Penders J, Masclee A, Pierik M. Probiotics in the management of inflammatory bowel disease. *Drugs.* 2012;72(6):803–823.
129. Shaltout, F.A.(1996): Mycological And Mycotoxicological profile Of Some Meat products. Ph.D.Thesis, Faculty of Veterinary Medicine, Moshtohor, Zagazig University Benha branch.
130. Shaltout, F.A. (1998): Proteolytic Psychrotrophes in Some Meat products. *Alex. Vet. Med. J.*14 (2):97-107.
131. Lindsay JO, Whelan K, Stagg AJ, Gobin P, Al-Hassi HO, Rayment N, Forbes A. Clinical, microbiological, and immunological effects of fructo-oligosaccharide in patients with Crohn's disease. *Gut.* 2006;55(3):348–355.
132. Shaltout, F.A. (1999): Anaerobic Bacteria in Vacuum Packed Meat Products. *Benha Vet. Med.J.*10 (1):1-10.
133. Peña AS. Intestinal flora, probiotics, prebiotics, synbiotics and novel foods. *Rev Esp Enferm Dig.* 2007;99(11):653.
134. Shaltout, F.A.(2000):Protozoal Foodborne Pathogens in some Meat Products. *Assiut Vet. Med. J.* 42 (84):54-59.
135. Shaltout, F.A.(2001): Quality evaluation of sheep carcasses slaughtered at Kalyobia abattoirs. *Assiut Veterinary Medical Journal,* 46(91):150-159.
136. Shaltout, F.A.(2002): Microbiological Aspects of Semi-cooked Chicken Meat Products. *Benha Vet.Med.J.* 13(2):15-26.
137. Homayouni A, Payahoo L, Azizi A (2012) Effects of probiotics on lipid profile: a review. *Am J Food Technol* 7(5)
138. Shaltout, F.A. (2003): *Yersinia Enterocolitica* in some meat products and fish marketed at Benha city.The Third international conference Mansoura 29-30 April.
139. Teitelbaum JE, Walker WA. Nutritional impact of pre-and probiotics as protective gastrointestinal organisms. *Annu Rev Nutr.* 2002;22(1):107–138.
140. Shaltout, F.A. (2009): Microbiological quality of chicken carcasses at modern Poultry plant. The 3rd Scientific Conference, Faculty of Vet. Med., Benha University, 1-3 january.

141. Tanaka Y, Kanazawa M, Fukudo S, Drossman DA. Biopsychosocial model of irritable bowel syndrome. *J Neurogastroenterol Motil.* 2011;17(2):131–139.
142. Shaltout, F.A. and Abdel Aziz ,A.M.(2004): Salmonella enterica Serovar Enteritidis in Poultry Meat and their Epidemiology .*Vet.Med.J.,Giza,52(3):429-436.*
143. Guglielmetti S, Mora D, Gschwender M, Popp K. Randomised clinical trial: Bifidobacterium bifidum MIMBb75 significantly alleviates irritable bowel syndrome and improves quality of life—a double-blind, placebo-controlled study. *Aliment Pharmacol Ther.* 2011;33(10):1123–1132.
144. Shaltout, F.A. and Abdel Aziz ,A.M.(2004): ESCHERICHIA COLI STRAINS IN SLAUGHTERED ANIMALS AND THEIR PUBLIC HEALTH IMPORTENCE. *J.Egypt. Vet. Med. Association* 64(2):7-21.
145. Shaltout, F.A., Amin, R., Marionet, Z., Nassif and Shima, Abdel-wahab(2014): Detection of aflatoxins in some meat products. *Benha veterinary medical journal, 27(2) :368-374.*
146. Hsu CK, Liao JW, Chung YC, Hsieh CP, Chan YC. Xylooligosaccharides and fructooligosaccharides affect the intestinal microbiota and precancerous colonic lesion development in rats. *J Nutr.* 2004;134(6):1523–1528.
147. Shaltout, F.A. and Afify, Jehan Riad, EM and Abo Elhasan, Asmaa, A.(2012): Improvement of microbiological status of oriental sausage. *Journal of Egyptian Veterinary Medical Association* 72(2):157-167.
148. Van den Abbeele P, Van de Wiele T, Grootaert C, Verstraete W, Gérard P, Bruneau A, Possemiers S (2010) Arabinoxylans and inulin modulate the luminal and mucosa-associated bacteria In vitro and In vivo (pp. 233–249). van der Kamp JW, M. Jones J, McCleary B. V, Topping DL (Eds.). Wageningen Academic Publishers: Waltham, MA, USA
149. Shaltout, F.A. and Daoud, J. R.(1996): Chemical analytical studies on rabbit meat and liver. *Benha Vet. Med.J.8 (2):17-27.*
150. Ley RE, Turnbaugh PJ, Klein S, Gordon JI. (2006). Microbial ecology: human gut microbes associated with obesity. *Nature.* 444(7122):1022–1023.
151. Shaltout, F.A. and Edris, A.M.(1999): Contamination of shawerma with pathogenic yeasts. *Assiut Veterinary Medical Journal, 40(64):34-39.*
152. Whisner CM, Martin BR, Schoterman MH, Nakatsu CH, McCabe LD, McCabe GP, Weaver CM. (2013). Galactooligosaccharides increase calcium absorption and gut bifidobacteria in young girls: a double-blind cross-over trial. *Br J Nutr.* 110(07):1292–1303.
153. Shaltout, F. A.; Eldiasty, E. and Mohamed, M.S.(2014): Incidence of lipolytic and proteolytic fungi in some chicken meat products and their public health significance. *Animal Health Research Institute: First International Conference on Food Safety and Technology 19-23 June 2014 Cairo Egypt* pages 79-89.
154. Kaufmann SH. (2008). Immunology's foundation: the 100-year anniversary of the Nobel Prize to Paul Ehrlich and Elie Metchnikoff. *Nat Immunol.* 9(7):705–712.
155. Shaltout, F.A.; Eldiasty, E.; Salem, R. and Hassan, Asmaa (2016): Mycological quality of chicken carcasses and extending shelf – life by using preservatives at refrigerated storage. *Veterinary Medical Journal -Giza (VMJG)62(3)1-7.*
156. Sudha MR, Chauhan P, Dixit K, Babu S, Jamil K. (2009). Probiotics as complementary therapy for hypercholesterolemia. *Biol Med.* 1(4):1–13.
157. Shaltout, F.A.; Salem, R. Eldiasty, E.; and Diab, Fatema. (2016): Mycological evaluation of some ready to eat meat products with special reference to molecular characterization. *Veterinary Medical Journal -Giza* 62(3)9-14.
158. Moeinian M, FarnazGhasemi-Niri S, Mozaffari S, Abdollahi M. Synergistic effect of probiotics, butyrate and L-Carnitine in treatment of IBD. *J Med Hypotheses Ideas.* 2013;7(2):50–53.
159. Shaltout, F. A.; Elshater, M. and Wafaa, Abdelaziz (2015): Bacteriological assessment of street vended meat products sandwiches in Kalyobia Governorate. *Benha Vet. Med.J.28 (2):58-66.*
160. Sudha R M, Bhonagiri S (2012) Efficacy of Bacillus coagulans strain unique is-2 in the treatment of patients with acute diarrhea. *Intern J Probiot Prebiot* 7(1)
161. Shaltout, F. A.; Gerges, M.T. and Shewail, A.A. (2018): Impact of Organic Acids and Their Salts on Microbial Quality and Shelf Life of Beef. *Assiut veterinary medical journal* 64(159): 164-177
162. Fotiadis CI, Stoidis CN, Spyropoulos BG, Zografos ED. (2008). Role of probiotics, prebiotics and synbiotics in chemoprevention for colorectal cancer. *World J Gastroenterol: WJG.* 14(42):6453.
163. Shaltout, F.A.; Ghoneim, A.M.; Essmail, M.E. and Yousseif, A.(2001): Studies on aflatoxin B1 residues in rabbits and their pathological effects. *J.Egypt. Vet. Med. Association* 61(2):85-103.
164. Stienstra R, Tack CJ, Kanneganti TD, Joosten LA, Netea MG. (2012). The inflammasome puts obesity in the danger zone. *Cell Metab.* 15(1):10–18.
165. Shaltout, F.A. and Hanan, M.T. El-Lawendy (2003): Heavy Metal Residues In Shawerma. *Beni-Suef Vet.Med.J.* 13(1):213-224.
166. Shaltout, F.A. and Hashim, M.F. (2002): Histamine in salted, Smoked and Canned Fish products. *Benha Vet. Med.J.13 (1):1-11.*
167. Hardy H, Harris J, Lyon E, Beal J, Foey AD. (2013). Probiotics, prebiotics and immunomodulation of gut mucosal defenses: homeostasis and immunopathology. *Nutrients.* 5(6):1869–1912.
168. Shaltout, F.A.; Hashim, M.F. and Elnahas, S.(2015): Levels of some heavy metals in fish (tilapia nilotica and Claris lazera) at Menufia Governorate. *Benha Vet. Med.J.29 (1):56-64.*
169. Van Immerseel F, Ducatelle R, De Vos M, Boon N, Van De Wiele T, Verbeke K, Flint HJ. (2010). Butyric acid-producing anaerobic bacteria as a novel probiotic treatment approach for inflammatory bowel disease. *J Med Microbiol.* 59(2):141–143.
170. Shaltout, F.A. and Ibrahim, H.M.(1997): Quality evaluation of luncheon and Alexandrian sausage. *Benha Vet. Med.J.10 (1):1-10.*
171. DeVrese M, Schrezenmeier J (2008) Probiotics, prebiotics, and synbiotics. in food biotechnology (pp. 1–66). Springer Berlin Heidelberg
172. Shaltout, F.A.; Nassif, M and Shakran, A (2014): Quality of battered and breaded chicken meat products. *Global Journal of Agriculture and Food Safety Science – 1(2) ISSN 2356-7775.*
173. Shaltout, F.A., Amani M. Salem, A. H. Mahmoud, K. A (2013): Bacterial aspect of cooked meat and offal at street vendors level. *Benha veterinary medical journal, 24(1): 320-328.*
174. Shaltout, F.A. and Salem, R.M.(2000): Moulds, aflatoxin B1 and Ochratoxin A in Frozen Livers and meat products. *Vet. Med. J.Giza* 48(3):341-346.

175. Harish K, Varghese T. (2006). Probiotics in humans–evidence-based review. *Calicut Med J.* 4(4):e3.
176. Yasser H. Al-Tarazi, A. Al-Zamil, Shaltout FA. and H. Abdel-Samei (2002). Microbiological status of raw cow milk marketed in northern Jordan. *AVMJ Volume 49 Issue 96 Pages 180-194*
177. Shaltout FA, Zakaria IM and Nabil ME. (2018): Incidence of Some Anaerobic Bacteria Isolated from Chicken Meat Products with Special Reference to *Clostridium perfringens*. *Nutrition and Food Toxicology*2(5):429-438.
178. Abrams SA, Griffin IJ, Hawthorne KM, Liang L, Gunn SK, Darlington G, Ellis KJ. (2005) A combination of prebiotic short-and long-chain inulin-type fructans enhances calcium absorption and bone mineralization in young adolescents. *Am J Clin Nutr.* 82(2):471–476.
179. Shaltout, F. A.; El-diasty, E.M. and Mohamed, M. S.(2014): Incidence of lipolytic and proteolytic fungi in some chicken meat products and their public health significance. 1st Scientific conference of food safety and Technology .2014, pp. 79-89.
180. Gershon MD (1998) *The second brain* (pp. 4–7). HarperCollins Publishers
181. Shaltout, F. A.; El-diasty, E.M.; Salem, R. M. and Asmaa, M. A. Hassan. (2016): Mycological quality of chicken carcasses and extending shelf -life by using preservatives at refrigerated storage. *Veterinary Medical Journal – Giza* ,62(3) :1-10.
182. Shaltout FA, R.M. Salem, E.M. El-Diasty and W.I.M. Hassan. (2019): Effect of Lemon Fruits and Turmeric Extracts on Fungal Pathogens in Refrigerated Chicken Fillet Meat. *Global Veterinaria* 21 (3): 156-160.
183. Shaltout FA, El-diasty, E.M.; Elmesalamy, M. and Elshaer, M.( 2014): Study on fungal contamination of some chicken meat products with special reference to 2 the use of PCR for its identification. Conference, *Veterinary Medical Journal – Giza* vol. December 2014/12/17 vol.60 1-10.
184. Chapman CMC, Gibson GR, Rowland I. Health benefits of probiotics: are mixtures more effective than single strains? *Eur J Nutr.* 2011;50(1):1–17.
185. Shaltout, F. A.; Salem, R. M; El-diasty, Eman and Fatema, A.H. Diab. (2016): Mycological evaluation of some ready to eat meat products with special reference to molecular characterization. *Veterinary Medical Journal – Giza.* 62(3): 9-14.
186. Shaltout FA, Ahmed, A.A. Maarouf, Eman, M.K. Ahmed(2018): Heavy Metal Residues in chicken cuts up and processed chicken meat products. *BENHA VETERINARY MEDICAL JOURNAL*, 34(1): 473-483.
187. Hill DR, Ryan ET (2008) Management of travellers' diarrhoea. *BMJ*, 337.
188. Shaltout ,F.A.; Hanan M. Lamada , Ehsan A.M. Edris.(2020): Bacteriological examination of some ready to eat meat and chicken meals. *Biomed J Sci & Tech Res.*, 27(1): 20461-20465.
189. Sobhy, Asmaa and Shaltout, Fahim(2020): Prevalence of some food poisoning bacteria in semi cooked chicken meat products at Qaliubiya governorate by recent Vitek 2 compact and PCR techniques. *Benha Veterinary Medical Journal* 38 (2020) 88-92.
190. Sobhy, Asmaa and Shaltout, Fahim (2020): Detection of food poisoning bacteria in some semi-cooked chicken meat products marketed at Qaliubiya governorate. *Benha Veterinary Medical Journal* 38 (2020) 93-96.
191. Boyle RJ, Robins-Browne RM, Tang ML. Probiotic use in clinical practice: what are the risks? *Am J Clin Nutr.* 2006;83(6):1256–1264.
192. Shaltout, F.A. (2024): Abattoir and Bovine Tuberculosis as A Reemerging Foodborne Diseases. *Clinical Medical Reviews and Report* 6(1):1-7.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

**Submit Manuscript**

DOI:10.31579/2637-8914/293

#### Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/nutrition-and-food-processing>