

Formalin Induces Alterations in Body and Lung Weights and Histological Structure of the Lung of Adult Male Albino Rats and Amelioration by Mint Aqueous Extract

Basma Amer El-Sayed Alsaid ^{1*}, Amina Ali Saleh Kalila ¹, Azab Elsayed Azab ²

¹Department of Zoology, Faculty of Science, University of Zawia, Libya.

²Physiology Department, Faculty of Medicine, Sabratha University, Libya.

***Corresponding Author:** Basma Amer El-Sayed Alsaid, Department of Zoology, Faculty of Science, University of Zawia, Libya.

Received date: April 08, 2025; **Accepted date:** April 21, 2025; **Published date:** May 02, 2025

Citation: El-Sayed Alsaid BA, Saleh Kalila AA, Azab E. Azab, (2025), Formalin Induces Alterations in Body and Lung Weights and Histological Structure of the Lung of Adult Male Albino Rats and Amelioration by Mint Aqueous Extract, *J, Biotechnology and Bioprocessing*, 6(3); DOI:10.31579/2766-2314/147

Copyright: © 2025, Basma Amer El-Sayed Alsaid. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Anatomy departments utilize formalin to harden museum specimens, and cadavers. Moreover, it is a component of pressed wood goods, paper, textile fibers, adhesives, plastics, carpeting, foam insulation, disinfectants, nail hardeners, and some finger paints. Additionally, it is used to stop germs from spoiling fish, fruits, milk, beverages, ice cream, sweetmeat, and spices. Because the respiratory epithelial cells are damaged and lose their ability to function, it leads in acute lung injury, a cytotoxic reaction in the respiratory system. The purpose of this study was to assess how formalin affected the adult male albino rats' body and lung weights as well as the histological structure of their lungs, and how this effect was mitigated by an aqueous extract of mint leaves.

Materials and Methods: For this experiment, thirty adult male albino rats were procured from the Libyan Medical Research Centre located in the city of Zawia. Three equal groups were formed out of the animals. As a control group, Group I was given drinking water. Group II was the formalin group; for 30 days, they were given solely formalin at a dose of 10 mg/kg BW. Groups III were treated as a formalin + mint group, receiving 10 mg/kg BW of formalin for one hour, followed by 30 days of daily dosages of 600 mg/kg BW of mint extract. Gastric tubes were used to deliver all samples orally. On the 30th day, the animals were weighed, anesthetized, then scarified by cervical dislocation, and dissected. This was done 24 hours after the last dose. The lungs were extracted, weighed, and some of the lung tissue was preserved for 72 hours in 10% neutral buffered formalin, followed by dehydration and paraffin embedding. Subsequently, 5µm slices were made and stained with hematoxylin–eosin for histological evaluations. The weight of the body and lungs was reported as mean ± SE. One-way analysis of variance was used to conduct multiple comparisons (ANOVA). A value of p less than 0.05 was deemed significant.

Results: The adult male albino rats' body weight significantly decreased ($P < 0.01$) following the administration of formalin and formalin+mint, according to the data. On the other hand, rats given formalin+mint for 30 days showed a significant ($P < 0.01$) increase in body weight as compared to the formalin group. Rats given formalin for 30 days showed an increase ($P < 0.01$) in lung weight as a percentage of body weight when compared to the controls. On the other hand, compared to the formalin group, adult male albino rats given formalin+mint for 30 days experienced a significant ($P < 0.01$) decrease in lung weight as a percentage of body weight. Control rats' lung sections underwent histological examination, which revealed normal pulmonary tissue architecture. When rats were given formalin, their lung tissues changed significantly in comparison to the control group. Co-administration of mint extract with formalin caused improvement in the lung tissues and restored the histoarchitecture to near normal as in the control group.

Conclusion: It can be concluded that administration of formalin to rats significantly decreased body weight and increased lung weight as a percentage of body weight and a severe histopathological changed in lung tissues in comparison to the control group. While, treatment with mint improved these changed.

Key Words: formalin; body weight; lung weight; histopathological alterations of the lung; mint aqueous extract; amelioration; adult male albino rats

Introduction

The word plastic originates from the Greek word "plastikos", which signifies 'ready to be formed into shifted shapes. Plastic is comprised of carbon, hydrogen, silicon, oxygen, chloride and nitrogen. For extraction of the fundamental materials of plastics oil, coal and petroleum gas are utilized. Plastics are comprised of connecting of monomers together by synthetic bonds.(Kale, Deshmukh, Dudhare, & Patil, 2015)Worldwide manufactured plastic creation is 140 million tons for every year, and this rate has been consistently expanding since the 1930's. It is assessed that 11% of landfill volume is comprised of waste plastics.(Norton, 2012)

Degradation is the cycle where in any substance is separating into isolated parts or components. Decay or pulverization of contaminant atoms by the activity of the chemical discharged by microorganisms is known as biodegradation. Any physical or concoction change in polymer are because of ecological factors, for example, light, heat, dampness, synthetic conditions and organic action is named as degradation of plastic(Kale et al., 2015).The natural corruption of polymeric substances is an unpredictable cycle including a few resulting steps instigated by the activity of proteins. Polymer corruption happens under high-impact and anaerobic condition(Devi et al., 2016).biodegradation of plastics relies upon both ecological variables temperature, dampness, oxygen, pH) and the substance structure of the polymer. Biodegradable polymers normally contain ester, amide, or carbonate hydrolysable bonds in the polymer spine. The nearness of these hydrolysable practical gatherings expands the powerlessness to biodegradation. Different components that influence biodegradability are crystallinity, sub-atomic weight, and, on account of copolymers, the copolymer creation.(Vaverková, Adamcová, Kotovicová, Toman, & S, 2014)

There are two gatherings of plastics based on biodegradability, i.e., non-biodegradable plastics and biodegradable plastics. **Non-biodegradable plastics:** Their atomic weight is high because of the broad redundancy of little monomer units. These plastics are profoundly steady and don't promptly go into the degradation patterns of the biosphere. Non-biodegradable plastics incorporate a significant number of the routinely utilized plastics like PVC, PP, PS, PET, PUR, and PE. **Biodegradable plastics:** Both bio-based and fossil-based polymers can be remembered for biodegradable plastics relying on the level of biodegradability and microbial assimilation.(Ahmed et al., 2018)

Fungi are the decomposers in the worldwide pattern of life and demise. They are normally there to accomplish the work when anything creature, plant, or even non-living article is fit to be separated again into its sub-atomic constituents. Fungi are found in soil, in new and ocean water, inside the collections of plants and creatures, and going through the air as spores. While they regularly are discovered working along with microbes and a variety of microorganisms, it is fungi that can particularly deal with separating the absolute biggest particles present in nature(Norton, 2012)Fungi are known



for their decent variety and amazing capacity to corrupt unpredictable and industrious characteristic materials, for example, lignin, chitin, and microcrystalline cellulose. Bacteria and fungi can develop under naturally focused on conditions, for example, low supplement availability(Atagana & Biotechnology, 2004)the open door for fungi to advance in their essence is progressing. the occupant fungi expanded their mycelial biomass, and that the network moved towards strength by three genera: Graphium, Fusarium, and Penicillium(Norton, 2012). Plastic corrupting organisms were disconnected from an assortment of assets, for example, rhizosphere soil of mangroves, polythene covered in the dirt, marine water, plastic and soil at the dumping destinations(Kale et al., 2015).Microorganisms are associated with the corruption and weakening of both engineered and characteristic polymers. So as to utilize such materials as a carbon and vitality source, microorganisms have built up an extraordinary strategy. Microorganisms, for example, bacteria and fungi are associated with the degradation of plastics.(Devi et al., 2016)

Four methods Mechanical, Chemical, physical and Biochemical are utilized for corruption of plastics Physical and mechanical procedures Leave their hurtful impacts on Environment and life of people, and these strategies additionally taint the earth again in general, Microbial catalysts liable for the degradation different kinds of plastics portray the substrates that use the plastics as carbon and vitality sources and aides in biodegradation. Microbial compounds initiate the pace of biodegradation of plastics successfully without making any damage nature and no hurtful intermediates or side-effects are produced during PHA corruption. Truth be told, 3-hydroxybutyrate is found in all higher creatures as blood plasma. Hence, PHAs have been considered for clinical applications, including long haul controlled medication discharge, careful pins, stitches, and bone and vein replacement(Devi et al., 2016).

Review Of Literature

Plastics are produced using hydrocarbon monomers. They are delivered by artificially altering characteristic substances or are combined from inorganic and natural crude materials. Based on their physical qualities, plastics are generally separated into **thermosets**, **elastomers**, and **thermoplastics**. These gatherings contrast essentially with respect to sub-atomic structure.(Shimao, 2001)

Most of the plastics are **thermoplastic**; that is, when the plastic is shaped it very well may be warmed and changed more than once. The other gathering is **thermosets**, which can't be re-dissolved. Manufactured plastics are generally utilized in pressing of items, for example, pharmaceuticals, food, beautifiers, synthetic compounds, and cleansers. The most generally utilized plastics in pressing are PE (LDPE, HDPE, MDPE, and LLDPE), PP, PS, PVC, PU, PET, polybutylene terephthalate, nylons, etc. **elastomers** also cannot be reshaped by heating.(Kale et al., 2015)

No.	Type of Plastic	Molecular Structure	Characteristics and Applications
1.	Thermosets		Thermosets are hard and have an exceptionally close fit, expanded sub-atomic structure. Relieving continues during forming, after which it is not, at this point conceivable to shape the material by warming.Further moulding may then just be performed by machining. Thermosets are utilized, for instance, to make light switches.
2.	Elastomers		Elastomers additionally have a cross-connected structure, they have a looser work than thermosets, offering ascend to a level of versatility.When moulded, elastomers additionally can't be reshaped by warming. Elastomers are utilized, for instance, to deliver car tires.


3.	Thermoplastics		Thermoplastics have a straight or stretched sub-atomic structure which decides their quality and warm conduct; they are adaptable at standard temperatures. At approx. 120–180°C, thermoplastics become a pale/fluid mass. The thermoplastics are utilized, for instance, in bundling applications.
----	----------------	---	---

Table 1: Characteristics and Applications of Plastics

During the 1980s, researchers began to investigate whether plastics could be intended to get vulnerable to microbial assault, making them degradable in a microbial dynamic condition. Biodegradable plastics opened the path for new contemplations of waste administration procedures.(Kaseem, Hamad, & Deri, 2012)

Methods of Plastics Degradation

The development of auxiliary homogeneities and arrangement of new utilitarian gatherings likewise happens during polymer degradation(Pospíšil & Nešpůrek, 1997). Depending upon the idea of the causing operators, polymer degradation has been named **Thermo degradation, Thermo-oxidative degradation, Thermal degradation, Photodegradation, UV degradation and oxidation Degradation.**(Wackett & Hershberger, 2001)




Figure 1: Types of plastic degradation

Thermo degradation implies the corruption of polymer by heat vitality. **Thermo-oxidative degradation** for the most part gets uphold from oxygen of the environment and known as thermo-oxidative degradation The underlying phase of degradation is the cycle of burst in the obligations of macromolecules bringing about radical destinations. **Thermal degradation** for the most part includes changes to sub-atomic load of polymer. **Photodegradation** The degradation that is completed within the sight of light is named as "photo degradation." It is started first by the retention of light vitality by the suitable gathering present in the polymer atom. The light ingestion brings about the scission of the polymer atom at a suitable situation of the chain prompting the transformation to littler sections. In this way, photodegradable polymers require either an in-assembled photograph responsive gathering in the chain or an added substance one.

Photodegradation of polymer incorporates UV degradation and oxidation Degradation, the UV light is utilized to degrade the final product. Oxidation, in this cycle heat is utilized to separate the plastic. Numerous engineered polymers are impervious to concoction and physical degradation. Both the thermal and physical strategies for degradation diminish the atomic load of the plastic and permit it to biodegrade.(Kyriako, Briassoulis, & Environment, 2007)

Biodegradation of Natural Plastics by Fungi

Polyhydroxyalkanoates are straight polyesters created in nature by bacterial maturation of sugar and lipids(Williams & Peoples, 1996). By and large, no destructive intermediates or side-effects are created during PHA degradation actually, 3-hydroxybutyrate is found in all higher creatures as blood plasma. A number of oxygen consuming and anaerobic organisms that degrade PHA

Takagi, 2006). Past discoveries have revealed a *Streptomyces* strain, *Streptovorticillium kashmeriense* AF1, equipped for degrading PHB and PHBV, several vigorous and anaerobic PHB-corrupting microorganisms have been detached from soil and different follows, cavities, and depressions as seen on the gouged surface of PHBV films showing that the debasement was a purposeful impact of a microbial consortium colonizing the film surface, including organisms, microbes, and actinomycetes. Various unpredictable disintegration pits have additionally been seen on the outside of PHA by *Comamonas* sp.(Shah, Hasan, Hameed, Ahmed, & biodegradation, 2007)

No	Biodegradable plastics	Applications
1.	PGA	Used for subcutaneous structure, intracutaneous closures, abdominal and thoracic surgeries.
2.	PHB	Manufacturing disposable utensils. Also used medical applications, it can also be used for drug delivery.
3.	PLA	Packaging and paper coatings, other possible markets include sustained release systems for pesticides and fertilizers, much films, and compost bag.
4.	PCL	Used in housing applications, drugs encapsulation, act as a scaffold for tissue repair via tissue engineering, in root canal filling etc
5.	PHA	Used for strutures, surgical mesh, repair patches, slings, cardiovascular patches, orthopedic pins, spinal fusion cages, implant meterials, skin substitutes, wound dressings. etc
6.	HV	Used in paper and film coatings, therapeutic drug delivery of worm medicine for cattle, and sustained release systems for pharmaceutical drugs and insecticides.
7.	PVOH	Packaging and bagging applications which dissolve in water to release. Products such as laundry detergent, pesticides, and hospitals washable.
8.	PVAc	Adhesives, the packaging applications include cardboard manufacture, paper bags, paper liminations, tube winding, and remoisten able label.

Table 2: Natural biodegradation plastics

Microorganisms Such as (fungi) are undeniably fit to the assignment of contaminant decimation since they have catalysts that permit them to utilize ecological




Figure 2: Different Mechanisms and Role of Fungi in Biodegradation of plastics

and prompts a populace that is better ready to use the contaminant (Fernández-Luqueño et al., 2011). Fungi have capacity to adjust to changing situations And additionally have had under 100 years in which to advance within the sight of manufactured plastics and numerous different poisons. Capacities that have not been found at this point in growths identifying with the corruption of poisons could be advancing right now in

some exceptionally contaminated pocket of soil. Fungi are particularly appropriate to PAH degradation comparative with other bacterial decomposers for a couple of reasons. They can degrade high atomic weight PAHs. They likewise work well in non-watery situations where hydrophobic PAHs amass; Also, they can work in the extremely low-oxygen conditions that happen in intensely PAH polluted zones. a rundown of more than 51 contagious species or species bunches that are fruitful at corrupting diverse PAHs. A wide assortment of fungi have advanced compelling instruments to assault explicit PAHs.(Mai, Schormann, Majcherczyk, Hüttermann, & biotechnology, 2004)

The shellfish mushroom, *Pleurotus ostreatus*, can degrade 0-95% of all PAHs present in soil after 80 days(Steffen, Schubert, Tuomela, Hatakka, & Hofrichter, 2007). This is a wood-spoiling fungi part of a gathering known as white decay organisms. *Phanerochaete chrysosporium* and *laevis*. These basidiomycetes have at any rate two pathways. One pathway is the cytochrome 450 framework, much like the framework in vertebrate livers which separate huge particles into metabolites; in any case, a considerable lot of these metabolites are poisonous themselves. The lignin extracellular corruption pathway is ideal on the grounds that the metabolites are completely separated into carbon dioxide(Peng et al., 2008). Fungi assault plastic polymers too; these arrive in a wide scope of structures as lignin and

are followed up on by various organisms' species for various polymers. This disintegrating capacity is maybe much more noteworthy than PAH decomposition. It was discovered that the white decay basidiomycetes known for lignin corruption explicitly *P. ostreatus* could adequately separate polyacrylimide. Copolymerization with characteristic polymers, for example, starch or collagen expanded biodegradation. These give an extra carbon source to the growths and may likewise give passages to the organisms to attack the manufactured polymer strains of *Fusarium* and *Hypocrea* that could corrupt one cancer-causing high weight PAH, pyrene, just as take-up copper and zinc.(Mai et al., 2004). Fungi have advanced to decay lignin; lignin's variable and enormous structure is fundamentally the same as polycyclic sweet-smelling hydrocarbons, and the catalysts the fungi produce are nonspecific. Fungi must have the option to deal with the high internal heat level inside the body to get by in the cerebrum; a portion of these irresistible organisms have been found in saunas, hot tubs, natural aquifers, and coal squander heaps. These equivalent outrageous conditions, just as other extraordinary conditions, for example, low oxygen, are likewise present in PAH-degraded soils. Fungi have a bewildering potential to tidy up defiled conditions. In the wake of taking a gander at the rundown of organisms that can degrade various Plastics, one could envision that there is a growth out there to corrupt each kind of relentless poison, and everyone just must be found.(Gautam, Bassi, Yanful, & biotechnology, 2007)

No.	Type of Plastic Used	Fungi	References
1.	Degradable plastic	<i>Phanerochaete chrysosporium</i>	(B. Lee, Pometto, Fratzke, Bailey, & Microbiology, 1991)
2.	Polyurethane	<i>Chaetomium globosum</i> and <i>Aspergillus terreus</i>	(Boubendir, 1992)
3.	Polyurethane	<i>Curvularia senegalensis</i> , <i>Fusarium solani</i> , <i>Aureobasidium pullulans</i> ,	(Crabbe et al., 1994)
4.	Disposable plastic films	<i>Aspergillus flavus</i> and <i>Mucor rouxii</i>	(El-Shafei, Abd El-Nasser, Kansoh, Ali, & stability, 1998)
5.	HDPE	<i>Phanerochaete chrysosporium</i> <i>Trametes versicolor</i>	(Iiyoshi, Tsutsumi, & Nishida, 1998)
6.	PVC	<i>Poliporusversicolor</i> , <i>Pleurotus sajor caju</i> ,	(Kırbaş, Keskin, Güner, & toxicology, 1999)
7.	Plasticized PVC	<i>Aureobasidium pullulans</i> , <i>Rhodotorula aurantiaca</i> , and <i>Cluyveromyces spp.</i>	(Webb et al., 2000)
8.	Polyethylene	<i>Penicillium simplicissimum</i>	(Yamada-Onodera et al., 2001)
9.	Low density	<i>Penicillium pinophilium</i> and <i>Aspergillus niger</i>	(Volke-Sepúlveda, Saucedo-Castañeda, Gutiérrez-Rojas, Manzur, & Favela-Torres, 2002)
10.	Degradable polyethylene	<i>Cladosporium cladosporides</i>	(Bonhomme et al., 2003)
11.	Polyethylene bags	<i>Aspergillus niger</i>	(Kathiresan, 2003)
12.	LDPE	<i>Fusarium sp</i>	(Shah, Hasan, Hameed, & Akhter, 2009)
13.	High density polyethylene (HDPE)	<i>Aspergillus niger</i> , <i>Aspergillus oryzae</i> <i>Aspergillus flavus</i>	(Konduri, Koteswarareddy, Rohini Kumar, Venkata Reddy, & Lakshmi Narasu, 2011)
14.	LDPE powder	<i>Aspergillus nidulans</i> and <i>Aspergillus flavus</i>	(Usha, Sangeetha, & Palaniswamy, 2011)
15.	Polyethylene carries bags	<i>Aspergillus niger</i>	(Aswale & Ade, 2011)
16.	LDPE	<i>Aspergillus oryzae</i>	(Konduri et al., 2011)
17.	Degradable plastic	<i>Aspergillus niger</i> , <i>Aspergillus nidulans</i> , <i>Aspergillus flavus</i> , <i>Aspergillus glaucus</i>	(Priyanka & Archana, 2011)
18.	LDPE powder	<i>Aspergillus versicolor</i> and <i>Aspergillus sp</i>	(Pramila & Ramesh, 2011)
19.	LDPE	<i>Aspergillus sp</i>	(Raaman, Rajitha, Jayshree, & Jegadeesh, 2012)

The Role of Fungal Enzymes in Biodegradation of plastics

Enzymes exist in each living cell and henceforth in all organisms. Compounds are quite certain in their activity on substrates, Fungi produce extracellular catalysts to corrupt lignin, which can't go through the cell dividers of microorganisms. This cycle of corruption is called mineralization, and the finished result is carbon dioxide.(K.-M. Lee, Gimore, Huss, & Environment, 2005). Laccase can help in the oxidation of the hydrocarbon spine of PE. Gel pervasion chromatography (GPC) decide if without cell laccase hatched with PE helps in the decrease of normal sub-atomic weight and normal sub-atomic number of PE by 20% and 15%, respectively.(Sivan, 2011). Laccases are generally present in lignin biodegrading organisms, where they catalyze the oxidation of fragrant mixes.

Laccase movement is known to follow up on nonaromatic substrates.(Mayer & Staples, 2002).Lignin and manganese-subordinate peroxidases (LiP and MnP, separately) and laccases are the three fundamental proteins of ligninolytic system. (Hofrichter, Lundell, Hatakka, & Microbiology,2001)

A few strains that are equipped for debasing the PE are *Brevibacillus spp.*, *Bacillus spp.*, where proteases are answerable for corruption of plastics. (Sivan, 2011).Papain and urease are the two proteolytic chemicals found to debase clinical polyester PU. Polymer corrupted by papain was because of the hydrolysis of urethane and urea linkages creating free amine and hydroxyl groups. (Phua, Castillo, Anderson, & Hiltner, 1987). Fungi and manganese peroxidase, incompletely filtered from the strain of *Phanerochaete chrysosporium* additionally helps in the debasement of high-

atomic weight PE under nitrogen and carbon restricted conditions. (Shimao, 2001) The chemicals liable for the debasement of different sorts of plastics portray the substrates that use the plastics as carbon and fuel sources and

helps in biodegradation. Microbial compounds actuate the pace of biodegradation of plastics viably without making any mischief the environment. (Devi et al., 2016)

No.	Enzyme	Microorganism	Plastic	references
1.	Glycosidase	<i>A. flavus</i>	PCL	(Tokiwa, Calabria, Ugwu, & Aiba, 2009)
2.	Cutinase	<i>Aspergillus oryzae</i>	PBS	(Maeda et al., 2005)
3.	Catalase, protease	<i>A. niger</i>	PCL	(Tokiwa et al., 2009)
4.	Manganese	<i>phnerochaete</i>	PEL	(Shimao, 2001)
5.	Cutinase	<i>Fusarium</i>	PCL	(Shimao, 2001)
6.	Urease	<i>Trichoderma sp.</i>	polyurethane	(Loredo-Treviño et al., 2011)
7.	Serine hydrolase	<i>Pestalotiopsis microspore</i> <i>Curvularia senegalensis</i> <i>Fusarium solani</i>	PUR	(Russell et al., 2011)

Table 4: Various enzymes used to degrade plastics

Biodegradation of Synthetic Plastics by Fungi

The degradation of most manufactured plastics in nature is a moderate cycle that includes ecological variables, trailed by the activities of microorganisms.

The essential instrument for the biodegradation of high-sub-atomic weight polymer is the oxidation.(A. C. Albertsson, Barenstedt, & Karlsson, 1994; Cruz-Pinto, Carvalho, Ferreira, & Physics, 1994)

No.	Plastic	Applications
1.	PET	SOFT DRINK, WATER AND DRESSING bottles, peanut butter, and jam bars.
2.	HDPE	Milk, juice and water bottles, trash and retail bags.
3.	PVC	Juice bottles, cling films, raincoats, visors, shoes soles, garden hoses, and electricity pipes.
4.	LDPE	Frozen food bags, squeezable bottles, flexible container lids
5.	PP	Bottle caps, drinking straws, medicine bottles, car batteries, disposable syringes.
6.	PS	Packing materials, laboratory ware, disposable cups, plates, trays, and cutlery.
7.	OTHER OFTEN POLYCARBONATE	Beverage bottles, baby milk bottles, electronic casing

Table 5: plastic degradation and its applications.

Polyethylene

Polyethylene is one of the manufactured polymers of high hydrophobic level and high-atomic weight. In common structure, it isn't biodegradable. Accordingly, their utilization in the creation of removal or pressing materials causes risky ecological problems(Krupp, Jewell, & technology, 1992) To make this biodegradable, it requires the adjustment of its crystallinity sub-atomic weight and mechanical properties that are liable for PE obstruction toward degradation(A. C. Albertsson et al., 1994).Biodegradation of PE is known to happen by two components: hydro-biodegradation and Oxo-biodegradation (Bonhomme et al., 2003). These two instruments concur with the alterations because of the two added substances, starch and favourable to oxidant, utilized in the amalgamation of biodegradable PE. Biodegradation of LDPE film was 0.2% weight reduction in 10 years(A.-C. J. E. P. J. Albertsson, 1980).Yamada-Onodera et al. (2001) confined a strain of organism *Penicillium simplicissimum* YK to biodegrade PE, with no additives.(Yamada-Onodera et al., 2001) The capacity of parasites and *Streptomyces* strains to assault degradable PE comprising of arranged PE packs containing 6% starch. They secluded eight unique strains of *Streptomyces* and two parasites *Mucor rouxii* NRRL 1835 and *As. flavus*. (El-Shafei et al., 1998)

Polypropylene

Polypropylene is a thermoplastic polymer utilized in a wide assortment of utilizations including bundling and naming, materials (e.g., ropes, warm clothing, and covers), writing material, plastic parts, and reusable compartments of different sorts, research center gear, amplifiers, car segments, and polymer banknotes. Most business PP is isotactic and has a middle of the road level of crystallinity between that of LDPE and HDPE. PP is ordinarily extreme and adaptable, particularly when copolymerized with ethylene.(Maier & Calafut, 1998). Polypropylene is subject to fasten debasement from introduction to warmth and UV radiation, for example, that present in sunlight(Morris, 2005).Microbial people group disconnected from soil tests blended in with starch have been fit for corrupting PP (Cacciari et

al., 1993). Biodegradation of isotactic PP with no treatment is accounted for with one of the networks assigned as 3S among the four microbial networks (assigned as 1S, 2S, 3S, and 6S) adjusted to develop on starch containing PE got from advancement culture. *Pseudomonas chlororaphis*, *Pseudomonas stutzeri*, and *Vibrio* species were distinguished in the network 3S.(Alariqi, Kumar, Rao, Singh, & Stability, 2006)

Polyvinyl Chloride

Polyvinyl chloride is a solid plastic that opposes scraped spot and synthetic substances and has low dampness ingestion. PVC is utilized in development since it is more compelling than conventional materials, for example, copper, iron, or wood in line and profile applications. It very well may be made gentler and more adaptable by the expansion of plasticizers, Mostly, PVC is utilized in structures for lines and fittings, electrical wire protection, floor covers, and engineered cowhide items. It is additionally used to make shoe soles, inflexible lines, materials, and nursery hoses (Owen, 1984).As indicated by Kirbas et al. (1999), PVC having low-atomic weight can be presented to biodegradation by the utilization of white-decay fungi (Kirbaş et al., 1999). The organism *As. fumigatus* viably corrupts plasticized PVC. *Phanerochaete chrysosporium* was developed on PVC in a mineral salt agar. *Phanerochaete chrysosporium*, *Lentinus tigrinus*, *As. niger*, and *Aspergillus sydowii* can successfully debase PVC.(Ali, 2011)

Polystyrene

Polystyrene is an engineered polymer that contains a rehashing gathering, likewise viewed as exceptionally steady and less powerless for biodegradation. It is utilized in the creation of dispensable cups, bundling materials, in research facility product, and in certain electronic employments. PS is utilized for its lightweight, firmness, and great warm protection. At the point when it is corrupted by warm or synthetic methods it discharges items, for example, styrene, benzene, toluene, and acrolein (Tsuchii, Suzuki, Takahara, & Chemistry, 1977). Biodegradation analyses of PS, styrene oligomers, and PS copolymers have been had a go at utilizing bacteria

(Higashimura et al., 1983), fungi (Milstein et al., 1992) blended culture and catalyst under various conditions. Enzymatic biodegradation of PS polymer was attempted with hydro-quinone peroxidase protein with success (Nakamiya, Ooi, Kinoshita, & bioengineering, 1997). The protein was separated from lignin decolorizing bacterium *Azotobacter beijerinckii* HM121 and utilized in a two-stage (watery and dissolvable) system. (Devi et al., 2016)

Polyurethane

Polyurethane is usually used as a constituent material in numerous items including furniture, covering, development materials, strands, and paints. (Saunders & Frisch, 1964). Primarily, PUR is the buildup result of polyisocyanate and polyol (Howard, Norton, & Burks, 2012). four types of organisms *Curvularia senegalensis*, *Fusarium solani*, *Aureobasidium pullulans*, and *Cladosporium* sp. were gotten from soil and found to debase ester-based PU. (Shah, Hasan, Akhter, Hameed, & Ahmed, 2008). Pathirana and Seal (1985) detailed that some polyester-PUR debasing organisms produce extracellular esterases, proteases, or ureases within the sight of PUR. Esterase action has been resolved in the way of life supernatant of growth, for example, *C. Senegalensis*. (Pathirana & Seal, 1985)

Conclusion

It is obvious that without plastic we can't meet our day-to-day life needs, but in view of its detrimental effect it is required to develop competent process for its safe disposal and explore alternative material like starch based and blended plastic. This knowledge is useful because currently, the annual worldwide use of plastic materials is gradually increasing. In the natural environment, different kinds of microorganisms play an important role in various steps involved in the degradation of plastics. Various plastic-degrading methods are available but the cheapest, eco-friendly acceptable method is degradation using microbes. The microbe releases the extracellular enzymes to degrade the plastic but the detailed characterization of these enzymes is still needed to be carried out. Utilization of molecular techniques to detect specific groups of microorganisms involved in the degradation process will allow a better understanding of the organization of the microbial community involved in the attack of materials.

Future prospects

To tackle the issues identified with the removal of plastic waste created from different sources, the most imaginative and earth safe path is to utilize biodegradable plastics in specific applications like bundling, horticulture, and wellbeing industry. Bio-and fossil-based biodegradable polymers, whenever used, are effectively corrupted in the earth, cells, or under enhanced modern facilities. The interest for nature well-disposed polymers is expanding consistently in specific applications. Use of these materials ought to be engaged in future particularly for the assembling of bundling stuff, the food thing bundling, and expendable clinical things. It is additionally useful to utilize biodegradable plastics in the earth as farming movies, fishery material (fishing nets), bio-absorbable materials in therapeutics, careful systems, and sterile merchandise. Additionally, biodegradable plastics ought to be applied where dissemination into nature is unavoidable or when it is trying to isolate the trash. Then again, appropriate game plan of their waste and littering control is fundamental to exploit such polymers in the community. Next-age bio-based biodegradable plastics will focus on building a more reasonable society for explicit applications. Further, these plastics ought to be biodegraded and reused in a reasonable manner to make them re-utilize conceivable

References

1. Ahmed, T., Shahid, M., Azeem, F., Rasul, I., Shah, A. A., Noman, M., . . . Research, P. (2018). Biodegradation of plastics: current scenario and future prospects for environmental safety. 25(8), 7287-7298.
2. Alariqi, S. A., Kumar, A. P., Rao, B., Singh, R. J. P. D., & Stability. (2006). Biodegradation of γ -sterilised biomedical

- polyolefins under composting and fungal culture environments. 91(5), 1105-1116.
3. Albertsson, A.-C. J. E. P. J. (1980). The shape of the biodegradation curve for low and high density polyethenes in prolonged series of experiments. 16(7), 623-630.
4. Albertsson, A. C., Barenstedt, C., & Karlsson, S. J. A. p. (1994). Abiotic degradation products from enhanced environmentally degradable polyethylene. 45(2), 97-103.
5. Ali, M. (2011). *Microbial degradation of polyvinyl chloride plastics*. Ph. D. thesis, Quaid-i-Azam University,
6. Aswale, P., & Ade, A. J. S. A. i. B., Pune. (2011). Polythene degradation potential of *Aspergillus niger*.
7. Atagana, H. I. J. W. J. o. M., & Biotechnology. (2004). Biodegradation of phenol, o-cresol, m-cresol and p-cresol by indigenous soil fungi in soil contaminated with creosote. 20(8), 851-858.
8. Bonhomme, S., Cuer, A., Delort, A., Lemaire, J., Sancelme, M., Scott, G. J. P. d., & Stability. (2003). Environmental biodegradation of polyethylene. 81(3), 441-452.
9. Boubendir, A. (1992). *Purification and biochemical evaluation of polyurethane degrading enzymes of fungal origin*. University of Salford,
10. Cacciari, I., Quatrini, P., Zirletta, G., Mincione, E., Vinciguerra, V., Lupattelli, P., . . . microbiology, e. (1993). Isotactic polypropylene biodegradation by a microbial community: physicochemical characterization of metabolites produced. 59(11), 3695-3700.
11. Crabbe, J. R., Campbell, J. R., Thompson, L., Walz, S. L., Schultz, W. W. J. I. B., & Biodegradation. (1994). Biodegradation of a colloidal ester-based polyurethane by soil fungi. 33(2), 103-113.
12. Cruz-Pinto, J., Carvalho, M., Ferreira, J. J. D. A. M. C. A. M. C., & Physics. (1994). The kinetics and mechanism of polyethylene photo-oxidation. 216(1), 113-133.
13. Devi, R. S., Kannan, V. R., Natarajan, K., Nivas, D., Kannan, K., Chandru, S., & Antony, A. R. J. E. W. M. (2016). The role of microbes in plastic degradation. 341.
14. El-Shafei, H. A., Abd El-Nasser, N. H., Kansoh, A. L., Ali, A. M. J. P. d., & stability. (1998). Biodegradation of disposable polyethylene by fungi and *Streptomyces* species. 62(2), 361-365.
15. Fernández-Luqueño, F., Valenzuela-Encinas, C., Marsch, R., Martínez-Suárez, C., Vázquez-Núñez, E., Dendooven, L. J. E. S., & Research, P. (2011). Microbial communities to mitigate contamination of PAHs in soil—possibilities and challenges: a review. 18(1), 12-30.
16. Gautam, R., Bassi, A., Yanful, E. J. A. b., & biotechnology. (2007). A review of biodegradation of synthetic plastic and foams. 141(1), 85-108.
17. Higashimura, T., Sawamoto, M., Hiza, T., Karaiwa, M., Tsuchii, A., Suzuki, T. J. A., & Microbiology, E. (1983). Effect of methyl substitution on microbial degradation of linear styrene dimers by two soil bacteria. 46(2), 386-391.
18. Hofrichter, M., Lundell, T., Hatakka, A. J. A., & Microbiology, E. (2001). Conversion of milled pine wood by manganese peroxidase from *Phlebia radiata*. 67(10), 4588-4593.
19. Howard, G. T., Norton, W. N., & Burks, T. J. B. (2012). Growth of *Acinetobacter gernerii* P7 on polyurethane and the purification and characterization of a polyurethanase enzyme. 23(4), 561-573.
20. Iiyoshi, Y., Tsutsumi, Y., & Nishida, T. J. J. o. w. s. (1998). Polyethylene degradation by lignin-degrading fungi and manganese peroxidase. 44(3), 222-229.
21. Kale, S. K., Deshmukh, A. G., Dudhare, M. S., & Patil, V. B. J. J. o. B. T. (2015). Microbial degradation of plastic: a review. 6(2), 952-961.

22. Kaseem, M., Hamad, K., & Deri, F. J. P. S. S. A. (2012). Thermoplastic starch blends: A review of recent works. *54*(2), 165-176.
23. Kathiresan, K. J. R. d. b. t. (2003). Polythene and plastics-degrading microbes from the mangrove soil. *51*(3-4), 629-633.
24. Kırbaş, Z., Keskin, N., Güner, A. J. B. o. e. c., & toxicology. (1999). Biodegradation of polyvinylchloride (PVC) by white rot fungi. *63*(3), 335-342.
25. Konduri, M. K., Koteswarareddy, G., Rohini Kumar, D., Venkata Reddy, B., & Lakshmi Narasu, M. J. J. o. A. P. S. (2011). Effect of pro-oxidants on biodegradation of polyethylene (LDPE) by indigenous fungal isolate, *Aspergillus oryzae*. *120*(6), 3536-3545.
26. Krupp, L. R., Jewell, W. J. J. E. s., & technology. (1992). Biodegradability of modified plastic films in controlled biological environments. *26*(1), 193-198.
27. Kyrikou, I., Briassoulis, D. J. J. o. P., & Environment, t. (2007). Biodegradation of agricultural plastic films: a critical review. *15*(2), 125-150.
28. Lee, B., Pometto, A. L., Fratzke, A., Bailey, T. B. J. A., & Microbiology, E. (1991). Biodegradation of degradable plastic polyethylene by *Phanerochaete* and *Streptomyces* species. *57*(3), 678-685.
29. Lee, K.-M., Gimore, D., Huss, M. J. J. o. P., & Environment, t. (2005). Fungal Degradation of the Bioplastic PHB (Poly-3-hydroxy-butyric acid). *13*(3), 213-219.
30. Lee, S. Y. J. B., & bioengineering. (1996). Bacterial polyhydroxyalkanoates. *49*(1), 1-14.
31. Loredó-Treviño, A., García, G., Velasco-Téllez, A., Rodríguez-Herrera, R., Aguilar, C. N. J. A. i. B., & Biotechnology. (2011). Polyurethane foam as substrate for fungal strains. *2*(2), 52.
32. Luzier, W. D. J. P. o. t. N. A. o. S. (1992). Materials derived from biomass/biodegradable materials. *89*(3), 839-842.
33. Maeda, H., Yamagata, Y., Abe, K., Hasegawa, F., Machida, M., Ishioka, R., . . . biotechnology. (2005). Purification and characterization of a biodegradable plastic-degrading enzyme from *Aspergillus oryzae*. *67*(6), 778-788.
34. Mai, C., Schormann, W., Majcherczyk, A., Hüttermann, A. J. A. m., & biotechnology. (2004). Degradation of acrylic copolymers by white-rot fungi. *65*(4), 479-487.
35. Maier, C., & Calafut, T. (1998). *Polypropylene: the definitive user's guide and databook*: William Andrew.
36. Mayer, A. M., & Staples, R. C. J. P. (2002). Laccase: new functions for an old enzyme. *60*(6), 551-565.
37. Milstein, O., Gersonde, R., Hüttermann, A., Chen, M.-J., Meister, J. J. J. A., & Microbiology, E. (1992). Fungal biodegradation of lignopolystyrene graft copolymers. *58*(10), 3225-3232.
38. Morris, P. J. (2005). *Polymer pioneers: a popular history of the science and technology of large molecules*: Chemical Heritage Foundation.
39. Nakamiya, K., Ooi, T., Kinoshita, S. J. J. o. f., & bioengineering. (1997). Non-heme hydroquinone peroxidase from *Azotobacter beijerinckii* HM121. *84*(1), 14-21.
40. Norton, J. M. J. J. o. H. i. a. H. H. i. C. C. H. (2012). Fungi for bioremediation of hydrocarbon pollutants. *10*, 18-21.
41. Owen, E. J. E. A. S. P. L., Ripple Road, Barking, Essex, UK, . (1984). Degradation and Stabilization of PVC.(Retroactive Coverage).
42. Pathirana, R., & Seal, K. J. I. B. B. (1985). Studies on polyurethane deteriorating fungi. *21*, 41-49.
43. Peng, R.-H., Xiong, A.-S., Xue, Y., Fu, X.-Y., Gao, F., Zhao, W., . . . Yao, Q.-H. J. F. m. r. (2008). Microbial biodegradation of polyaromatic hydrocarbons. *32*(6), 927-955.
44. Phua, S., Castillo, E., Anderson, J., & Hiltner, A. J. J. o. b. m. r. (1987). Biodegradation of a polyurethane in vitro. *21*(2), 231-246.
45. Pospíšil, J., & Nešpůrek, S. (1997). *Highlights in chemistry and physics of polymer stabilization*. Paper presented at the Macromolecular Symposia.
46. Pramila, R., & Ramesh, K. V. J. A. J. o. M. R. (2011). Biodegradation of low density polyethylene (LDPE) by fungi isolated from marine water a SEM analysis. *5*(28), 5013-5018.
47. Priyanka, N., & Archana, T. J. J. E. A. T. (2011). Biodegradability of polythene and plastic by the help of microorganism: a way for brighter future. *1*(4), 1000111.
48. Raaman, N., Rajitha, N., Jayshree, A., & Jegadeesh, R. J. J. A. I. R. (2012). Biodegradation of plastic by *Aspergillus* spp. isolated from polythene polluted sites around Chennai. *1*(6), 313-316.
49. Russell, J. R., Huang, J., Anand, P., Kucera, K., Sandoval, A. G., Dantzler, K. W., . . . microbiology, e. (2011). Biodegradation of polyester polyurethane by endophytic fungi. *77*(17), 6076-6084.
50. Saunders, J. H., & Frisch, K. C. (1964). *Polyurethanes: Chemistry and Technology: Chemistry* (Vol. 16): Interscience Publishers.
51. Shah, A. A., Hasan, F., Akhter, J. I., Hameed, A., & Ahmed, S. J. A. o. m. (2008). Degradation of polyurethane by novel bacterial consortium isolated from soil. *58*(3), 381.
52. Shah, A. A., Hasan, F., Hameed, A., Ahmed, S. J. I. b., & biodegradation. (2007). Isolation and characterization of poly (3-hydroxybutyrate-co-3-hydroxyvalerate) degrading bacteria and purification of PHBV depolymerase from newly isolated *Bacillus* sp. AF3. *60*(2), 109-115.
53. Shah, A. A., Hasan, F., Hameed, A., & Akhter, J. I. J. A. J. o. M. R. (2009). Isolation of *Fusarium* sp. AF4 from sewage sludge, with the ability to adhere the surface of polyethylene. *3*(10), 658-663.
54. Shimao, M. J. C. o. i. b. (2001). Biodegradation of plastics. *12*(3), 242-247.
55. Sivan, A. J. C. o. i. b. (2011). New perspectives in plastic biodegradation. *22*(3), 422-426.
56. Steffen, K. T., Schubert, S., Tuomela, M., Hatakka, A., & Hofrichter, M. J. B. (2007). Enhancement of bioconversion of high-molecular mass polycyclic aromatic hydrocarbons in contaminated non-sterile soil by litter-decomposing fungi. *18*(3), 359-369.
57. Takaku, H., Kimoto, A., Kodaira, S., Nashimoto, M., & Takagi, M. J. F. m. l. (2006). Isolation of a Gram-positive poly (3-hydroxybutyrate)(PHB)-degrading bacterium from compost, and cloning and characterization of a gene encoding PHB depolymerase of *Bacillus megaterium* N-18-25-9. *264*(2), 152-159.
58. Tokiwa, Y., Calabia, B. P., Ugwu, C. U., & Aiba, S. J. I. j. o. m. s. (2009). Biodegradability of plastics. *10*(9), 3722-3742.
59. Tsuchii, A., Suzuki, T., Takahara, Y. J. A., & Chemistry, B. (1977). Microbial degradation of styrene oligomer. *41*(12), 2417-2421.
60. Usha, R., Sangeetha, T., & Palaniswamy, M. J. L. A. R. C. J. I. (2011). Screening of polyethylene degrading microorganisms from garbage soil. *2*(4), 200-204.
61. Vaverková, M., Adamcová, D., Kotovicová, J., Toman, F. J. E. C., & S, E. (2014). Evaluation of biodegradability of plastics bags in composting conditions. *21*(1), 45-57.
62. Volke-Sepúlveda, T., Saucedo-Castañeda, G., Gutiérrez-Rojas, M., Manzur, A., & Favela-Torres, E. J. J. o. a. p. s. (2002). Thermally treated low density polyethylene biodegradation by *Penicillium pinophilum* and *Aspergillus niger*. *83*(2), 305-314.
63. Wackett, L. P., & Hersherberger, C. D. (2001). *Biocatalysis and biodegradation: microbial transformation of organic compounds*: ASM press Washington, DC.

64. Webb, J. S., Nixon, M., Eastwood, I. M., Greenhalgh, M., Robson, G. D., Handley, P. S. J. A., & microbiology, e. (2000). Fungal colonization and biodeterioration of plasticized polyvinyl chloride. 66(8), 3194-3200.
65. Williams, S., & Peoples, O. J. C. (1996). Biodegradable plastics from plants. 26(9), 38-44.
66. Yamada-Onodera, K., Mukumoto, H., Katsuyaya, Y., Saiganji, A., Tani, Y. J. P. d., & stability. (2001). Degradation of polyethylene by a fungus, *Penicillium simplicissimum* YK. 72(2), 323-327.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

Submit Manuscript

DOI: [10.31579/2766-2314/149](https://doi.org/10.31579/2766-2314/149)

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://www.auctoresonline.org/journals/biotechnology-and-bioprocessing>