

More Information about Ascorbic Acid in Human Food

Fahim Aziz Eldin Shaltout

Food Control Department, Faculty of Veterinary Medicine Benha University, Egypt.

***Corresponding Author:** Fahim Aziz Eldin Shaltout, Food Control Department, Faculty of Veterinary Medicine, Benha University, Egypt.

Received date: June 23, 2025; **Accepted date:** July 07, 2025; **Published date:** July 18, 2025

Citation: Fahim Aziz Eldin Shaltout, (2025), More Information about Ascorbic Acid in Human Food, *J Thoracic Disease and Cardiothoracic Surgery*, 6(4); DOI:10.31579/2693-2156/139

Copyright: © 2025, Fahim Aziz Eldin Shaltout. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Ascorbic acid (C₆H₈O₆) is an organic compound belonging to the family of monosaccharide. It is highly soluble in water, and is often called one of the secrets of the Mediterranean diet. Its use is widespread in the food industry is also important, having always been exploited for its antioxidant and stabilising ability. Many indeed are the additive formulations that take advantage of these properties.

Keywords: ascorbic acid; food industry; additive formulations; antioxidant

Introduction

The ascorbic acid (C₆H₈O₆) – better known as vitamin C – is an organic compound belonging to the family of monosaccharide. It is strongly water-soluble and it is often considered as one of the elements that characterises the Mediterranean diet. The diffusion of its use is also most important in the food industry, which has always used its stabilised and antioxidant property. Indeed, there are several formulations of additives that contain ascorbic acid.

One of the most important characteristics of the ascorbic acid is its reducing ability. In the presence of oxygen, ascorbic acid tends to oxidise with a strong result, especially in relation to catalyst metals, removing the environmental resources of oxygen. Furthermore, the ascorbic acid can react with free radicals, arresting the chain reactions that may provoke dangerous effects on organisms, such as neoplastic pathologies of the oral cavity, alimentary system, etc (1-7). It allows maintaining stable other important elements, such as vitamin A, E, folic acid and thiamine in organisms and foods. and it is essential for synthesis of collagen, connective tissue's protein, important to heal wounds, sores and lesions, and to the prevention of hemorrhages. In addition to prevent the onset of the atherosclerosis, vitamin C contributes to the development of the adrenaline and the endogenous serotonin and to the hydroxylation of aromatic compounds in the liver. It operates in processes of the cellular defense, facilitates the intestinal absorption of the iron and the elimination of toxic heavy metals (such as cadmium, nickel and lead) with which it is able to tie. Its action is also important in the synthesis of the carnitine in the zymotic transformation of the cholesterol in bile acid or in vitamin D (8-15). The use of the ascorbic acid is important, especially in production activities of the ground meat and cold cuts. In meat and ground meat the ascorbic acid prevents the oxidation and the discoloration of the product during the storage. This effect delays the appearance of an abnormal coloration that could be unpleasant to the consumer but it is not associated

to the organoleptic alteration of the product. The use of the ascorbic acid in meats, with the addition of nitrites, is important for the activity of reduction dependent upon nitrousmetamyoglobin-Fe (III) converted into nitrousmetamyoglobin-Fe (II), which maintains the colour of the product most brilliant. Furthermore, the ascorbic acid prevents the development of nitrosamines (16-22). The unauthorised use of the ascorbic acid as additive in the preparation of ground meat is often object of control by the competent authorities, with official samples. Whether the result of the laboratory analysis, made at the Institutes for Experimental Veterinary Medicine will be positive, the food business operator (FBO) will be indicted for the misdemeanor. In the preparation of food and drink is banned using, selling, possessing to sell the ascorbic acid or administers it like compensation of employees or distributes it to the consumption of food substances with the addition of not authorized chemical additives, under the decree of the Minister of Health or, whether they would be authorized, without the compliance of required rule of law for their use (23-29).

Discussion

Ascorbate is one of the most important vitamins in the human diet, being obtained largely from vegetables, fruit, and other plant material. Green leaves may contain as much ascorbate as chlorophyll. The presence of such a remarkable amount of ascorbate may suggest considerable metabolic significance, particularly if the amount is related to biological importance. Glutathione peroxidase, which is important in H₂O₂ detoxification in animals is largely absent from plant tissues. Seasonal variations in the ascorbate content of leaves have been found; on the whole, the ascorbate pool in chloroplasts and leaves is maintained at a remarkably constant level. Oxidation of ascorbic acid occurs in two sequential steps, forming, in the first instance, monodehydroascorbate, and subsequently dehydroascorbate. The monodehydroascorbate radical

spontaneously disproportionates to ascorbate and dehydroascorbate. The conversion of glucosone and sorbosone to ascorbic acid has been observed in bean and spinach leaves (30-36).

Ascorbic Acid Sources

Ascorbic acid is synthesized from glucose and is soluble in water. Most animals cannot synthesize ascorbic acid endogenously, but poultry, animals can synthesize it due to the L-gulonolactone oxidase enzyme, which is available in the animals' body. This enzyme is available in the kidney tissue where L-gulono-g-lactone is converted into ascorbic acid by the L-gulonolactone oxidase enzyme. In birds, ascorbic acid is produced in the kidney, while in mammals, it is produced in the liver. Ascorbic acid can be found in natural plants or synthesized through industrial processes. For plants, ascorbic acid is found from rose hips and sea buckthorn as the richest resource. Furthermore, other fruits such as guava, star fruit, black currant, strawberry, and kiwi also contained a high amount of ascorbic acid. As we know, citrus fruits such as orange, lime, grapefruit, and so on contain ascorbic acid. Thus, the ascorbic acid content of the latter is much less than the former. The amount of ascorbic acid per fruit is shown in. This table shows that guava, cashew apple, and sea buckthorn contain high ascorbic acid levels (37-43). An ordinary diet of natural and synthetic ascorbic acid is the only way to maintain the physiological requirements. The well-known symptom of ascorbic acid deficiency is associated with connective tissue damage, such as scurvy, which is characterized by fragile tissues and poor wound healing. The currently recommended dietary allowances (RDA) for ascorbic acid are 90 mg/day and 75 mg/day for men and women, respectively. Researchers have found that the steady-state concentration of ascorbic acid in plasma is about 80 $\mu\text{mol/L}$, when sufficient fruits and vegetables are consumed every day. Oral dosing of ascorbic acid (1.25 g) can improve the concentration of ascorbic acid in plasma to $134.8 \pm 20.6 \mu\text{mol/L}$. In order to maintain the ascorbic acid concentration required by the body, ascorbic acid-fortified dietary supplements or foods have attracted interest from consumers, in addition to fruits and vegetables found in nature (44-50). The main challenge in the development of ascorbic acid products is its high instability and reactivity. Ascorbic acid is reversibly oxidized into dehydroascorbic acid (DHA) upon exposure to light, heat, transition metal ions and pH (alkaline condition), then DHA further irreversibly hydrolyzes to form 2,3-diketogulonic acid. In recent decades, the strategy of shielding ascorbic acid from sensitive environments by encapsulating ascorbic acid within a layer of wall material has attracted much interest among researchers. A series of innovation delivery technologies have emerged, including microfluidic, melt extrusion, spray drying and chilling. The particles prepared by these methods are usually on the microscale. Under certain conditions, nano-encapsulation of ascorbic acid can be realized through ion gelation of chitosan or complex coacervation with anionic polymers. On the other hand, some bioactive compounds with low molecular weight can protect ascorbic acid by scavenging the pro-degradation factors of ascorbic acid in the solution (51-57).

Bioactivity of Ascorbic Acid

Antioxidant

The by-products of normal cell metabolism are reactive oxygen species (ROS), including superoxide radicals ($\text{O}_2^{\cdot-}$), singlet oxygen ($^1\text{O}_2$), hydrogen peroxide (H_2O_2) and highly reactive hydroxyl radicals (OH^{\cdot}). The adverse effect of ROS is that it can initiate a cascade of radicals, producing hydroxyl free radicals and other destructive species. These further induce protein and DNA damage, lipid peroxidation and finally lead to cell apoptosis. The antioxidant defense system cannot fully eliminate the toxic ROS accumulated in the cells, that is, the so-called "oxidative stress" occurs. In addition to enzymatic reactions, ROS can also be eliminated through non-enzymatic means such as antioxidants. Ascorbic acid is a free radical and other oxygen species scavenger, which can protect cells from oxidative damage caused by ROS. Antiradical capability commonly reflects the antioxidant ability, and ascorbic acid in foodstuff and bio-systems acts as antioxidant. As the most effective and

natural antioxidant with the least side effects, ascorbic acid can inhibit various diseases caused by oxidative stress in the body, such as cancer, cardiovascular disease, aging and cataracts (58-64). Studies have shown that the mortality from these diseases is inversely related to plasma concentration of ascorbic acid. Ascorbic acid and its derivatives can reduce the level of lipid peroxidation in vivo due to aging. In the absence of transition metals, ascorbic acid can reduce the frequency of mutations induced by H_2O_2 in human cells. Compared with other polyphenols or flavonoid antioxidants, ascorbic acid terminates the free radical chain reaction through disproportionation reaction, and the reaction products such as DHA and 2,3-diketogulonic acid are non-damaging and non-radical products. Another manifestation of antioxidant property is that ascorbic acid can form relatively stable ascorbic acid free radicals to donate single electrons. As reported, antioxidants can also repair tryptophan free radicals produced by the one-electron oxidation of free tryptophan in lysozyme to maintain protein integrity (65-71). Ascorbic acid is also used as an antioxidant to protect the sensory and nutritional properties of foods. As an anti-browning agent, it can inhibit the browning of vegetables and fruits caused by oxidation. The formation of quinones mediated by polyphenol oxidase causes the accumulation of H_2O_2 , which in turn causes the browning of polyphenols mediated by peroxidase. Ascorbic acid inhibits browning by reducing the o-quinone produced by polyphenol oxidase to the original diphenol through a process called "deactivation reaction". In addition to the regeneration mechanism of polyphenols, the protective effect is also attributed to the competitive inhibition of polyphenol oxidase activity by ascorbic acid. Meanwhile, addition of ascorbic acid causes a decrease in pH and is not conducive to the expression of polyphenol oxidase activity (72-78). In meat products, ascorbic acid is widely used as a natural agent for color retention, which can inhibit lipid oxidation and maintain color stability. Compared with other organic acids such as malic acid, citric acid and tartaric acid, ascorbic acid had the best protective effect on the quality of cured meat and was a suitable ingredient for cured meat products. The surface of the pork sprayed with ascorbic acid and a mix of that and rosemary extract maintained good stability in color, water content and pH after frozen storage. It is worth noting that this dietary source of ascorbic acid added in meat products is often overlooked. Norwegian researchers found that the content of ascorbic acid in sausages is 11-40 mg/mL, but ascorbic acid is usually ignored in the table of food ingredients because the added ascorbic acid is used as a color retention agent rather than a nutrient component. As a result, the actual ascorbic acid intake of Norwegian residents increased by 3-10%. The ascorbic acid added to the edible polysaccharide film can eliminate or quench the free radicals generated by radiation (79-85). As a radiation inhibitor, ascorbic acid can inhibit the decrease in the viscosity of carrageenan caused by radiation and protect its rheological properties. Ascorbic acid can inhibit food-borne pathogens in the early stage of biofilm formation due to its anti-quorum sensing activity and inhibition of extracellular polymer production. The efficacy of ascorbic acid is related to its concentration and the strain. For *Escherichia coli* and *Staphylococcus aureus*, the inhibitory effect of ascorbic acid at 25 mg/mL is the greatest, and lower concentrations of ascorbic acid are ineffective. For *Listeria monocytogenes*, ascorbic acid at 0.25 mg/mL shows an inhibitory effect (86-92).

Pro-Oxidant

Pro-oxidant activity is defined as the ability of antioxidants to reduce transition metal ions to a lower oxidation state, which refers to the Fenton reaction. In the Fenton reaction, transition metal ions such as Fe^{3+} are reduced by ascorbic acid and then Fe^{2+} further react with oxygen and hydrogen peroxide to form highly active and destructive hydroxyl radicals. Ascorbic acid does not always express antioxidant activity, and may be converted into a pro-oxidant and show toxic effects under certain conditions. The effect of ascorbic acid on the redox properties of bovine hemoglobin is dual with an antioxidant effect at the initial stage of the reaction. During the reaction process, ascorbic acid generates hydrogen peroxide under the mediation of oxygen or oxygenated hemoglobin (93-99). With the consumption of ascorbic acid, its own scavenging ability

cannot balance the accumulated hydrogen peroxide, which leads to the formation of bilirubin and accelerates the oxidation of hemoglobin. Studies have shown that ascorbic acid will transform from an antioxidant under physiological conditions to a pro-oxidant at higher concentrations. Researchers found that supplementation of 500 mg of ascorbic acid in the diet for 6 weeks increases the level of oxidative damage to peripheral blood lymphocytes, although this result is still controversial in the academic community (100-106). Furthermore, the presence of transition metal ions in the system is also a key factor for ascorbic acid exerting pro-oxidant activity. In mayonnaise, the added ascorbic acid works as a lipid antioxidant or pro-oxidant depending on the presence or absence of the fat-soluble antioxidant vitamin E. In the system containing vitamin E, the synergistically antioxidant effect of these two vitamins is stronger than the pro-oxidant effect of ascorbic acid. Without the addition of vitamin E, the hydrogen peroxide at the interface of the oil droplets promotes the lipid oxidation of lipoprotein particles in the mayonnaise, which in turn induces the oxidation of apolipoproteins and produces volatile odors (107-113). In addition, dehydroascorbic acid may irreversibly degrade and produce highly reactive carbonyl intermediates, which can induce glycosylation of proteins. This is a non-enzymatic, non-specific reaction between carbonyl and amino groups, which is involved in a variety of age-related diseases. It is worth noting that the pro-oxidation of ascorbic acid can induce the apoptosis of cancer cells, thereby exerting anti-cancer effects to a certain extent. As reported, the copper-dependent cellular redox state is an important factor in the cytotoxic effect of ascorbic acid on cancer cells. Ascorbic acid mobilizes nuclear copper to cause pro-oxidative cleavage of cellular DNA, and nuclear copper serves as a new molecular target for the toxic effects of cancer cells. From this perspective, the pro-oxidation effect of ascorbic acid is beneficial (114-120). Up to now, the mechanism and conditions that induce ascorbic acid to express pro-oxidant properties have not been clearly elucidated, and the definition of the conversion concentration between antioxidant and pro-oxidant is also unclear. Moreover, most of the reports on the promotion of oxidation of ascorbic acid are concentrated in vitro (121-127).

Co-Factors

Ascorbic acid can also be used as a co-factor for enzymes and other bioactive components to indirectly exert biological activities by acting as a free radical scavenger and electron transfer donor/acceptor to directly express its antioxidant properties. In the metabolic process of animals and plants, ascorbic acid does not directly participate in the catalytic cycle. As an enzyme co-factor, ascorbic acid exerts its indispensable function by regulating hydroxylation processes in multiple enzymatic reactions (128-134). For the active part of the enzyme with iron or copper, the role of ascorbic acid is to maintain the transition metal ions of these enzymes in a reduced form to exert their maximum physiological activity. Ascorbic acid is a co-factor for non-heme iron α -ketoglutarate-dependent dioxygenases such as prolyl 4-hydroxylase with the role in the synthesis of collagen. As an electron donor, ascorbic acid can keep iron in the ferrous state, thereby maintaining the full activity of collagen hydroxylase (135-141). This promotes the hydroxylation of proline and lysine residues, allowing pro-collagen correct intracellular folding. Ascorbic acid can also promote catecholamine synthesis by circulating tetrahydrobiopterin and enhance adrenal steroid production by increasing the expression of tyrosine hydroxylase. As a co-factor, it helps dopamine β -hydroxylase convert dopamine to norepinephrine. In addition, ascorbic acid can regulate cardiomyopathy and neurometabolic diseases. For example, as a co-factor for carnitine synthesis, it can shuttle fatty acids into the mitochondria and reduce oxidative stress. In some clinical situations, as a co-factor for the biosynthesis of amidated opioid peptides, taking ascorbic acid can exhibit analgesic effects (142-148).

Synergistic Effect

As a natural antioxidant, ascorbic acid mostly exists in the form of coexistence with other components in nature. Combining it with other

antioxidants may produce additive or even synergistic effects. Ascorbic acid and vitamin E, as chain-scission antioxidants, have an important inhibitory effect on the auto-oxidation of cell membrane polyunsaturated liposomes in vivo and the oxidation of lipids in vitro. Studies have shown that the combination of 15% ascorbic acid and 1% α -tocopherol can significantly inhibit erythema and the formation of sunburn cells. The synergy between α -tocopherol and ascorbic acid relies on the ability of ascorbic acid to regenerate α -tocopherol, and maintain the antioxidant capacity of α -tocopherol through circulation and inhibition of pro-oxidation. The combined use of ascorbic acid and gallic acid is a promising strategy to prevent the formation of advanced glycation end products, showing the synergistic effect in the inhibition of amyloid cross- β -structure and protein carbonyl formation in fructose-induced BSA glycosylation samples. Lycopene can inhibit inflammation and further stimulate the release of anti-inflammatory cytokine IL-10 when it combined with ascorbic acid and/or α -tocopherol. Understanding the synergy between ascorbic acid and other bioactive compounds allows the antioxidant system of foods and drugs to be selected more specifically (149-155).

Sensitivity to Environment:

Concentration and pH:

Ascorbic acid is unstable in aqueous solutions, and its degradation has been considered the main cause of quality and color changes during food storage and processing. Stability analysis of ascorbic acid is a key point in its application. In addition to the interference of external factors, the concentration of ascorbic acid in the solution will also affect its stability. As reported, after storage at room temperature in the presence of light for 27 days, an aqueous solution of 1% concentration of ascorbic acid lost around 21% of its initial concentration, while the 10% ascorbic acid system only degraded about 8%. In order to ensure sufficient content of ascorbic acid in the product, reinforcing the ascorbic acid content is a method commonly used in the food industry. The content of ascorbic acid in fortified milk decreased from 36.4 mg/L to 26.1 mg/L after sterilization, while the ascorbic acid content of normal milk dropped from 12.2 mg/L to 8.3 mg/L. Although the loss content of ascorbic acid increased with the addition of ascorbic acid, the loss efficiency decreased compared with the initial concentration of ascorbic acid. According to the stability and the degradation kinetics of ascorbic acid, a higher concentration of ascorbic acid has a lower degradation rate constant. However, some studies have shown that excessive ascorbic acid (AH2) is prone to auto-oxidation to produce dehydroascorbic acid anions (156-162).

Temperature

Ascorbic acid is severely degraded by heat, and the instability of ascorbic acid in thermal-processed foods impedes its application. The degradation of ascorbic acid involves complex oxidation and intermolecular rearrangement reactions, and is considered to be one of the main reasons for quality and color changes during food processing and storage. From an analysis of the effects of temperature and pressure on the retention of ascorbic acid in processed juices, it revealed that the dominant factor determining the stability of ascorbic acid is the temperature, which directly affects the degradation rate of ascorbic acid. Products containing ascorbic acid, such as fruit juice, need to undergo high-temperature pasteurization in order to guarantee safety and stability. The content of ascorbic acid in fresh orange juice ranges from 25–68 mg/100 mL. Studies have shown that the retention of ascorbic acid in the product after pasteurization (90 °C, 1 min) is about 82–92%. It was observed that the maximum temperature of the ultra-high pressure homogenization treatment at 100, 200 and 300 MPa was 45 °C, 70 °C and 94 °C, respectively, and the continuous treatment time under the maximum temperature was 0.7 s or less. The loss of ascorbic acid in the ultra-high pressure treated juice is less than that in the traditionally heat-pasteurized juice. As detected, the content of ascorbic acid in guava juice was around 42.2 ± 0.01 mg/mL. After 7 days of storage at 25 and 35 °C in the dark,

ascorbic acid was degraded by 23.4% and 56.4%, respectively. Its degradation is significantly reduced at 4–10 °C. The use of relatively mild temperature (75 °C) for heat treatment and a storage temperature below 25 °C is optimal for maintaining the ascorbic acid content of the product. The degradation of ascorbic acid during storage and heat treatment follows first-order kinetics based on a classic dynamic model (163-169).

The degradation or oxidation products of ascorbic acid heated at 100 °C for 2 h include furfural, 2-furoic acid, 3-hydroxy-2-pyrone and an unknown compound. Among them, furfural is one of the main degradation products of ascorbic acid, which can polymerize or combine with amino acids to form brown melanoids, causing the browning of ascorbic acid-containing juice products. Furthermore, thermally oxidized ascorbic acid was identified as a potential precursor of furan; it is a possible carcinogen usually produced in some heated food products. Meanwhile, natural and synthetic antioxidants, such as chlorogenic acid, have a certain mitigation effect on the formation of furan induced by heated ascorbic acid, but the mitigation effect may decrease with the increase in heating time. In addition, the thermal degradation process of ascorbic acid is also affected by pH, oxygen concentration, transition metal ions and oxidases, which is a complex system. Some researchers in food science believe that oxygen saturation decreases with increasing temperature and drops to 0 at 100 °C. However, according to the Tromans and Battino model, although 100–130 °C is the minimum oxygen solubility temperature, there is still dissolved oxygen in the system. At temperatures above 100 °C, oxygen has a greater effect on ascorbic acid degradation than temperature. Therefore, removing all oxygen including dissolved oxygen is the best way to preserve ascorbic acid at high temperatures (170-176).

Light

In addition to heat-treatment sterilization, ultraviolet radiation (240 nm–300 nm) is a promising alternative and is gradually being used more for fruit juice sterilization. The ultraviolet sterilization method includes the use of high-intensity pulsed ultraviolet radiation with wavelengths between 200 and 400 nm and a monochromatic ultraviolet system of which approximately 90% of the energy comes from a single wavelength. However, ascorbic acid absorbs ultraviolet radiation in the wavelength range of 229–330 nm and undergoes degradation. The formation of UV-induced free radicals may accelerate the loss of ascorbic acid. Ascorbic acid continues to degrade after UV treatment; higher initial UV dose values and storage temperature accelerate the degradation of ascorbic acid in the later stage. In addition, the pH of the solution also affects the photo-degradation of ascorbic acid. Under alkaline conditions, AH⁻ produced by ionization of AH₂ is more prone to photo-degradation than AH₂. It is worth noting that the ingredients in products may absorb or scatter UV radiation, thereby affecting the degradation of ascorbic acid. Niacinamide, as a component of vitamin B-complex with vitamin C, acts as a photo-degradation accelerator to reduce the stability of ascorbic acid under UV-irradiation (177-183).

Commercial Application of Ascorbic Acid:

Based on the afore-mentioned biological activity of ascorbic acid, ascorbic acid is mainly used as an antioxidant to inhibit food browning and as a dietary supplement for humans. Ascorbic acid is mainly used as an antioxidant to protect the senses of foods. As is well known, polyphenol oxidase catalyzes the enzymatic browning of phenol substrates to yield dark-colored melanin. Browning affects product sensory qualities and reduces consumer acceptance. Adding xyloglucan microcapsules containing ascorbic acid to baked foods such as tilapia fish burgers can significantly inhibit the browning that occurs during the preparation process and maintain the sensory qualities of the product. The chitosan/tripolyphosphate nano-aggregates containing ascorbic acid enhanced the inhibition of mushroom slices browning induced by tyrosinase. Acute heat stress during transport is known to predispose rainbow trout quality to deterioration, with negative effects on the histological, physicochemical and microbiological quality of fillets.

Treatment with added ascorbic acid partially mitigated damage caused by acute heat stress. It can maintain tissue structure, delay protein oxidation and then prolong the shelf life of fish fillets to about 2 days. In addition, as a nutritional supplement, ascorbic acid plays an important role as a co-factor in many biological processes. Unfortunately, fishes lack L-gluconolactone oxidase and cannot biosynthesize ascorbic acid by themselves, which is not conducive to the growth of their bone matrix and connective tissue. Lack of ascorbic acid can cause reduced wound-healing capacity and bone deformities in fish. At present, in the aquaculture area, ascorbic acid is widely added to fish diets. Based on the healthcare function of ascorbic acid, it is also vital in nutrition fortification products. As an important source of protein supplementation, dairy products are popular beverages all over the world. At present, milk and soymilk have been fortified with ascorbic acid, including ascorbate and ascorbic acid isomers, to improve the iron absorption in the small intestine. Food fortification can improve micronutrient malnutrition. It is worth noting that a category of foods tailored according to the necessary nutrients for a healthy life and their specific concentrations and ratios are called designer foods, also known as health foods, and are sought after and recognized by consumers. Such products often contain a variety of bioactive compounds. By adding calcium and antioxidants such as vitamins E and C to low-fat chicken patties, a high-quality product with high-quality animal protein, fat, multivitamins and minerals can be prepared. Ascorbic acid not only acted as a nutritional additive, but also maintained better color and flavor of chicken patties, and inhibited the formation of nitrosamines in the meat. The addition of sodium ascorbate and vitamin A to pig feed can significantly improve the growth performance, antioxidant capacity and immune function of weaned piglets. Meanwhile, as an antioxidant, sodium ascorbate can delay the degradation of vitamin A. A cornstarch-based baking premix was developed by addition of vitamin B, vitamin C and digestible iron, zinc, selenium and iodine. Although the added ascorbic acid in the baked bread degraded due to high temperature, it strengthened the structure of the bread and was benefit to product quality. Meanwhile, it was found that the combination of butylated hydroxytoluene and ascorbic acid significantly inhibited the oxidation and isomerization of vitamin A in skim milk powder during thermally accelerated storage. There are two major aspects in the current development of ascorbic acid-fortified products. On the one hand, the natural ascorbic acid is directly added, in order to use its antioxidant activity to maintain the sensory appearance of the product during the shelf life. The cost is low, but the retention activity of the final product is low. Another aspect is the addition of ascorbic acid derivatives, which is to ensure that sufficient physiological activity can be expressed after ingestion of the product. However, the cost of ascorbic acid derivatives is high, and they need to be converted before they can exert their functional properties. Although various delivery technologies are available, they are still in the developmental stage of industrial transformation and have not been widely used. Combined with the above analysis of delivery strategies, these may be limited by the cost of wall materials, and the complexity of the process, which is not suitable for large-scale industrial production. Therefore, researchers still need to explore low-cost, simple, and high-yield encapsulation techniques of ascorbic acid for industrial application (179- 184).

Conclusions:

In this review, the bioactivity and stability of ascorbic acid are introduced. There are many strategies for improving the bioavailability of ascorbic acid, and the influence of delivery systems on the stability and release properties of ascorbic acid is discussed. Besides the addition of low-molecular-weight antioxidants and preservatives, encapsulation technology is more and more widely used in the food field. The stabilization mechanism includes chemical chelation and physical barrier. Since the positively charged chitosan can interact with ascorbic acid through electrostatic interaction and hydrogen bond, it is the most superior carrier material. The complex system is mostly in the form of nano-sized particles. On the other hand, biomacromolecules can construct microcapsules with coatings through a series of technologies, such as

spray drying, microfluidic technique and complex coacervation. The physical barrier restricts ascorbic acid within the inner core, reducing the contact between it and the external environment. The two mechanisms have their own limitations. For example, chemically complexed nanoparticles are beneficial to the absorption by mucosal membranes, but the encapsulation efficiency of ascorbic acid is low and ascorbic acid is accessible to solvent. The coating of the microcapsules can effectively shield the inner ascorbic acid from external environment, but the operation is more complicated and requires the assistance of a variety of equipment. In addition, the larger size and poor water-solubility of the microcapsules limit the absorption in the body to a certain extent. Therefore, understanding the pro-degradation factors of ascorbic acid and its properties are conducive to the targeted design of delivery systems. Adopting low-cost methods to design an effective fortification strategy to improve the stability of ascorbic acid during processing and storage is still the focus and challenge for researchers.

Conflicts of Interest

The authors declare no conflicts of interest.

References:

- Shaltout, F.A., Riad, E.M., and AbouElhassan, Asmaa, A. (2017): prevalence Of Mycobacterium Tuberculosis In Imported cattle Offals And Its lymph Nodes. *Veterinary Medical Journal -Giza (VMJG)*, 63(2): 115 – 122.
- Sripakdee, T.; Mahachai, R.; Chanthai, S. Phenolics and Ascorbic Acid Related to Antioxidant Activity of MaoFruit Juice and Their Thermal Stability Study (Review Article). *Orient. J. Chem.* 2017, 33, 74–86
- Shaltout, F.A., Riad, E.M., and Asmaa Abou-Elhassan (2017): Prevalence Of Mycobacterium Spp. In Cattle Meat and Offal's Slaughtered In And Out Abattoir. *Egyptian Veterinary medical Association*, 77(2): 407 – 420.
- Abd Elaziz, O., Fatin, S. Hassanin, Fahim A. Shaltout and Othman A. Mohamed (2021): Prevalence of Some Foodborne Parasitic Affection in Slaughtered Animals in Local Egyptian Abattoir. *Journal of Nutrition Food Science and Technology* 2(3): 1-5.
- Abd Elaziz, O., Fatin, S. Hassanin, Fahim, A. Shaltout, Othman, A. Mohamed (2021): Prevalence of some zoonotic parasitic affections in sheep carcasses in a local abattoir in Cairo, Egypt. *Advances in Nutrition & Food Science* 6(2): 6(2): 25-31.
- AlShorman, A.A.M.; Shaltout, F.A. and Hilat, N. (1999): Detection of certain hormone residues in meat marketed in Jordan. *Jordan University of Science and Technology*, 1st International Conference on Sheep and goat Diseases and Productivity, 23-25 October, 1999.
- Ebeed Saleh, Fahim Shaltout, Essam Abd Elaal (2021); Effect of some organic acids on microbial quality of dressed cattle carcasses in Damietta abattoirs, Egypt. *Damanhour Journal of Veterinary Sciences* 5(2): 17-20.
- Edris A, Hassanin, F. S; Shaltout, F.A., Azza H Elbaba and Nairoz M Adel (2017): Microbiological Evaluation of Some Heat-Treated Fish Products in Egyptian Markets. *EC Nutrition* 12.3 (2017): 124-132.
- De Lorenzo, A.; Andreoli, A.; Sinibaldi Salimei, P.; D'Orazio; Guidi, A.; Ghiselli, A. Determination of the blood ascorbic acid level after administration of slow-release vitamin C. *Clin. Ter.* 2001, 152, 87–90.
- Edris, A., Hassan, M.A., Shaltout, F.A. and Elhosseiny, S. (2013): Chemical evaluation of cattle and camel meat. *BENHA VETERINARY MEDICAL JOURNAL*, 24(2): 191-197.
- Edris, A.M., Hassan, M.A., Shaltout, F.A. and Elhosseiny, S. (2012): Detection of E.coli and Salmonella organisms in cattle and camel meat. *BENHA VETERINARY MEDICAL JOURNAL*, 24(2): 198-204.
- Edris A.M.; Hemmat M. I., Shaltout F.A.; Elshater M.A., Eman F.M.I. (2012): STUDY ON INCIPIENT SPOILAGE OF CHILLED CHICKEN CUTS-UP. *BENHA VETERINARY MEDICAL JOURNAL*, VOL. 23, NO. 1, JUNE 2012: 81-86.
- Edris A.M.; Hemmat M.I.; Shaltout F.A.; Elshater M.A., Eman, F.M.I. (2012): CHEMICAL ANALYSIS OF CHICKEN MEAT WITH RELATION TO ITS QUALITY. *BENHA VETERINARY MEDICAL JOURNAL*, 23(1): 87-92
- Edris, A.M.; Shaltout, F.A. and Abd Allah, A.M. (2005): Incidence of Bacillus cereus in some meat products and the effect of cooking on its survival. *Zag. Vet. J.* 33 (2):118-124.
- Edris, A.M.; Shaltout, F.A. and Arab, W.S. (2005): Bacterial Evaluation of Quail Meat. *Benha Vet. Med.J.* 16 (1):1-14.
- Edris, A.M.; Shaltout, F.A.; Salem, G.H. and El-Toukhy, E.I. (2011): Incidence and isolation of Salmonellae from some meat products. *Benha University, Faculty of Veterinary Medicine, Fourth Scientific Conference 25-27th May 2011 Veterinary Medicine and Food Safety*) 172-179 benha, Egypt.
- Edris AA, Hassanin, F. S; Shaltout, F.A., Azza H Elbaba and Nairoz M Adel. (2017): Microbiological Evaluation of Some Heat-Treated Fish Products in Egyptian Markets. *EC Nutrition* 12.3 (2017): 134-142.
- Edris, A.M.; Shaltout, F.A.; Salem, G.H. and El-Toukhy, E.I. (2011): Plasmid profile analysis of Salmonellae isolated from some meat products. *Benha University, Faculty of Veterinary Medicine, Fourth Scientific Conference 25-27th May 2011 Veterinary Medicine and Food Safety*) 194-201 benha, Egypt.
- Yang, S.; Verhoeff, A.A.; Merks, D.W.H.; van Duynhoven, J.P.M.; Hohlbein, J. Quantitative Spatiotemporal Mapping of Lipid and Protein Oxidation in Mayonnaise. *Antioxidants* 2020
- Ragab A, Abobakr M. Edris, Fahim A.E. Shaltout, Amani M. Salem (2022): Effect of titanium dioxide nanoparticles and thyme essential oil on the quality of the chicken fillet. *BENHA VETERINARY MEDICAL JOURNAL* 41(2): 38-40.
- Hassan, M.A, Shaltout, F. A, Arfa M.M, Mansour A.H and Saudi, K. R (2013): BIOCHEMICAL STUDIES ON RABBIT MEAT RELATED TO SOME DISEASES. *BENHA VETERINARY MEDICAL JOURNAL* 25(1):88-93.
- Hassan, M. A and Shaltout, F.A. (1997): Occurrence of Some Food Poisoning Microorganisms In Rabbit Carcasses *Alex.J.Vet.Science*, 13(1):55-61.
- Hassan M, Shaltout FA* and Saqr N (2020): Histamine in Some Fish Products. *Archives of Animal Husbandry & Dairy Science* 2(1): 1-3.
- Hassan, M.A and Shaltout, F.A. (2004): Comparative Study on Storage Stability of Beef, Chicken meat, and Fish at Chilling Temperature. *Alex.J.Vet.Science*, 20(21):21-30.
- Hassan, M.A; Shaltout, F.A.; Arafa, M.M.; Mansour, A.H. and Saudi, K.R. (2013): Biochemical studies on rabbit meat related to some diseases. *Benha Vet. Med.J.* 25 (1):88-93.
- Hassan, M. A; Shaltout, F.A.; Maarouf, A.A. and El-Shafey, W.S. (2014): Psychrotrophic bacteria in frozen fish with special reference to pseudomonas species. *Benha Vet. Med.J.* 27 (1):78-83.
- Hwang, K.E.; Kim, H.W.; Song, D.H.; Kim, Y.J.; Ham, Y.K.; Choi, Y.S.; Lee, M.A.; Kim, C.J. Effect of Mugwort and Rosemary Either Singly, or Combination with Ascorbic Acid on Shelf Stability of Pork Patties. *J. Food Process. Preserv.* 2017, 41, e12994
- Hassan, M.A; Shaltout, F.A.; Arafa, M.M.; Mansour, A.H. and Saudi, K.R. (2013): Bacteriological studies on rabbit meat related to some diseases *Benha Vet. Med.J.* 25 (1):94-99.
- Hassanin, F. S; Hassan, M.A., Shaltout, F.A., Nahla A. Shawqy and 2Ghada A. Abd-Elhameed (2017): Chemical criteria of

- chicken meat. BENHA VETERINARY MEDICAL JOURNAL, 33(2):457-464.
30. Shaltout, F. A. (2024). Egyptian Medicinal Plants and Respiratory Disease. *Journal of Agriculture and Education Research*. 2 (3), 1-7.
 31. Hassanin, F. S; Hassan, M.A.; Shaltout, F.A. and Elrais-Amina, M (2014): CLOSTRIDIUM PERFRINGENS IN VACUUM PACKAGED MEAT PRODUCTS. BENHA VETERINARY MEDICAL JOURNAL, 26(1):49-53.
 32. Hassanien, F.S.; Shaltout, F.A.; Fahmey, M.Z. and Elsukkary, H.F. (2020): Bacteriological quality guides in local and imported beef and their relation to public health. *Benha Veterinary Medical Journal* 39: 125-129.
 33. Hassanin, F. S; Shaltout, F.A. and , Mostafa E.M (2013): Parasitic affections in edible offal. *Benha Vet. Med.J.* 25 (2):34-39.
 34. Hassanin, F. S; Shaltout, F.A., Lamada, H.M., Abd Allah, E.M. (2011): THE EFFECT OF PRESERVATIVE (NISIN) ON THE SURVIVAL OF LISTERIA MONOCYTOGENES. BENHA VETERINARY MEDICAL JOURNAL (2011)-SPECIAL ISSUE [I]: 141-145.
 35. Shaltout FA. Dry-Aged Meat and their Importance. *Open J of Frail Sci* 2024, 2(1): 000111. DOI: 10.23880/oajfs-16000111
 36. Fujii, K.; Yasuda, A.; Orikoshi, E.; Kimura, Y.; Chaen, H. Milk Constituent-Containing Food Reinforced in Vitamin C. Patent 2006320222 A, 30 November 2006
 37. Khattab, E., Fahim Shaltout and Islam Sabik (2021): Hepatitis A virus related to foods. BENHA VETERINARY MEDICAL JOURNAL 40(1): 174-179.
 38. Shaltout, F. A. Human Parasites in Relation to Contaminated Food and Drinking Water. *J Biomed Sci Biotech Res*. 2024. 2(1): 1-5. DOI: doi.org/10.61440/JBSBR. 2024.v2.02
 39. Saad M. Saad, Fahim A. Shaltout , Amal A. A. Farag & Hashim F. Mohammed (2022): Organophosphorus Residues in Fish in Rural Areas. *Journal of Progress in Engineering and Physical Science* 1(1): 27-31.
 40. Shaltout FAE. Everything about Nutritional Value of the Meat Ingredients and How we can Reduce its Microbial Hazards. *J Vet Sci Res* 2025, 10(1): 000283. DOI: 10.23880/oajvsr-16000283
 41. Al Fata, N.; George, S.; Dlalsh, N.; Renard, C.M.G.C. Influence of partial pressure of oxygen on ascorbic acid degradation at canning temperature. *Innov. Food Sci. Emerg. Technol.* 2018, 49, 215–221.
 42. Saif, M., Saad S.M., Hassanin, F. S; Shaltout FA, Marionette Zaghloul (2019): Molecular detection of enterotoxigenic *Staphylococcus aureus* in ready-to-eat beef products. *Benha Veterinary Medical Journal* 37 (2019) 7-11.
 43. Saif, M., Saad S.M., Hassanin, F. S; Shaltout, F.A., Marionette Zaghloul (2019); Prevalence of methicillin-resistant *Staphylococcus aureus* in some ready-to-eat meat products. *Benha Veterinary Medical Journal* 37 (2019) 12-15.
 44. Farag, A. A., Saad M. Saad¹, Fahim A. Shaltout¹, Hashim F. Mohammed (2023 a): Studies on Pesticides Residues in Fish in Menofia Governorate. *Benha Journal of Applied Sciences*, 8(5): 323-330.
 45. Shaltout, F. A. (2024): The concept of meat analysis in economy and public health, *Dietary Nourishment and Food Processing Techniques (DNFPT)* 1(1) 1-7.
 46. Farag, A. A., Saad M. Saad¹, Fahim A. Shaltout¹, Hashim F. Mohammed (2023 b): Organochlorine Residues in Fish in Rural Areas. *Benha Journal of Applied Sciences*, 8 (5): 331-336.
 47. Shaltout, F.A., Mona N. Hussein, Nada Kh. Elsayed (2023): Histological Detection of Unauthorized Herbal and Animal Contents in Some Meat Products. *Journal of Advanced Veterinary Research* 13(2): 157-160.
 48. Shaltout, F. A., Heikal, G. I., Ghanem, A. M. (2022): Mycological quality of some chicken meat cuts in Gharbiya governorate with special reference to *Aspergillus flavus* virulent factors. *benha veteriv medical journal veterinary* 42(1): 12-16.
 49. Shaltout, F.A., Ramadan M. Salem, Eman M. Eldiasty, Fatma A. Diab (2022): Seasonal Impact on the Prevalence of Yeast Contamination of Chicken Meat Products and Edible Giblets. *Journal of Advanced Veterinary Research* 12(5): 641-644.
 50. Shaltout, F.A., Abdelazez Ahmed Helmy Barr and Mohamed Elsayed Abdelaziz (2022): Pathogenic Microorganisms in Meat Products. *Biomedical Journal of Scientific & Technical Research* 41(4): 32836-32843.
 51. Shaltout, F.A., Thabet, M.G. and Koura, H.A. (2017). Impact of Some Essential Oils on the Quality Aspect and Shelf Life of Meat. *J Nutr Food Sci.*, 7: 647.
 52. Shaltout, F.A., Islam Z. Mohammed², El -Sayed A. Afify (2020): Bacteriological profile of some raw chicken meat cuts in Ismailia city, Egypt. *Benha Veterinary Medical Journal* 39 (2020) 11-15.
 53. Shaltout, F.A., Islam, Z. Mohammed², El -Sayed A. Afify (2020): Detection of *E. coli* O157 and *Salmonella* species in some raw chicken meat cuts in Ismailia province, Egypt. *Benha Veterinary Medical Journal* 39 (2020) 101-104.
 54. Shaltout, F.A., E.M. El-diasty and M. A. Asmaa- Hassan (2020): HYGIENIC QUALITY OF READY TO EAT COOKED MEAT IN RESTAURANTS AT Cairo. *Journal of Global Biosciences* 8(12): 6627-6641.
 55. Shaltout, F.A., Marrionet Z. Nasief, L. M. Lotfy, Bossi T. Gamil (2019): Microbiological status of chicken cuts and its products. *Benha Veterinary Medical Journal* 37 (2019) 57-63.
 56. Shaltout, F.A. (2019): Poultry Meat. *Scholarly Journal of Food and Nutrition* 22 1-2.
 57. Shaltout, F.A. (2019): Food Hygiene and Control. *Food Science and Nutrition Technology* 4(5): 1-2.
 58. Hassanin, F. S; Shaltout, F.A., Seham N. Homouda and Safaa M. Arakeeb (2019): Natural preservatives in raw chicken meat. *Benha Veterinary Medical Journal* 37 (2019) 41-45.
 59. Shaltout, D. E. (2024): Additives Extend the Food Shelf Life by Addition of Preservatives Nitrate, and Nitrite to Food, *Dietary Nourishment and Food Processing Techniques*, 1(3): 1-12. DOI: 10.9567/3064-7061/WSJ.83
 60. Hazaa, W., Shaltout, F.A., Mohamed El-Shate (2019): Prevalence of some chemical hazards in some meat products. *Benha Veterinary Medical Journal* 37 (2) 32-36.
 61. Akyildiz, A.; Mertoglu, T.S.; Agcam, E. Kinetic study for ascorbic acid degradation, hydroxymethylfurfural and furfural formations in orange juice. *J. Food Compos. Anal.* 2021, 102, 103996
 62. Shaltout, F. A. E. (2024): Using of Meat Diets as a Functional Food, *Dietary Nourishment and Food Processing Techniques*, vol 1(3): 1-14
 63. Shaltout, F. A. (2024) Evaluation of Hazards in food, *Journal of Medical Discoveries*, 1(1);1-8
 64. Hazaa, W., Shaltout, F.A., Mohamed El-Shater (2019): Identification of Some Biological Hazards in Some Meat Products. *Benha Veterinary Medical Journal* 37 (2) 27-31.
 65. Shaltout, F. A. (2024): Through a light on Meat as Functional food, *International Journal of Nursing Didactics*, 14 (08): 1-12. DOI: <https://doi.org/10.52845/IJND/2024/14-08-1>
 66. Gaafar, R., Hassanin, F. S; Shaltout, F.A., Marionette Zaghloul (2019): Molecular detection of enterotoxigenic *Staphylococcus aureus* in some ready to eat meat-based sandwiches. *Benha Veterinary Medical Journal* 37 (2) 22-26.
 67. Shaltout F. (2019) Microbial Contamination of Beef and Beef Products. *J. Nutrition and Food Processing*, 2(2): 1;

68. Gaafar, R., Hassanin, F. S.; Shaltout, F.A., Marionette Zaghoul (2019): Hygienic profile of some ready to eat meat product sandwiches sold in Benha city, Qalubia Governorate, Egypt. Benha Veterinary Medical Journal 37 (2) 16-21.
69. Shaltout, F. A. (2024): Abattoir and Bovine Tuberculosis as a Reemerging Foodborne Disease. Biomed J Sci & Tech Res 54(3)-2024. BJSTR. MS.ID.008545.
70. Saad S.M., Shaltout, F.A., Nahla A Abou Elroos, Saber B El-nahas (2019): Antimicrobial Effect of Some Essential Oils on Some Pathogenic Bacteria in Minced Meat. J Food Sci Nutr Res. 2019; 2 (1): 012-020.
71. Shaltout, F. A. E. (2024): Good News about Application of Advanced Methods in Food Examination, Dietary Nourishment and Food Processing Techniques, vol 1(3): 1-9.
72. Saad S.M., Shaltout, F.A., Nahla A Abou Elroos and Saber B El-nahas (2019): Incidence of Staphylococci and E. coli in Meat and Some Meat Products. EC Nutrition 14.6 (2019).
73. Przekwas, J.; Wiktorczyk, N.; Budzynska, A.; Walecka-Zacharska, E.; Gospodarek-Komkowska, E. Ascorbic Acid Changes Growth of Food-Borne Pathogens in the Early Stage of Biofilm Formation. Microorganisms 2020, 8, 10
74. Shaltout, F. A. E. (2024): Our options to improve food safety and quality by using preservatives which are used in food processing and preservation, Dietary Nourishment and Food Processing Techniques, vol 1(3): 1-16.
75. Saad S.M., Hassanin, F. S.; Shaltout, F.A., Marionette Z Nassif, Marwa Z Seif. (2019): Prevalence of Methicillin-Resistant Staphylococcus Aureus in Some Ready-to-Eat Meat Products. American Journal of Biomedical Science & Research 4(6):460-464.
76. Shaltout, Fahim (2019): Pollution of Chicken Meat and Its Products by Heavy Metals. Research and Reviews on Healthcare: Open Access Journal, 4, 3(381-3382).
77. Shaltout, F. A.; E.M EL-diahy; M. S. M Mohamed (2018): Effects of chitosan on quality attributes fresh meat slices stored at 4 C. BENHA VETERINARY MEDICAL JOURNAL, VOL. 35, NO. 2: 157-168.
78. Shaltout and Abdel-Aziz, 2004: Salmonella enterica serovar Enteritidis in poultry meat and their epidemiology. Vet. Med. J. Giza, 52 (2004), pp. 429-436.
79. Shaltout, F.A., Hala F El-Shorah, Dina I El Zahaby, Lamiaa M Lotfy (2018): Bacteriological Profile of Chicken Meat Products. SciFed Food & Dairy Technology Journal, 2:3.
80. Scheffler, J.; Bork, K.; Bezold, V.; Rosenstock, P.; Gnanapragassam, V.S.; Horstkorte, R. Ascorbic acid leads to glycation and interferes with neurite outgrowth. Exp. Gerontol. 2019, 117, 25–30
81. Shaltout, F.A., Mohamed, A.H. El-Shater., Wafaa Mohamed Abd El-Aziz (2015): Bacteriological assessment of Street Vended Meat Products sandwiches in kalyobia Governorate. BENHA VETERINARY MEDICAL JOURNAL, 28(2):58-66,
82. Shaltout, F.A., Mohamed A El shatter and Heba M Fahim (2019): Studies on Antibiotic Residues in Beef and Effect of Cooking and Freezing on Antibiotic Residues Beef Samples. Scholarly Journal of Food and Nutrition 2(1) 1-4
83. Shaltout FA, Zakaria IM and Nabil ME. (2018): Incidence of Some Anaerobic Bacteria Isolated from Chicken Meat Products with Special Reference to Clostridium perfringens. Nutrition and Food Toxicology 2.5 (2018): 429-438.
84. Shaltout FA, Ahmed A A Maarouf and Mahmoud ES Elkhoully. (2017): Bacteriological Evaluation of Frozen Sausage. Nutrition and Food Toxicology 1.5; 174-185.
85. Shaltout FA, El-Toukhy EI and Abd El-Hai MM. (2019): Molecular Diagnosis of Salmonellae in Frozen Meat and Some Meat Products. Nutrition and Food Technology Open Access 5(1): 1-6.
86. Shaltout, F.A., A.M.Ali and S.M.Rashad (2016): Bacterial Contamination of Fast Foods. Benha Journal of Applied Sciences (BJAS) 1 (2)45-51.
87. Shaltout, F.A., Zakaria. I. M. , Jehan Eltanani , Asmaa . Elmelegy(2015): Microbiological status of meat and chicken received to University student hostel. BENHA VETERINARY MEDICAL JOURNAL, 29(2):187-192, DECEMBER, 2015.
88. Saad,S.M.;Edris, A.M.; Shaltout,F.A. and Edris, Shima(2012): Isolation and identification of salmonellae and E.coli from meat and poultry cuts by using A.multiplex PCR. Benha Vet. Med.J.special issue 16-26.
89. Saad, S.M. and Shaltout, F.A. (1998): Mycological Evaluation of camel carcasses at Kalyobia Abattoirs. Vet.Med.J. Giza,46(3):223-229.
90. Zhao, R.N.; Yuan, Y.H.; Liu, F.Y.; Han, J.G.; Sheng, L.S. A computational investigation on the geometries, stabilities, antioxidant activity, and the substituent effects of the L-ascorbic acid and their derivatives. Int. J. Quantum Chem. 2013, 113, 2220–2227
91. Shaltout, F. A. (2024): Whey We Extend the Food Shelf Life by Aid of Natural Antioxidants? Biomed J Sci & Tech Res 59(1)-2024. BJSTR. MS.ID.009235
92. Saad S.M., Shaltout, F.A., Nahla A Abou Elroos, Saber B El-nahas. 2019: Antimicrobial Effect of Some Essential Oils on Some Pathogenic Bacteria in Minced Meat. J Food Sci Nutr Res. 2019; 2 (1): 012-020.
93. Saad S.M., Hassanin, F. S; Shaltout, F.A., Marionette Z Nassif, Marwa Z Seif. (2019): Prevalence of Methicillin-Resistant Staphylococcus Aureus in Some Ready-to-Eat Meat Products. American Journal of Biomedical Science & Research 4(6):460-464.
94. Saad S.M., Shaltout, F.A., Nahla A Abou Elroos and Saber B El-nahas. (2019): Incidence of Staphylococci and E. coli in Meat and Some Meat Products. EC Nutrition 14.6 (2019).
95. Shaltout FA, Riad EM, TES Ahmed and AbouElhassan A (2017): Studying the Effect of Gamma Irradiation on Bovine Offal's Infected with Mycobacterium tuberculosis Bovine Type. Journal of Food Biotechnology Research 1 (6): 1-5.
96. Shaltout FA, Zakaria IM and Nabil ME. (2018): Incidence of Some Anaerobic Bacteria Isolated from Chicken Meat Products with Special Reference to Clostridium perfringens. Nutrition and Food Toxicology 2.5 (2018): 429-438.
97. Shaltout FA, Mohamed, A.Hassan and Hassanin, F. S(2004): THERMAL INACTIVATION OF ENTEROHAEMORRHAGIC ESCHERICHIA COLI O157:H7 AND ITS SENSITIVITY TO NISIN AND LACTIC ACID CULTURES. 1st Ann. Confr., FVM., Moshtohor, Sept, 2004.
98. Shaltout FA, El-diahy, E, M;Elmeslamy, M. and Elshaer, M.(2014): Study on fungal contamination of some chicken meat products with special reference to 2 the use of PCR for its identification . Conference,Veterinary Medical Journal – Giza vol. December 2014/12/17 vol.60: 1-10.
99. shaltout, F.A. (2002): Microbiological Aspects of Semi-cooked chicken Meat Products. Benha Veterinary Medical Journal13, 2, 15-26.
100. Yuan, J.P.; Chen, F. Degradation of Ascorbic Acid in Aqueous Solution. J. Agric. Food Chem. 1998, 46, 5078–5082
101. Shaltout FA, Thabet, M.G2 and Hanan, A. Koura3. (2017): Impact of some essential oils on the quality aspect and shelf life of meat.BENHA VETERINARY MEDICAL JOURNAL, 33, (2): 351-364.
102. Shaltout FA, Mohammed Farouk; Hosam A.A. Ibrahim and Mostafa E.M. Afifi4.2017: Incidence of Coliform and Staphylococcus aureus in ready to eat fast foods. BENHA VETERINARY MEDICAL JOURNAL, 32(1): 13 - 17, MARCH, 2017.

103. Shaltout, F.A., Zakaria, I.M., Nabil, M.E. (2017): Detection and typing of *Clostridium perfringens* in some retail chicken meat products. *BENHA VETERINARY MEDICAL JOURNAL*, 33(2):283-291.
104. Shaltout, F.A. (1992): Studies on Mycotoxins in Meat and Meat by Products. M.V.Sc Thesis Faculty of Veterinary Medicine, Moshtohor, Zagazig University Benha branch.
105. Shaltout, F.A. (1996): Mycological and Mycotoxicological profile Of Some Meat products. Ph.D.Thesis, Faculty of Veterinary Medicine, Moshtohor, Zagazig University Benha branch.
106. Shaltout, F.A. (1998): Proteolytic Psychrotrophes in Some Meat products. *Alex. Vet. Med. J.* 14 (2):97-107.
107. Shaltout, F.A. (1999): Anaerobic Bacteria in Vacuum Packed Meat Products. *Benha Vet. Med.J.* 10 (1):1-10.
108. Shaltout, F.A. (2000): Protozoal Foodborne Pathogens in some Meat Products. *Assiut Vet. Med. J.* 42 (84):54-59.
109. Shaltout, F.A. (2001): Quality evaluation of sheep carcasses slaughtered at Kalyobia abattoirs. *Assiut Veterinary Medical Journal*, 46(91):150-159.
110. Zhao, R.N.; Yuan, Y.H.; Liu, F.Y.; Han, J.G.; Sheng, L.S. A computational investigation on the geometries, stabilities, antioxidant activity, and the substituent effects of the L-ascorbic acid and their derivatives. *Int. J. Quantum Chem.* 2013, 113, 2220–2227
111. Shaltout, F.A. (2002): Microbiological Aspects of Semi-cooked Chicken Meat Products. *Benha Vet. Med.J.* 13(2):15-26.
112. Shaltout, F.A. (2003): *Yersinia Enterocolitica* in some meat products and fish marketed at Benha city. The Third international conference Mansoura 29-30 April.
113. Shaltout, F.A. (2009): Microbiological quality of chicken carcasses at modern Poultry plant. The 3rd Scientific Conference, Faculty of Vet. Med., Benha University, 1-3 january.
114. Shaltout, F.A. and Abdel Aziz ,A.M. (2004): *Salmonella enterica* Serovar Enteritidis in Poultry Meat and their Epidemiology .*Vet. Med.J.*, Giza, 52(3):429-436.
115. Shaltout, F.A. and Abdel Aziz ,A.M. (2004): *ESCHERICHIA COLI* STRAINS IN SLAUGHTERED ANIMALS AND THEIR PUBLIC HEALTH IMPORTENCE. *J.Egypt. Vet. Med. Association* 64(2):7-21.
116. Shaltout, F.A., Amin, R., Marionet , Z., Nassif and Shimaa, Abdel-wahab (2014): Detection of aflatoxins in some meat products. *Benha veterinary medical journal*, 27(2) :368-374.
117. Shaltout, F.A. and Afify , Jehan Riad, EM and Abo Elhasan , Asmaa, A. (2012): Improvement of microbiological status of oriental sausage. *Journal of Egyptian Veterinary Medical Association* 72(2):157-167.
118. Shaltout, F.A. and Daoud, J. R. (1996): Chemical analytical studies on rabbit meat and liver. *Benha Vet. Med.J.* 8 (2):17-27.
119. Shaltout, F.A. and Edris, A.M. (1999): Contamination of shawerma with pathogenic yeasts. *Assiut Veterinary Medical Journal*, 40(64):34-39.
120. Alvim, I.D.; Stein, M.A.; Koury, I.P.; Dantas, F.B.H.; Cruz, C. Comparison between the spray drying and spray chilling microparticles contain ascorbic acid in a baked product application. *Lwt-Food Sci. Technol.* 2016, 65, 689–694
121. Su, L.J.; Zhang, J.H.; Gomez, H.; Murugan, R.; Hong, X.; Xu, D.; Jiang, F.; Peng, Z.Y. Reactive Oxygen Species-Induced Lipid Peroxidation in Apoptosis, Autophagy, and Ferroptosis. *Oxid. Med. Cell. Longev.* 2019, 2019, 5080843.
122. Shaltout, F. A.; Eldiasty , E. and Mohamed , M.S. (2014): Incidence of lipolytic and proteolytic fungi in some chicken meat products and their public health significance. *Animal Health Research Institute: First International Conference on Food Safety and Technology* 19-23 June 2014 Cairo Egypt pages 79-89.
123. Shaltout, F.A.; Eldiasty, E.; Salem, R. and Hassan, Asmaa (2016): Mycological quality of chicken carcasses and extending shelf – life by using preservatives at refrigerated storage. *Veterinary Medical Journal -Giza (VMJG)* 62(3)1-7.
124. Shaltout, F.A.; Salem, R. Eldiasty, E.; and Diab, Fatema. (2016): Mycological evaluation of some ready to eat meat products with special reference to molecular chacterization. *Veterinary Medical Journal -Giza* 62(3)9-14.
125. Shaltout, F. A.; Elshater, M. and Wafaa, Abdelaziz (2015): Bacteriological assessment of street vended meat products sandwiches in Kalyobia Governorate. *Benha Vet. Med.J.* 28 (2):58-66.
126. Shaltout, F. A.; Gerges, M.T. and Shewail, A.A. (2018): Impact of Organic Acids and Their Salts on Microbial Quality and Shelf Life of Beef. *Assiut veterinary medical journal* 64(159): 164-177
127. Shaltout, F.A.; Ghoneim, A.M.; Essmail, M.E. and Yousseif ,A. (2001): Studies on aflatoxin B1 residues in rabbits and their pathological effects. *J.Egypt. Vet. Med. Association* 61(2):85-103.
128. Shaltout, F.A. and Hanan ,M.T. El-Lawendy (2003): Heavy Metal Residues In Shawerma. *Beni-Suef Vet. Med.J.* 13(1):213-224.
129. Shaltout, F.A. and Hashim, M.F. (2002): Histamine in salted, Smoked and Canned Fish products. *Benha Vet. Med.J.* 13 (1):1-11.
130. Chang, D.W.; Abbas, S.; Hayat, K.; Xia, S.Q.; Zhang, X.M.; Xie, M.Y.; Kim, J.M. Encapsulation of ascorbic acid in amorphous maltodextrin employing extrusion as affected by matrix/corn ratio and water content. *Int. J. Food Sci. Technol.* 2010, 45, 1895–1901.
131. Shaltout, F.A.; Hashim, M.F. and Elnahas, s. (2015): Levels of some heavy metals in fish (*tilapia nilotica* and *Claris lazera*) at Menufia Governorate. *Benha Vet. Med.J.* 29 (1):56-64.
132. Shaltout, F.A. and Ibrahim, H.M. (1997): Quality evaluation of luncheon and Alexandrian sausage. *Benha Vet. Med.J.* 10 (1):1-10.
133. Shaltout, F.A.; Nassif, M and Shakran, A (2014): Quality of battered and breaded chicken meat products. *Global Journal of Agriculture and Food Safety Science – 1(2)* ISSN 2356-7775.
134. Shaltout, F.A., Amani M. Salem, A. H. Mahmoud, K. A (2013): Bacterial aspect of cooked meat and offal at street vendor's level. *Benha veterinary medical journal*, 24(1): 320-328.
135. Shaltout, F.A. and Salem, R.M. (2000): Moulds, aflatoxin B1 and Ochratoxin A in Frozen Livers and meat products. *Vet . Med. J.Giza* 48(3):341-346.
136. Yasser H. Al-Tarazi, A. Al-Zamil, Shaltout FA. and H. Abdel-Samei (2002). Microbiological status of raw cow milk marketed in northern Jordan. *AVMJ Volume 49 Issue 96 Pages* 180-194
137. Shaltout FA, Zakaria IM and Nabil ME. (2018): Incidence of Some Anaerobic Bacteria Isolated from Chicken Meat Products with Special Reference to *Clostridium perfringens*. *Nutrition and Food Toxicology* 2(5):429-438.
138. Shaltout, F. A.; El-diasty, E.M. and Mohamed, M. S. (2014): Incidence of lipolytic and proteolytic fungi in some chicken meat products and their public health significance. 1st Scientific conference of food safety and Technology .2014, pp. 79-89.
139. Shaltout, F. A.; El-diasty, E.M.; Salem, R. M. and Asmaa, M. A. Hassan. 2016: Mycological quality of chicken carcasses and extending shelf -life by using preservatives at refrigerated storage. *Veterinary Medical Journal – Giza* ,62(3) :1-10.
140. Bei, R. Effects of vitamin C on health: A review of evidence. *Front. Biosci.* 2013, 18, 1017–1029

141. Shaltout FA, R.M. Salem, E.M. El-Diasty and W.I.M. Hassan. 2019: Effect of Lemon Fruits and Turmeric Extracts on Fungal Pathogens in Refrigerated Chicken Fillet Meat. *Global Veterinaria* 21 (3): 156-160.
142. Shaltout FA, El-diasty, E. M.; Elmesalamy, M. and Elshaer, M. (2014): Study on fungal contamination of some chicken meat products with special reference to the use of PCR for its identification. *Conference, Veterinary Medical Journal – Giza* vol. December 2014/12/17 vol.60 1-10.
143. Shaltout, F. A.; Salem, R. M; El-diasty, Eman and Fatema, A.H. Diab. (2016): Mycological evaluation of some ready to eat meat products with special reference to molecular characterization. *Veterinary Medical Journal – Giza*. 62(3): 9-14.
144. Shaltout FA, Ahmed, A.A. Maarouf, Eman, M.K. Ahmed (2018): Heavy Metal Residues in chicken cuts up and processed chicken meat products. *BENHA VETERINARY MEDICAL JOURNAL*, 34(1): 473-483.
145. Shaltout, F.A.; Hanan M. Lamada, Ehsan A.M. Edris. (2020): Bacteriological examination of some ready to eat meat and chicken meals. *Biomed J Sci & Tech Res.*, 27(1): 20461-20465.
146. Sobhy, Asmaa and Shaltout, Fahim (2020): Prevalence of some food poisoning bacteria in semi cooked chicken meat products at Qaliubiya governorate by recent Vitek 2 compact and PCR techniques. *Benha Veterinary Medical Journal* 38 (2020) 88-92.
147. Shaltout, F. A. (2024): Good Idea on Preservatives and the Natural Preservatives and Meat Preservation Against the Foodborne Pathogens and the Spoilage Microorganisms. *Biomed J Sci & Tech Res* 57(5)-2024. BJSTR. MS.ID.009067.
148. Sobhy, Asmaa and Shaltout, Fahim (2020): Detection of food poisoning bacteria in some semi-cooked chicken meat products marketed at Qaliubiya governorate. *Benha Veterinary Medical Journal* 38 (2020) 93-96.
149. Shaltout, F.A. (2024): Abattoir and Bovine Tuberculosis as A Reemerging Foodborne Diseases. *Clinical Medical Reviews and Report* 6(1):1-7.
150. Padayatty, S.J.; Sun, H.; Wang, Y.H.; Riordan, H.D.; Hewitt, S.M.; Katz, A.; Wesley, R.A.; Levine, M. Vitamin C pharmacokinetics: Implications for oral and intravenous use. *Ann. Intern. Med.* 2004, 140, 533–537
151. Shaltout, F.A. (2023): Viruses in Beef, Mutton, Chevon, Venison, Fish and Poultry Meat Products. *Food Science & Nutrition Technology* 8(4):1-10.
152. Shaltout, F. A. (2024): Human Salmonellosis Acquired through the Food". *Acta Scientific Pharmaceutical Sciences* 8. (3): 1-6: 12-17
153. Elkholy, R. A; Hussein, M. N; Abou El-Roos, N. A. and Shaltout, F.A.E. (2025) Enhancing Microbiological and Histological Quality of Frozen Turkey Meat Using Vinegar. *Egyptian Journal of Veterinary Sciences* pp 1-8. DOI: 10.21608/EJVS.2024.291707.2118
154. Shaltout, F. A. (2024): Availability, Price, Tradition, Religion, Income, Social, Development and Economic Influences on Meat Consumption. *Med J Clin Trials Case Stud* 2024, 8(2): 000370
155. Mohamed Q. M., Fahim A. Shaltout, f.A. and Ali, E.A. (2025): Multidrug-Resistant Bacteria from Raw Chevon and Mutton Meat. *Egyptian Journal of Veterinary Sciences* pp 1-8. DOI: 10.21608/EJVS.2024.310881.2302
156. Shaltout, F. A. E; Ab delazez Ahmed Helmy Barr, Mohamed Elsayed Abdelaziz (2024): Pathogenic Microorganisms in Meat Products. *Biomed J Sci & Tech Res* 41(4)-2022. BJSTR. MS.ID.006623.
157. Mohamed Q. M., Fahim A. Shaltout, f.A. and Ali, E.A. (2025): Bacteriological Quality Profiles and Prevalence of *Staphylococcus aureus*, *Salmonella* Species, and *E. coli* in Meat Samples of Sheep and Goats. *Egyptian Journal of Veterinary Sciences* pp 1-7. DOI: 10.21608/EJVS.2024.312380.2317
158. Ibrahim, S. M.; Hassanin, F. S.; Abou-Elroos, N. S. and Shaltout, F.A (2025): Quantifying The antimicrobial Efficacy of Selected Herbal Essential Oils Against Bacteria in Simulated Beef Steak Conditions. *Egyptian Journal of Veterinary Sciences*, pp 1-9. DOI: 10.21608/EJVS.2024.329367.2437
159. Shaltout, F. A (2024): The Availability, the Price, the Tradition, the Religion, the Income, the Social, the Development and the Economic Influences on the Meat Consumption. *Biomed J Sci & Tech Res* 55(4)-2024. BJSTR. MS.ID.008734.
160. Ibrahim, S. M.; Hassanin, F. S.; Abou-Elroos, N. S. and Shaltout, F.A (2025): Evaluating The impact of Certain Herbal Essential Oils on The Shelf Life and Chemical Composition of Beef Steak. *Egyptian Journal of Veterinary Sciences*, pp. 1-8. DOI: 10.21608/EJVS.2024.329509.2439
161. Shaltout, F. A. (2024): Our Opinion on Using of Irradiation in Food Preservation and Production. *Journal of Medical and Clinical Case Reports*, 1(6): 1-9. <https://doi.org/10.61615/JMCCR/2024/AUG027140805>
162. Anees, K. P; El-diasty, E. M. and Shaltout, F. A (2023): Mycological Evaluation and Occurrence of Aflatoxins and Ochratoxin A in *Tilapia Oreochromis niloticus* Fish and Fish Products. *Journal of Advanced Veterinary Research* ,13(7):1381-1385.
163. AMR, A. K; HASSANIN, F. S.; HASSAN, M. A. and SHALTOUT, F. A. E. (2024): TRIALS TO ESTIMATE AND CONTROL THE RESIDUAL LEVELS OF HETEROCYCLIC AROMATIC AMINES IN MEAT PRODUCTS. *Assiut Vet. Med. J.*, 70 (182): 98-105.
164. Shaltout, F. A Mohammed, I.; Afify, E. A (2020): Detection of *E. coli* O157 and *Salmonella* species in some raw chicken meat cuts in Ismailia province, Egypt. *Benha Veterinary Medical Journal* 39(2): 101-104.
165. Hassanin, F. S.; Shaltout, F. A.; Maarouf, A. A.; El-Sisy, S. F.; Ahmed, A. E. (2020): Bacteriological profile of frozen chicken meat cuts at Qalubiya governorate markets. *Benha Veterinary Medical Journal* 39 (2) 1-5.
166. Shaltout, F. A.; Heikal, G. I.; Ghanem, A. M. (2022): Mycological quality of some chicken meat cuts in Gharbiya governorate with special reference to *Aspergillus flavus* virulent factors. *Benha Veterinary Medical Journal* 40 (42) 12-16.
167. Shaltout, F. (2024) Application of Irradiation in Food Preservation and Production. *Journal of Pathology Research Reviews & Reports*. SRC/JPR-190. 6(5): 1-8. DOI: [doi.org/10.47363/JPR/2024\(6\)173](https://doi.org/10.47363/JPR/2024(6)173)
168. Taha, S. T.; Shaltout, F. A; Shimaa, N. Edris, S. N.; Mohamed, E. Nabil, M. E. (2024): Effect of lavender oil, clove oil and frankincense extract on sensory and microbial properties of raw drumsticks in refrigerator. *Benha Veterinary Medical Journal* 46 (1) 135-139.
169. Shaltout, F. A.; Salem, R. M; Eldiasty, E. M and Diab, F. A. (2023): Experimental Study on the Effect of Propionibacterium and Acetic acid on *Candida albicans* contamination in chicken fillet Stored at Chilling Conditions. *Benha Veterinary Medical Journal* 43 (2) 91-96.
170. Mubarak, S. R.; Abou EL-Roos, N. A.; Hussein, M. N. and Shaltout, F. A. E. (2024): Comparative microbiological evaluation between fresh and frozen bovine liver. *Benha Veterinary Medical Journal* 47 (1) 99-102.
171. El Asely, M. M. ; Fath Elbab, G. F.; Shaltout, F. A. E.(2024): Antibiotic Residues in Commercially Available Freshwater and Marine Fish: A Risk Assessment. *Egyptian Journal of Aquatic Biology & Fisheries*, 28(1): 397 – 410.
172. El Asely, M. M.; Fath Elbab, G. F. and Shaltout, A. E (2025): Impact of Freezing Intervals on Oxytetracycline and

- Ciprofloxacin Residues in Nile Tilapia and Catfish Muscles. Egypt. J. Vet. Sci. Vol. 56, No. 7, pp. 1419-1424.
173. Elkholy, R. A.; Abou EL-Roos, N. A.; Hussein, M. N. and Shaltout, F. A. E (2025): Differential Microbiological Quality on Marketed Frozen Turkey Breast and Thigh Meat. Egypt. J. Vet. Sci. 56 (1), pp. 1-10.
 174. Shaltout, F. A. (2024): THE FOOD ADDITIVES USED IN FOOD PRODUCTION, ADVANTAGES AND DISADVANTAGES. World Journal of Internal Medicine and Surgery 1 (6): 1-17
 175. Shaltout, F. A. (2024): Right Methods to Extend the Meat Shelf- Life by Using of Natural Preservatives and Their Public Health Importance. Journal of Medicine Care and Health Review 1(2): 1-17.
 176. Saad M. Saad, Fahim A. Shaltout, Amal A. A. Farag & Hashim F. Mohammed (2022): Organophosphorus Residues in Fish in Rural Areas. Journal of Progress in Engineering and Physical Science 1(1): 27-31.
 177. Shaltout, F. A. (2024): Importance of Extending the Shelf Life of the Meat. Journal of Medical and Clinical Case Reports 01 | (9): 1-10.
 178. Shaltout, F. A. E. Mona N. Hussein, Nada Kh. Elsayed (2023): Histological Detection of Unauthorized Herbal and Animal Contents in Some Meat Products. Journal of Advanced Veterinary Research (2023) 13(2): 157-160.
 179. Shaltout, F. A (2023): Abattoir and Bovine Tuberculosis as A Reemerging Foodborne Disease. Clinical Medical Reviews and Reports 6(1): 1-7
 180. Shaltout, F. A. Ramadan M. Salem, Eman M. Eldiasty, Fatma A. Diab (2022): Seasonal Impact on the Prevalence of Yeast Contamination of Chicken Meat Products and Edible Giblets. Journal of Advanced Veterinary Research ,12(5):641-644.
 181. Shaltout, S. and Shaltout, F. (2024), "Food Borne Bacterial Diseases Due to Consumption of Meat, Fish and Poultry Products", Arch Health Sci; 8(1): 1-8.
 182. Shaltout, F. A (2024): Our Opinion on Using of Irradiation in Food Preservation and Production. Journal of Medical and Clinical Case Reports 01 | (6): 1-9
 183. Hasan, L.; Vgeli, P.; Neuenschwander, S.; Stoll, P.; Stranzinger, G. The L-gulonogamma-lactone oxidase gene (GULO) which is a candidate for vitamin C deficiency in pigs maps to chromosome 14. Anim. Genet. 2015, 30, 309–312
 184. Hakeem, K. P.; El-diasty, E. M.; Shaltout, F. A. E (2023): Effects of natural compounds of some plants on microbial contamination and sensory quality of fish fillet during refrigeration. Benha Veterinary Medical Journal 45 (1) 152-156.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here: [Submit Manuscript](#)

DOI: [10.31579/2693-2156/139](https://doi.org/10.31579/2693-2156/139)

Ready to submit your research? Choose Auctores and benefit from:

- ❖ fast, convenient online submission
- ❖ rigorous peer review by experienced research in your field
- ❖ rapid publication on acceptance
- ❖ authors retain copyrights
- ❖ unique DOI for all articles
- ❖ immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <http://www.auctoresonline.org/journals/journal-of-thoracic-disease-and-cardiothoracic-surgery>